
(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference 150 . 192W01	FOR FURTHER see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.				
International application No.	International filing date (day/month/year)	(Earliest) Priority Date (day/month/year)			
PCT/US 00/14296	24/05/2000	24/05/1999			
Applicant MAYO MEDICAL VENTURES					
according to Article 18. A copy is being transfer and the second	_				
	nternational search was carried out on the ba less otherwise indicated under this item.	sis of the international application in the			
the international search was Authority (Rule 23.1(b)).	as carried out on the basis of a translation of	the international application furnished to this			
b. With regard to any nucleotide and was carried out on the basis of the Contained in the internation filed together with the internation furnished subsequently to the statement that the subsinternational application as	sequence listing: nal application in written form. national application in computer readable for this Authority in written form. this Authority in computer readble form. sequently furnished written sequence listing of filed has been furnished.				
Certain claims were foun Unity of invention is lack	d unsearchable (See Box I). Ing (see Box II).				
	ed by this Authority to read as follows:	DURING XENOTRANSPLANTATION			
	• • • • • • • • • • • • • • • • • • • •	ity as it appears in Box III. The applicant may, port, submit comments to this Authority.			
6. The figure of the drawings to be public as suggested by the applic because the applicant faile	shed with the abstract is Figure No.	None of the figures.			



CLASSIFICATION OF SUBJECT MATTER
PC 7 C12N15/48 C12N15/62 A. CLAS C07K14/15 C07K19/00 A61K48/00 C1201/68 C07K16/10 C12N7/01 According to International Patent Classification (IPC) or to both national classification and IPC **B. FIELDS SEARCHED** Minimum documentation searched (classification system followed by classification symbols) C12N C07K A61K C12Q IPC 7 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practical, search terms used) BIOSIS, CHEM ABS Data C. DOCUMENTS CONSIDERED TO BE RELEVANT Category ° Citation of document, with indication, where appropriate, of the relevant passages Relevant to claim No. X AKIYOSHI D. E. ET AL.: "Identification of 15-20 a full-length cDNA for an endogenous retrovirus of miniature swine" JOURNAL OF VIROLOGY, vol. 72, no. 5, May 1998 (1998-05), pages 4503-4507, XP002126425 ISSN: 0022-538X Υ the whole document 1-14.21-28. 30,32 - 39Further documents are listed in the continuation of box C. Patent family members are listed in annex. Х X Special categories of cited documents: "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international "X" document of particular relevance; the claimed invention filing date cannot be considered novel or cannot be considered to "L" document which may throw doubts on priority claim(s) or involve an inventive step when the document is taken alone which is cited to establish the publication date of another "Y" document of particular relevance; the claimed invention citation or other special reason (as specified) cannot be considered to involve an inventive step when the document is combined with one or more other such docu-"O" document referring to an oral disclosure, use, exhibition or other means ments, such combination being obvious to a person skilled document published prior to the international filing date but later than the priority date claimed "&" document member of the same patent family Date of the actual completion of the international search Date of mailing of the international search report 21 September 2000 04/10/2000 Name and mailing address of the ISA Authorized officer European Patent Office, P.B. 5818 Patentlaan 2 NL – 2280 HV Rijswijk Tel. (+31–70) 340–2040, Tx. 31 651 epo nl, Fax: (+31–70) 340–3016

5

Mandl, B

Interpretional Application No PC 5 00/14296

		PC 5 00/14296
	ation) DOCUMENTS CONSIDERED TO BE RELEVANT Citation of decument, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Gategory *	— Спанса от десентем, мин телезавен, винете груперияте, от ще телезавен разхадез	
Υ	VANBROCKLIN M. ET AL.: "Expression of a murine leukemia virus Gag-Escherichia coli RNase HI fusion polyprotein significantly inhibits virus spread." JOURNAL OF VIROLOGY, vol. 71, no. 4, 1997, pages 3312-3318, XP002148090 ISSN: 0022-538X the whole document	1-14, 21-28, 32-39
X	WO 98 53104 A (MEDICAL RES COUNCIL ;STOYE JONATHAN PAUL (GB); WEISS ROBIN ANTHONY) 26 November 1998 (1998-11-26)	29,31
Υ	the whole document	30
A	NATSOULIS G. ET AL.: "NEW ANTIVIRAL STRATEGY USING CAPSID-NUCLEASE FUSION PROTEINS" NATURE, vol. 352, no. 6336, 15 August 1991 (1991-08-15), pages 632-635, XP000608400 ISSN: 0028-0836 the whole document	15-28
A	SUN LQ. ET AL.: "Ribozyme-mediated suppression of Moloney murine leukemia virus and human immunodeficiency virus type I replication in permissive cells." PROC. NATL. ACAD. SCI. USA, vol. 91, October 1994 (1994-10), pages 9715-9719, XP002148091 the whole document	13
Α	WO 96 32494 A (UNIV ALABAMA RES FOUND) 17 October 1996 (1996-10-17) the whole document	15-28
A	PATIENCE C. ET AL.: "INFECTION OF HUMAN CELLS BY AN ENDOGENOUS RETROVIRUS OF PIGS" NATURE MEDICINE, vol. 3, no. 3, 1 March 1997 (1997-03-01), pages 282-286, XP002037074 ISSN: 1078-8956 the whole document	1-39

5

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-28,32-39

A method to inhibit or prevent infectious agent transmission in a mammalian transplant recipient; an isolated and purified nucleic acid molecule comprising at least a portion of a pig endogenous retrovirus, the corresponding polypeptides and methods of using them; a method to detect human tropic pig endogenous retroviruses; a fusion protein comprising a capsid or envelope protein of PERV and a degradative enzyme, the corresponding nucleic acid and a host cell comprising it.

2. Claims: 29-31

An antibody that specifically bings pig endogenous retrovirus.

nform on patent family members

Interactional Application No PC 5 00/14296

Patent document cited in search report	rt	Publication date		atent family member(s)	Publication date
WO 9853104	A	26-11-1998	AU EP GB US	7442698 A 0981651 A 2333102 A 6100034 A	11-12-1998 01-03-2000 14-07-1999 08-08-2000
WO 9632494	Α	17-10-1996	AU CA EP US	5482896 A 2230925 A 0873415 A 6001985 A	30-10-1996 17-10-1996 28-10-1998 14-12-1999



PCT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

Applicant's	or and	ent's file reference							
, , ,		FOR FURTHER AC			tion of Transmittal of Internation Examination Report (Form PC				
		ication No	International filing date (a	lav/month/ve	par)	Priority date (day/month/year			
1			24/05/2000	аулпопплуе	zar,	24/05/1999	,		
			↓			2 17 0 07 1 0 0 0			
	International Patent Classification (IPC) or national classification and IPC C12N15/48								
Applicant	Applicant								
''	MAYO FOUNDATION FOR MEDICAL EDUCATION et al.								
IVIATOR	OUN	DATION FOR MEDIC	AL EDUCATION et	aı.					
				orepared b	y this Inter	national Preliminary Exam	ining Authority		
and is	trans	smitted to the applicant a	according to Article 36.						
2. This f	REPO	RT consists of a total of	8 sheets, including this	cover she	et.				
п	hie ro	nort is also accompanio	d hy ANNEXES in sho	ate of the	description	, claims and/or drawings w	vhich have		
						ctifications made before this			
(5	see R	ule 70.16 and Section 6	07 of the Administrative	Instruction	s under the	PCT).			
These	e anne	exes consist of a total of	sheets.						
3. This r	eport	contains indications rela	ating to the following item	ns:					
[\boxtimes	Basis of the report		•					
11		Priority							
iii		•	pinion with regard to nov	velty, inver	ntive step a	and industrial applicability			
IV	\boxtimes	Lack of unity of invention	· -	• •	•				
V	\boxtimes				velty, inve	ntive step or industrial appl	licability;		
VI		Certain documents cite	ons suporting such state	ment	•				
VII		Certain defects in the in							
VIII	\boxtimes		n the international applic	ation					
					, -				
Data of out	minolo	on of the demand		Data of oar	npletion of t	hio roport			
Date of Suc	ninssic	on or the demand		Date of cor	npietion or t	nis report			
13/12/20	00			10.07.2001	1				
		g address of the internationa	d .	Authorized	officer		AND LECKES MATOURS		
premimary		ning authority: pean Patent Office							
)	D-80	0298 Munich	S opmud	Roscoe,	R		Salar		
Tel. +49 89 2399 - 0 Tx: 523656 epmu d Fax: +49 89 2399 - 4465			э ерни и	Telephone	No. +49 89	2399 2554	BOWN STORES WHOLE ELWON		

Telephone No. +49 89 2399 2554



INTERNATIONAL PRELIMINARY

EXAMINATION REPORT

International application No. PCT/US00/14296

1.	With regard to the elements of the international application (Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report since they do not contain amendments (Rules 70.16 and 70.17)): Description, pages:							
	1-83	3	as originally filed					
	Clai	ms, No.:						
	1-45	5	as received on ,	02/01/2001	with letter of	22/12/2000		
	Dra	wings, sheets:	r					
	1/21	-21/21	as originally filed					
	Seq	uence listing part	of the description, pages:					
	1-30), as originally filed						
						A contact of A contact of the Atlanta		
2.	With	n regard to the lang Juage in which the i	ruage, all the elements marke international application was f	d above were a led, unless oth	erwise indicated unde	er this item.		
	These elements were available or furnished to this Authority in the following language: , which is:							
		the language of a	translation furnished for the po	urposes of the i	international search (ı	under Rule 23.1(b)).		
		the language of pu	blication of the international a	pplication (und	er Rule 48.3(b)).			
	the language of a translation furnished for the purposes of international preliminary examination (under Rule 55.2 and/or 55.3).							
3.	. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:							
	\boxtimes	contained in the in	ternational application in writte	en form.				
	\boxtimes	filed together with	the international application in	computer read	dable form.			
		furnished subsequ	ently to this Authority in writte	n form.				
		furnished subsequ	ently to this Authority in comp	uter readable f	orm.			
			t the subsequently furnished v pplication as filed has been fu		ce listing does not go	beyond the disclosure in		
		The statement tha listing has been fu	t the information recorded in ornished.	omputer reada	ble form is identical to	the written sequence		
4	TL -		regulted in the cancellation o	٤.				



INTERNATIONAL PRELIMINARY

EXAMINATION REPORT

International application No. PCT/US00/14296

		the description,	pages:				
		the claims,	Nos.:				
		the drawings,	sheets:				
5.			established as if (some of) the amendments had not been made, since they have been yond the disclosure as filed (Rule 70.2(c)):				
		(Any replacement sh report.)	neet containing such amendments must be referred to under item 1 and annexed to this				
6.	Add	litional observations, i	f necessary:				
			,				
IV.	Lac	k of unity of inventi	on				
1.	In re	esponse to the invitati	on to restrict or pay additional fees the applicant has:				
		restricted the claims.					
		paid additional fees.					
		paid additional fees	under protest.				
		neither restricted nor	paid additional fees.				
2.			that the requirement of unity of invention is not complied and chose, according to Rule applicant to restrict or pay additional fees.				
3.	This Authority considers that the requirement of unity of invention in accordance with Rules 13.1, 13.2 and 13.3 is						
		complied with.					
		not complied with for	the following reasons:				
4.		sequently, the followi mination in establishir	ng parts of the international application were the subject of international preliminary ng this report:				
	\boxtimes	all parts.					
		the parts relating to c	slaims Nos				
	cita		der Article 35(2) with regard to novelty, inventive step or industrial applicability; ons supporting such statement				
	Nov	relty (N)	Yes: Claims 1-14, 16-28, 32-45				





EXAMINATION REPORT

International application No. PCT/US00/14296

No: Claims 15, 29-31

Inventive step (IS) Yes: Claims

No: Claims 1-45

Industrial applicability (IA) Yes: Claims 15, 16, 20-31, 40, 41, 44

No: Claims 1-14, 17-19, 32-39, 42, 43, 45

2. Citations and explanations see separate sheet

VIII. Certain observations on the international application

The following observations on the clarity of the claims, description, and drawings or on the question whether the claims are fully supported by the description, are made: see separate sheet



The documents mentioned in the present International Preliminary Examination Report are numbered as in the search report, i.e. D1 corresponds to the first document of the search report etc.

Lack of Unity IV.

The present application lacks unity. The only concept that links detection assays for PERVs and related products (i.e. antibodies) to methods of prevention of PERV (or other infectious agent) transmission from transplant tissues to recipients by introduction of transgenes into the donor cells is PERV. However PERVs were already known in the art (e.g. D1, D3 or D7). Hence, common concept is not even novel. For practical reasons this objection is not persued in the International Phase.

Reasoned statement on Novelty, Inventive Step and Industrial Applicability V.

Novelty (Art.33(2) PCT)

Claim 15 - anticipated by nucleic acids disclosed in D1, D3 and D7.

Claims 29-31 are anticipated by D3 which suggests making antibody probes for detecting PERV expression (particularly direct probes to N-terminus of env).

Inventive Step (Art.33(3) PCT)

Matter not directly related to inhibition of PERV transfer from xenografts:

Claims 15-17 contain no inventive subject-matter. PERV sequences had already been identified and identification of further sequences by use of probes (see e.g. D1 or D7) is simple (due to extremely high levels of sequence conservation). Reuse of any such further sequences as probes is merely a trivial alternative which does not solve any new problem vis a vis the prior art probes.

Claims 18 and 19 merely relate to transformation of PERV sequences into host cells. It is trivial to insert a known DNA into a cell unless some unexpected

technical effect can be achieved thereby. That is not the case in claims 18 and 19 since proteins not manipulated to be of e.g. medical utility. Hence, claims 18 and 19 are not inventive.

Claims 29-31 contain no inventive subject-matter. Raising antibodies to known proteins is trivial especially in view of possible probe utility (e.g. D3)

Matter relating to inhibition of PERV transfer from xenografts:

(i) Fusion proteins

The construction of fusion proteins between retroviral proteins and virus inhibitory proteins has been disclosed in the prior art (see D2, D4 or D6). Such fusion proteins could find use for treatment of non-transplant-associated retroviral infection aswell as in applicants proposed xenograft context. Mere application of this technology to an alternative retroviral system of viral system in general cannot be considered inventive. It is at best only the xenograft context which could be considered to impart inventivity and hence, given that product claims per se cannot be limited to this context, mere provision of the fusion proteins of claims 21-26 and systems for expression thereof in claims 27 and 28 cannot be considered inventive.

Methods for inhibiting infectious agent transmission (ii)

The cited prior art does not suggest the use of transplant donor cells which have been rendered transgenic in an attempt to prevent transmission of infective agents therefrom. D7 is the closest prior art identified in this context. D7 recognizes the problem of PERV transmission to humans during organ transplantation. As a remedy, removal of PERV sequences from the pig genome is suggested (p.285, col.1). Further, (presumably until this has happened) it is stated that precautions should be taken to minimize such zoonosis and to monitor retrovirus infection in early clinical trials. It is not stated how this problem should be addressed. Given this statement, a skilled person is presented with a limited choice of options. These are basically continuous administration of antiretroviral medicaments, or integration of an antiretroviral transgene construct into donor or



recipient tissues (or both). Applicant has chosen to make the donor organ transgenic (the technology used is defined very broadly, and it is noted that no new techniques as such have been introduced). Hence, applicant has made a selection from a very limited number of options to deal with a recognized problem. No surprising effect has been shown, and if it had been this would be limited to a particular method, not any method of donor-targetted transgene based antiretroviral treatment. It is noted that the concept of donor organ genetic manipulation in general is not a novel one. WO-A-94/16065 discloses donor cells with adenoviral E19 protein (reduces MHC I presentation) expressing transgenes. WO-A-95/28412 discloses transgenic swine as organ donors where alpha(1,3)galactosyltransferase expression is prevented by an antisense transgene.

The present application is thus considered devoid of inventive subject-matter.

Industrial Applicability (Art.33(4) PCT)

For the assessment of the present claims 1-14, 17-19, 32-39, 42, 43 and 45 on the question whether they are industrially applicable, no unified criteria exist in the PCT Contracting States. The patentability can also be dependent upon the formulation of the claims. The EPO, for example, does not recognize as industrially applicable the subject-matter of claims to the use of a compound in medical treatment, but may allow, however, claims to a known compound for first use in medical treatment and the use of such a compound for the manufacture of a medicament for a new medical treatment.

Claims 1-14, 17-19, 32-39, 42, 43 and 45 relate to subject-matter considered by this Authority to be covered by the provisions of Rule 67.1(iv) PCT. Consequently. no opinion will be formulated with respect to the industrial applicability of the subject-matter of these claims (Article 34(4)(a)(i) PCT).

VIII. Certain observations

Clarity (Art.6 PCT)

The number of independent method claims relating to methods of inhibiting infectious agent transmission is excessive and unacceptable (i.e. claims 1, 2, 3, 36, 37, 40, 42).

Claims 1-4 - size of portion not defined so can be any protein. Any protein would not solve problem

Claim 15 - size of portion / hybridizing conditions not defined

Claim 16 "comprising a nucleic acid segment" is unnecessary and renders wording confusing. Further, given that nucleic acid defined in open-ended manner can contain reading frame of any known protein, thus claim effectively lacks novelty over all proteins. Further actual proteins encoded by nucleic acids of Seq.ID nos 18-23 and 32 not identified anywhere in application or claims.

Claim 18 - method makes no sense - seem simply to transform a cell with

Claims 20, 37, 41, 42, 44, 45 - size of portion

Note: It has been noted that claim 41 encompasses transgenic human beings. Such matter would not be admissible in a European Regional Phase.





For receiving Office use only International Application No. REQUEST International Filing Date The undersigned requests that the present international application be processed Name of receiving Office and "PCT International Application" according to the Patent Cooperation Treaty. Applicant's or agent's file reference (if desired) (12 characters maximum) 150.192WO1 Box No. I TITLE OF INVENTION METHODS TO INHIBIT INFECTIOUS AGENT TRANSMISSION **APPLICANT** Box No. II Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is, country) of residence is indicated by the law. This person is also inventor. of residence is indicated below.) MAYO FOUNDATION FOR MEDICAL EDUCATION AND RESEARCH Telephone No. 200 First Street SW Rochester, Minnesota 55905 Facsimile No. United States of America Teleprinter No. State (that is, country) of nationality: State (that is, country) of residence: US This person is applicant all designated States all designated States except the United States of America the United States the States indicated in for the purposes of: of America only the Supplemental Box FURTHER APPLICANT(S) AND/OR (FURTHER) INVENTOR(S) Box No. III Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is, country) of residence if no State of residence is indicated below.) This person is: applicant only FEDERSPIEL, Mark J. 5722 Colonial Lane SE applicant and inventor Rochester, Minnesota 55904 inventor only (If this check-box United States of America is marked, do not fill in below.) State (that is, country) of nationality: State (that is, country) of residence: This person is applicant all designated States except the United States of America all designated the United States of America only the States indicated in for the purposes of: States the Supplemental Box Further applicants and/or (further) inventors are indicated on a continuation sheet. Box No. IV AGENT OR COMMON REPRESENTATIVE; OR ADDRESS FOR CORRESPONDENCE The person identified below is hereby/has been appointed to act on behalf of the applicant(s) before the competent International Authorities as: X agent common representative Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country.) Telephone No. VIKSNINS, Ann S. (612) 373-6900 Schwegman, Lundberg, Woessner & Kluth Facsimile No. P.O. Box 2938 Minneapolis, Minnesota 55402 (612) 339-3061 United States of America Teleprinter No. Address for correspondence: Mark this check-box where no agent or common representative is/has been appointed and the space above is used instead to indicate a special address to which correspondence should be sent.

***	W. A.						
PCT	For receiving Office use only						
	International Application No.						
REQUEST							
ALL QUEST	International Filing Date						
The undersigned requests that the present							
international application be processed according to the Patent Cooperation Treaty.	Name of receiving Office and "PCT International Application"						
	Applicant's or agent's file reference (if desired) (12 characters maximum) 150.192WO1						
Box No. 1 TITLE OF INVENTION	l						
METHODS TO INHIBIT INFECTIOUS AGENT TRANS	MISSION						
Box No. II APPLICANT							
Name and address: (Family name' followed by given name; for a designation. The address must include postal code and name of cou address indicated in this Box is the applicant's State (that is, country of residence is indicated below.)	legal entity, full official miry. The country of the o) of residence if no State This person is also inventor.	-					
MAYO MEDICAL VENTURES	Telephone No.						
200 First Street SW Rochester, Minnesota 55905							
United States of America	Facsimile No.						
·							
	Teleprinter No.						
State (that is, country) of nationality:	State (that is, country) of residence:						
US	US						
This person is applicant for the purposes of: all designated the United States All designated the United States	d States except ates of America the United States the States indicated in the States indicated in the Supplemental Bo						
Box No. III FURTHER APPLICANT(S) AND/OR (FURTH							
Name and address: (Family name followed by given name; for a designation. The address must include postal code and name of counaddress indicated in this Box is the applicant's State (that is, country, of residence is indicated below).	egal entity, full official						
address indicated in this Box is the applicant's State (that is, country, of residence is indicated below.)							
FEDERSPIEL, Mark J.	applicant only						
5722 Colonial Lane SE	x applicant and inventor						
Rochester, Minnesota 55904	inventor only Afthir sheek have						
United States of America	inventor only (If this check-box is marked, do not fill in below.)						
State (that is, country) of nationality:	I Shared at the state of the st						
US	State (that is, country) of residence: US						
This person is applicant all designated the United States all designated the United States	States except ates of America the United States of America only the States indicated in the Supplemental Box	n X					
Further applicants and/or (further) inventors are indicated on a continuation sheet.							
Box No. IV AGENT OR COMMON REPRESENTATIVE;							
The person identified below is hereby/has been appointed to act or of the applicant(s) before the competent International Authorities a	s: agent common representative						
Name and address: (Family name followed by given name; for a designation. The address must include postal code	legal entity, full official Telephone No.						
VIKSNINS, Ann S. Schwegman, Lundberg, Woessner & Kluth	(612) 373-6900						
P.O. Box 2938	Facsimile No.						
Minneapolis, Minnesota 55402 United States of America	(612) 339-3061						
O-ution of thinories	Teleprinter No.	_					

Address for correspondence: Mark this check-box where no agent or common representative is/has been appointed and the space above is used instead to indicate a special address to which correspondence should be sent.

Form PCT/RO/101 (first sheet) (July 1998; reprint January 2000)

See Notes to the request

See Notes to the request form



 _

	Box No.V DESIGNATION OF STATES									
1	he fo	llowing designations are hereby made under Rule 4.9(a)	mar	k the o	unlicable check-hores; at least one must be marked).					
		nal Patent			pproduct encer boxes, at least one must be marked).					
7		AP ARIPO Patent: GH Ghana, GM Gambia, KE Kenya, LS Lesotho, MW Malawi, SD Sudan, SL Sierra Leone, SZ Swaziland TZ United Republic of Tanzania, UG Uganda, ZW Zimbabwe, and any other State which is a Contracting State of the Harar Protocol and of the PCT								
[K EA	Eurasian Patent: AM Armenia, AZ Azerbaijan, BY Belarus, KG Kyrgyzstan, KZ Kazakhstan, MD Republic of Moldova RU Russian Federation, TJ Tajikistan, TM Turkmenistan, and any other State which is a Contracting State of the Eurasian Patent Convention and of the PCT								
[2	E EP	European Patent: AT Austria, BE Belgium, CH and LI Switzerland and Liechtenstein, CY Cyprus, DE Germany DK Denmark, ES Spain, FI Finland, FR France, GB United Kingdom, GR Greece, IE Ireland, IT Italy, LU Luxembourg MC Monaco, NL Netherlands, PT Portugal, SE Sweden, and any other State which is a Contracting State of the European Patent Convention and of the PCT								
[2	☑ OA	OAPI Patent: BF Burkina Faso, BJ Benin, CF Central African Republic, CG Congo, CI Côte d'Ivoire, CM Cameroon GA Gabon, GN Guinea, GW Guinea-Bissau, ML Mali, MR Mauritania, NE Niger, SN Senegal, TD Chad, TG Togo, and any other State which is a member State of OAPI and a Contracting State of the PCT (if other kind of protection or treatment desired specify on dotted line)								
N	lation	al Patent (if other kind of protection or treatment desired, spe	 c.		α. J.τ., Α.					
5	AE	United Arab Emirates	<i>cijy</i> —	on aoi	iea ine).					
		Albania	=	_	Liberia					
		I Armenia			Lesotho					
6	TAS	Austria	_	LT	Lithuania					
		Australia	_		0					
		Azerbaijan			Latvia					
					Morocco					
		Bosnia and Herzegovina	X	MD	Republic of Moldova					
	_	Barbados	X	MG	Madagascar					
I.S.		Bulgaria	X] MK	The former Yugoslav Republic of Macedonia					
Į.		Brazil								
		Belarus	X	MN	Mongolia					
_	_	Canada			' Malawi					
		and LI Switzerland and Liechtenstein	X	MX	Mexico					
X	CN	China	X	NO	Norway					
	CK	Costa Rica	X	NZ	New Zealand					
	CU	Cuba		PL	Poland					
K	CZ	Czech Republic	X	PT	Portugal					
	DE	Germany	X	RO	Romania					
		Denmark	X	RU	Russian Federation					
		Dominica	X	SD	Sudan					
		Estonia	X	SE	Sweden					
	ES	Spain	X	SG	Singapore					
	FI	Finland	X	SI	Slovenia					
		United Kingdom	X	SK	Slovakia					
		Grenada		SL	Sierra Leone					
X	GE	Georgia		TJ	Tajikistan					
K	GH	Ghana	X	TM	Turkmenistan					
		Gambia	X	TR	Turkey					
		Croatia	X	TT	Trinidad and Tobago					
	HU	Hungary	X	TZ	United Republic of Tanzania					
=	ID	Indonesia		UA	Ukraine					
	IL	Israel		UG	Uganda					
	IN	India	X	US	United States of America					
	IS	Iceland								
	JP	Japan	×	UZ	Uzbekistan					
		Kenya	X	VN	Viet Nam					
_		Kyrgyzstan	X	YU	Yugoslavia					
X	KP	Democratic People's Republic of Korea	X	ZA	South Africa					
				$\mathbf{z}\mathbf{w}$	Zimbabwe					
		Republic of Korea	Ch	eck-b	oxes reserved for designating States which have					
X	ΚZ	Kazakhstan	bec	ome p	party to the PC1 after issuance of this sheet:					
X	LC	Saint Lucia	X	AG A	ntigua and Barbuda					
			DZ Algeria							
Pre	caut	onary Designation Statement: In addition to the designa	tion	s mad	e above, the applicant also makes under Rule 4.9(b) all other					
des	ignati	ons which would be permitted under the PCT except any	des	ignatio	on(s) indicated in the Supplemental Box as being excluded					







Box No. VI PRIORITY C	LAIM	Further prior	Further priority claims are indicated in the Supplemental Box.			
Filing date Number of earlier application of earlier application			Where earlier application is:			
of earlier application (day/month/year)	or carner application	national application:		international application:		
item (1) (24.05.99)		country	regional Office	receiving Office		
24 May 1999	60/135,631	us				
item (2)						
item (3)						
(5)						
The receiving Office is						
purposes of the present into) (only if the earlier a _l ernational application	ransmit to the International Bur pplication was filed with the t is the receiving Office) identifi	Office which for the ed above as item(s): ()			
* Where the earlier application is a Convention for the Protection of Inc	an ARIPO application, it i dustrial Property for whic	is mandatory to indicate in the Suj ch that earlier application was file	pplemental Box at least one of Rule 4 10(h)(ii)) See Su	e country party to the Paris		
Box No. VII INTERNATIO	NAL SEARCHING A	AUTHORITY	(Itale 4.10(0)(11)). Bee Bu	рриетета Вох.		
Choice of International Search (if two or more International Sea competent to carry out the interna	rching Authorities are	Request to use results of ear search has been carried out by or	lier search; reference trequested from the Internati	to that search (if an earlier ional Searching Authority):		
the Authority chosen; the two-letter of	code may be used):	Date (day/month/year)	Number	Country (or regional Office)		
ISA / EP						
Box No. VIII CHECK LIST			·			
This international application co the following number of sheets	:	tional application is accompan	ied by the item(s) marke	d below:		
request : 3		alculation sheet				
description (excluding sequence listing part) : 83		ate signed power of attorney of general power of attorney;	rafaranca numbar if and			
claims : 5		nent explaining lack of signatu				
abstract : 1	j .	ty document(s) identified in Bo		į		
drawings : 21	1	ation of international application				
sequence listing part of description : 30	7. 🔲 separa	ate indications concerning depo	sited microorganism or	other biological material		
	8. 🗷 nucleo	otide and/or amino acid sequen	ce listing in computer re	eadable form		
Total number of sheets: 143		(specify): general transmittal; ch	eck; return postcard; Comm	n. re: sequence lisitng		
Figure of the drawings which should accompany the abstract:	0	T T	nglish			
	OF APPLICANT OR					
Next to each signature, indicate the name of the person signing and the capacity in which the person signs (if such capacity is not obvious from reading the request). Ann S. Viksnins						
Date of actual receipt of the printernational application:		r receiving Office use only —	\	2. Drawings:		
 Corrected date of actual receitimely received papers or drathe purported international ap 	wings completing	`		received:		
Date of timely receipt of the recorrections under PCT Article	e [1(2):			not received:		
5. International Searching Author (if two or more are competent	ority S): ISA/	6. Transmittal until search	of search copy delayed fee is paid.			
Date of receipt of the record copy by the International Bureau use only						

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference 150.192W01	FOR FURTHER see Notification (Form PCT/ISA/2	of Transmittal of International Search Report 220) as well as, where applicable, item 5 below.
International application No.	International filing date (day/month/year)	(Earliest) Priority Date (day/month/year)
PCT/US 00/14296	24/05/2000	24/05/1999
MAYO MEDICAL VENTURES		
This International Search Report has be according to Article 18. A copy is being t	en prepared by this International Searching Auti ransmitted to the International Bureau.	nority and is transmitted to the applicant
	s of a total of <u>5</u> sheets. y a copy of each prior art document cited in this	report.
· ·	. Tada a santa a	
language in which it was filed, ur	e international search was carried out on the bas pless otherwise indicated under this item.	is of the international application In the
the international search (Authority (Rule 23.1(b)).	was carried out on the basis of a translation of the	ne international application furnished to this $^{\frac{1}{4}}$
The second of the		ternational application, the international search
	onal application in written form. ernational application in computer readable form	
	o this Authority in written form.	1.
	this Authority in computer readble form.	
the statement that the su	bsequently furnished written sequence listing do as filed has been furnished.	pes not go beyond the disclosure in the
		identical to the written sequence listing has been
	nd unsearchable (See Box I).	
3. X Unity of invention is lac	king (see Box II).	
4. With regard to the title,	,	
the text is approved as su	hmitted by the emplicent	
	hed by this Authority to read as follows:	
METHODS TO INHIBIT IN	FECTIOUS AGENT TRANSMISSION	DURING XENOTRANSPLANTATION
	,	
5. With regard to the abstract,		•
the text is approved as su	V hmitted by the applicant	
the text has been establis	hed, according to Rule 38.2(b), by this Authority date of mailing of this international search repo	as it appears in Box III. The applicant may,
6. The figure of the drawings to be publ		, and realisting.
as suggested by the appli		X None of the figures.
because the applicant fallo	ed to suggest a figure.	in the lightes.
because this figure better	characterizes the invention.	

Application No 00/14296 A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C12N15/48 C12N15/62 C07K14/15 C07K19/00 A61K48/00 C12Q1/68 C07K16/10 C12N7/01 According to International Patent Classification (IPC) or to both national classification and IPC **B. FIELDS SEARCHED** Minimum documentation searched (classification system followed by classification symbols) IPC 7 C12N C07K A61K C120 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practical, search terms used) BIOSIS, CHEM ABS Data C. DOCUMENTS CONSIDERED TO BE RELEVANT Category ° Citation of document, with indication, where appropriate, of the relevant passages Relevant to claim No. X AKIYOSHI D. E. ET AL.: "Identification of 15-20 a full-length cDNA for an endogenous retrovirus of miniature swine" JOURNAL OF VIROLOGY, vol. 72, no. 5, May 1998 (1998-05), pages 4503-4507, XP002126425 ISSN: 0022-538X Y the whole document 1-14. 21-28, 30,32-39 Further documents are listed in the continuation of box C. X X Patent family members are listed in annex. Special categories of cited documents: *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the "A" document defining the general state of the art which is not considered to be of particular relevance invention earlier document but published on or after the international filing date

document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

document published prior to the international filing date but later than the priority date claimed

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

21 September 2000


Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentiaan 2 NL – 2280 HV Rijswijk Tel. (+31–70) 340–2040, Tx. 31 651 epo ni, Fax: (+31–70) 340–3016

Date of mailing of the international search report

04/10/2000

Authorized officer

Mandl, B



C.(Continu	nation) DOCUMENTS CONSIDERED TO BE RELEVANT	101, 00/14296
Category °		Relevant to claim No.
Υ	VANBROCKLIN M. ET AL.: "Expression of a murine leukemia virus Gag-Escherichia coli RNase HI fusion polyprotein significantly inhibits virus spread." JOURNAL OF VIROLOGY, vol. 71, no. 4, 1997, pages 3312-3318, XP002148090 ISSN: 0022-538X the whole document	1-14, 21-28, 32-39
X	WO 98 53104 A (MEDICAL RES COUNCIL ;STOYE JONATHAN PAUL (GB); WEISS ROBIN ANTHONY) 26 November 1998 (1998-11-26) the whole document	29,31
A	NATSOULIS G. ET AL.: "NEW ANTIVIRAL STRATEGY USING CAPSID-NUCLEASE FUSION PROTEINS" NATURE, vol. 352, no. 6336, 15 August 1991 (1991-08-15), pages 632-635, XP000608400 ISSN: 0028-0836 the whole document	15-28
A	SUN LQ. ET AL.: "Ribozyme-mediated suppression of Moloney murine leukemia virus and human immunodeficiency virus type I replication in permissive cells." PROC. NATL. ACAD. SCI. USA, vol. 91, October 1994 (1994-10), pages 9715-9719, XP002148091 the whole document	13
A	WO 96 32494 A (UNIV ALABAMA RES FOUND) 17 October 1996 (1996-10-17) the whole document	15-28
A	PATIENCE C. ET AL.: "INFECTION OF HUMAN CELLS BY AN ENDOGENOUS RETROVIRUS OF PIGS" NATURE MEDICINE, vol. 3, no. 3, 1 March 1997 (1997-03-01), pages 282-286, XP002037074 ISSN: 1078-8956 the whole document	1-39





FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-28,32-39

A method to inhibit or prevent infectious agent transmission in a mammalian transplant recipient; an isolated and purified nucleic acid molecule comprising at least a portion of a pig endogenous retrovirus, the corresponding polypeptides and methods of using them; a method to detect human tropic pig endogenous retroviruses; a fusion protein comprising a capsid or envelope protein of PERV and a degradative enzyme, the corresponding nucleic acid and a host cell comprising it.

2. Claims: 29-31

An antibody that specifically bings pig endogenous retrovirus.





Box I	Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)
This Into	ternational Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:
1. X	Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely:
	Although claims 1-14, 19 and 32-39 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2.	Claims Nos.: because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
ı	!
з. 🗌	Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).
Box II	Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)
This Inte	ernational Searching Authority found multiple Inventions in this international application, as follows:
	see additional sheet
1.	As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. X	As all searchable claims could be searched without effort justifying an additional fee, this Authority did not Invite payment of any additional fee.
3	As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. [] <u> </u>	No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
n-mark s	
Remark o	The additional search fees were accompanied by the applicant's protest. No protest accompanied the payment of additional search fees.

patent family members

Int	Application No
PCT	00/14296

Patent document		Publication Patent family		Patent family	y Publication	
cited in search repo	π	date mer		member(s)	date	
WO 9853104	A	26-11-1998	AU EP GB US	7442698 A 0981651 A 2333102 A 6100034 A	11-12-1998 01-03-2000 14-07-1999 08-08-2000	
WO 9632494	A	17-10-1996	AU CA EP US	5482896 A 2230925 A 0873415 A 6001985 A	30-10-1996 17-10-1996 28-10-1998 14-12-1999	

CORRECTED VERSION

9/9805 2900

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date 30 November 2000 (30.11.2000)

PCT

(10) International Publication Number WO 00/71726 A1

- (51) International Patent Classification⁷: C12N 15/48, 15/62, C07K 14/15, 19/00, A61K 48/00, C12Q 1/68, C07K 16/10, C12N 7/01
- (21) International Application Number: PCT/US00/14296
- (22) International Filing Date: 24 May 2000 (24.05.2000)
- (25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data: 60/135,631

24 May 1999 (24.05.1999) US

- (71) Applicant (for all designated States except US): MAYO FOUNDATION FOR MEDICAL EDUCATION AND RESEARCH [US/US]: 200 First Street SW, Rochester, MN 55905 (US).
- (72) Inventor; and
- (75) Inventor/Applicant (for US only): FEDERSPIEL, Mark, J. [US/US]; 5722 Colonial Lane SE, Rochester, MN 55904 (US).
- (74) Agent: ELLINGER, Mark, S.: Fish & Richardson P.C., P.A., Suite 3300, 60 South Sixth Street, Minneapolis, MN 55402 (US).

- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- (48) Date of publication of this corrected version:

27 June 2002

(15) Information about Correction:

see PCT Gazette No. 26/2002 of 27 June 2002, Section II

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

5

10

15

20

25

30

RECEIVED

JUL 2 3 2002

TECH CENTER 1600/2900

PCT/US00/14296

METHODS TO INHIBIT INFECTIOUS AGENT TRANSMISSION DURING XENOTRANSPLANTATION

Background of the Invention

The success of solid organ transplantation protocols continues to improve while the availability of donor organs remains limited, leading to a critical shortage (Policy et al., 1996). In 1995, only one-half of the patients on organ transplant waiting lists received transplants. Approximately nine patients per day died while waiting for a suitable organ to become available.

Xenotransplantation of animal organs is under consideration as a supplemental approach to alleviate this shortage (Policy et al., 1996). The pig is currently the donor of choice for cells, tissues and vascularized organs used in permanent and transient xenotransplantation treatments (Fishman, 1994). Swine offer a number of advantages as a source for xenotransplantations: economic and animal husbandry issues, the ability to produce transgenic animals, experience as a transplantation model, and the development of specific pathogen-free herds to eliminate known pathogens.

Pig to human organ transplants are discordant. However, several groups have engineered transgenic pig lines in efforts to control hyperacute rejection (HAR) (Lin et al., 1996). Two immunologic factors initiate HAR of xenografts: the interaction of natural xenoreactive antibodies to the Galα1-3Gal carbohydrate (α-Gal) of discordant cell proteins; and the inability of the xenograft complement regulatory proteins to control the recipient's complement system. Several approaches to control HAR are being studied including the ablation of Galα1-3Gal expression (Sandrin et al., 1995), and the design of transgenic pigs which express complement regulatory proteins (e.g., DAF, CD59) which have been shown to be protective against HAR (Diamond et al., 1996; Foder et al., 1994; McCurry et al., 1995; Oldham et al., 1996; Ryan, 1995).

Several promising xenotransplantation approaches use pig cells and tissues in protocols where xenotic cells and tissues are not rejected as readily as whole organs: fetal pig neuronal cells producing dopamine have been tested in patients to replace cells destroyed by Parkinson's disease (Deacon et al., 1997);

and fetal pig pancreatic islet cell clusters have been tested in protocols to treat diabetes (Groth et al., 1994). Another xenotransplantation approach has used pig liver and kidney xenografts as temporary support treatments for patients with fulminant organ failure (Breimer et al., 1996; Groth et al., 1994; Makowka et al., 1995). These studies have sparked research in the use of bioartificial organs and whole organ perfusion using xenogeneic tissue for bridging patients prior to transplantation. The early successes of these innovative clinical approaches indicate that xenotransplantation protocols using pig cells, tissues and organs will continue regardless of the results of solid organ xenotransplantation.

10

15

20

25

30

Besides the expected immunological hurdles that must be overcome, xenotransplantation produces a risk of transmitting infectious disease (xenoses) from the source animal tissue to the recipient, and possibly to the recipient's contacts. Xenotransplantation increases the risk of transferring infectious agents from pigs to humans due to the breach of the normal physical barriers against infectious agents upon introduction of pig cells, tissues or organs; the therapeutic immunosuppression of the recipient to prevent graft rejection which interferes with the mechanisms which defend human cells from infection by zoonotic viruses; and the genetic modifications of pigs, e.g., those being tested to control HAR or modify complement activation, may allow zoonotic viruses to evade immune surveillance or prevent the inactivation of invading enveloped viruses (Bergelson et al., 1995; Dorig et al., 1993; Fishman, 1994; Ward et al., 1994; Weiss, 1998).

Since most vertebrates carry endogenous retroviruses in their germline DNA (Boeke et al., 1997), endogenous retroviruses in xenografts may be an infectious risk to xenotransplant patients. The risks of transmission to humans with endogenous viruses of non-humans are unknown and difficult to assess. Endogenous retroviruses from a variety of animals, including baboons, cats, and mice, can infect human cells (Boeke et al., 1997). The ability of endogenous murine leukemia viruses (MLVs) to be activated and infect human tumor xenografts in immunodeficient mice is well documented (Achong et al., 1976; Beattie et al., 1982; Crawford et al., 1979; Gautsch et al., 1980; Suzuki et al., 1977; Tralka et al., 1983; Wunderi et al., 1979). The observation that the gibbon ape leukemia virus SEATO, a virus most likely derived from an endogenous

5

10

15

normal juvenile gibbon apes highlights the potential risks of exposure to xenotic endogenous retroviruses (Kawakami et al., 1980; Lieber et al., 1975).

There have been several reports of MLV retroviral vector stocks that contain the result from recombination between the vector and endogenous retroviruses in the genome of the helper cell line (Chong et al., 1998; Donahue et al., 1992; Patience et al., 1996; Purcell et al., 1996; Vanin et al., 1994). In one study, 3 of 10 rhesus monkeys developed T-cell lymphomas after autologous transplantation of enriched bone marrow stem cells transduced with a MLV retroviral vector stock containing an array of MLV-related recombinant viruses (Vanin et al., 1994). Finally, the activation of endogenous retrovirus by human tumor xenografts as well as studies of graft versus host reaction *in vivo* and mixed lymphocyte reaction *in vitro* (Hirsch et al., 1970; Hirsch et al., 1972; Levy et al., 1977; Sherr et al., 1974), demonstrate that normally inactive endogenous retroviruses can be induced by xenotransplantation conditions.

Pig endogenous retroviruses (PERVs) released by established pig cell lines were observed in the 1970s, but the host range of these viruses was thought to be restricted to pig cells (Armstrong et al., 1971; Lieber et al., 1975; Moennig et al., 1974; Strandstrom et al., 1974; Todaro et al., 1974). Recently, replication-competent pig endogenous retroviruses (PERV) were identified, and shown to be capable of productively infecting human cells *in vitro* (Wilson et al., 1998; Martin et al., 1998; Patience et al., 1997; Takeuchi et al., 1997; Weiss, 1997). PERVs have been associated with pig lymphomas (Bostock et al., 1973; Frazier, 1985; Moennig et al., 1974; Strandstrom et al., 1974; Suzuka et al., 1985).

25 Although Heneine et al. (1998) and Patience et al. (1998) screened blood lymphocytes and sera from 10 diabetic patients that received porcine fetal islet cells, and 2 renal dialysis patients whose circulation had been linked with pig kidneys extracorporeally, respectively, neither group detected PERVs. However, the risk of viral infection is increased in transplantation by the presence of factors commonly associated with viral activation, e.g., immune suppression, graft-versus-host disease, graft rejection, viral coinfection and cytotoxic therapies. Thus, in immunosuppressed xenotransplant patients, there may be a greater risk that infection would lead to disease.

5

10

15

20

25

30

Therefore, there is a need for a method to inhibit or prevent transmission of infectious agents, from donor to recipient and/or recipient to donor, after cell, tissue or organ transplant.

Summary of the Invention

The invention provides isolated and purified nucleic acid molecules and methods useful to inhibit or prevent the transmission of an infectious agent during xenogeneic or allogeneic cell, tissue or organ transplant. Infectious agents include, but are not limited to, viruses, bacteria and parasites. The invention includes prophylactic and therapeutic uses of the nucleic acid molecule of the invention which encodes at least a portion of a polypeptide of the infectious agent or which encodes at least a portion of the receptor employed by the infectious agent to enter the cells of the transplant recipient. It is preferred that the expression of the nucleic acid molecule in a donor cell, tissue or organ inhibits or prevents the transmission of the infectious agent from the donor cells, tissue or organ to a transplant recipient, or, alternatively, the transmission of the infectious agent from the recipient to the donor cell, tissue or organ. Preferred transplant recipients are mammals such as primates, e.g., humans, apes and monkeys, as well as canines, felines, bovines, ovines, swine, and equines.

Preferably, the transmission of the following infectious agents is inhibited or prevented: retroviruses, lentiviruses, herpesvirus, e.g., cytomegalovirus (CMV) and Epstein Barr virus (EBV), and hepatitis viruses, e.g., hepatitis A, B or C. Thus, transplanted organs, e.g., bioengineered organs, tissues, or cells, e.g., stem cells, including allografts, e.g., human-to-human transplants, or xenografts, are genetically modified to inhibit the infection of several major infectious pathogens that limit the success of transplantation (e.g., CMV, EBV, HIV and hepatitis viruses) or the infection by potential pathogens, e.g., endogenous viruses such as PERVs. Preferred nucleic acid molecules useful in the practice of the invention include a nucleic acid molecule which encodes gag, pol, env, protease, and accessory proteins (e.g., Vpr, Vif, Nef, Tat, or Rev) of retroviruses, lentiviruses and spumaviruses; a nucleic acid molecule which comprises a capsid or envelope gene of a herpesviruses, e.g., HSV 1 and 2, EBV and CMV; a nucleic acid molecule which comprises the C gene (nucleocapsid), ORFS/pre-S gene (viral surface glycoproteins), ORFX gene

(regulatory protein), or ORFP gene (viral polymerase) of Hepadnaviruses, e.g., hepatitis B virus; a nucleic acid molecule which comprises the C gene (viral capsid) or the E1 and/or E2 gene of Flaviviruses, e.g., hepatitis C virus; a nucleic acid molecule which comprises ORF1, e.g., which encodes Pro (protease), the X gene, the H gene (helicase), the Pol gene (polymerase), or the ORF2 gene, which codes for capsid, of hepatitis E virus; and a nucleic acid molecule which encodes the P1 gene, the 1A gene, the 1B gene, the 1C gene, or the 1D gene (capsid and glycoproteins) of hepatitis A virus. The alteration of human cells and tissue is preferred for modifying pluripotential cells and in ex vivo gene therapy protocols.

10

5

15

20

25

30

Preferably, viral entry and/or assembly is inhibited. To inhibit viral entry, receptor interference may be employed. Receptor interference refers to the expression of a receptor protein for the infectious agent or the ligand thereof, e.g., a viral ligand such as a viral glycoprotein. Preferably, the genome of the donor cell, tissue or organ is augmented with DNA encoding the viral ligand thereby preventing infection. Alternatively, or in addition, capsid-targeted viral inactivation may be employed to inhibit or prevent viral replication. Capsidtargeted viral inactivation (CTVI), reduces or eliminates the production of infectious virus by incorporating a degradative enzyme into newly synthesized viral particles, e.g., by expressing a fusion polypeptide comprising a viral polypeptide and a degradative enzyme. Degradative enzymes such as nucleases, e.g., RNase H, staphylococcal nuclease or ribozymes (see Marshall et al., 1994, and U.S. Patent No. 5,811,275), lipases or proteases, may be employed in the practice of the invention. Preferred fusion polypeptides include fusions of viral capsid, envelope or accessory proteins such as Vif, Vpx, Vpr and Nef, with degradative enzymes.

A preferred xenograft for use in the methods of the invention is swine neuronal cells, pancreatic islet cells, hepatocytes, heart, liver or kidney. Therefore, the invention provides a method in which the genomic DNA of swine cells, tissue or organ to be transplanted is augmented with a recombinant DNA molecule encoding a polypeptide of an infectious agent which binds to a cell surface receptor of a transplant recipient, e.g., a viral glycoprotein, or a fusion polypeptide comprising at least a portion of an infectious agent polypeptide and

5

10

15

20

a degradative enzyme. Infectious agents in this embodiment of the invention include, but are not limited to, those described in Table 1 of Fishman (1994), which is specifically incorporated by reference herein. In particular, PERVs are useful in CTVI methods to inhibit or prevent human-tropic PERV infection in xenotransplant patients.

As described hereinbelow, human tropic PERVs are produced by activated peripheral blood mononuclear cells. The relative PERV proviral copy number doubled upon passaging of the 1°-293 PERV-infected cells to fresh 293 cells, indicating that passaging the virus may amplify the replication-competent PERV. Only PERV-A env sequences were detected in DNA isolated from all three 293/PERV infected cultures by PCR analysis with PERV-A, PERV-B and PERV-C specific env primers. The 1°-293/PERV DNA contained a low level of PERV-C env. PERV-B env was not detected in any culture. Characterization of these PERVs indicated that human-tropic PERVs may be more diverse than previously thought. For example, significant sequence differences were found in the cloned LTR and env sequences compared to known PERV sequences. The 293/PERV LTRs contained 71 additional nucleotides compared to the PERV-MSL LTR, and the envelope amino acid sequence contains a PERV-A surface region (SU) sequence but a PERV-C transmembrane (TM) region. These PERVs are useful to determine the tissue tropism of various isolates and to determine the prevalence of anti-PERV antibodies in humans exposed to sources of PERV, e.g., pig slaughterhouse workers and xenotransplant patients such as those having a bioartificial liver with pig hepatocytes, as well as in the methods of the invention.

Thus, a vector comprising a nucleic acid molecule of the invention may also be employed to deliver soluble *env* genes to mammalian donor cells, tissues or organs prior to cell, tissue or organ transplantation so that the transplanted cells, tissues or organs are less susceptible to infection by an infectious agent, the genome of which comprises the *env* gene. For example, soluble PERV *env* genes are introduced to donor swine cells, tissues or organs to inhibit infection of the cells, tissues or organs by PERV, e.g., to inhibit infection by PERVs after the virus is reactivated following transplantation.

As also described hereinbelow, two genes encoding an avian leukosis virus (ALV) receptor protein, sTva proteins, stva and stva-mIgG, were efficiently delivered and broadly expressed by ALV-based retroviral vectors both in cultured cells and in chickens. Both the sTva and sTva-mIgG proteins significantly inhibited ALV(A) infection in vitro and in vivo. The antiviral effect was specific for ALV(A), consistent with a receptor interference mechanism.

Brief Description of the Figures

Figure 1. Stimulated pig peripheral blood mononuclear cells (PBMCs) produce replication-competent human-tropic PERV. (A) Porcine PBMC were cultured with either phytohemagglutinin (PHA) + phorbol myristate acetate (PMA) (black bars) or PMA + calcium ionophore (grey bars) and the supernatants assayed for RT activity. (B) NIH and Yucatan minipig PBMC stimulated by PHA + PMA were cocultured with ST-IOWA cells. Cell culture supernatants were assayed for RT activity. Background RT values for uninfected ST-IOWA cells have been subtracted from the values shown. NIH (black squares); Yucatan (open circles). (C) NIH minipig PBMC stimulated with PHA + PMA were cocultured with human 293 cells: 293 cells cocultured with live PBMC (closed squares); with lethally irradiated PBMC (open squares). The cell culture supernatants were assayed for RT activity.

Figure 2. Schematic overview of the generation of the human 293 cell cultures infected with PERV produced by stimulated NIH minipig primary PBMC.

20

25

30

Figure 3. Relative titers of PERV produced from the infected 293 cell cultures. The titer of MLV- β gal/PERV pseudotyped virus stocks produced from the 1°, 2° and 3°-293/PERV cultures were quantitated on human 293 cells (open bars) and pig ST-IOWA cells (black bars). The assays scored β -galactosidase producing cells (BFU) and were done in triplicate.

Figure 4. Analysis of the relative number of PERV proviruses per cell in the 293/PERV infected cultures. Genomic DNA isolated from human 293 cells and the 293/PERV infected cultures was digested with EcoRI, and the fragments separated by agarose gel electrophoresis, transferred to nitrocellulose, and probed with ³²P-labeled PERV *pol* sequences. Two internal proviral DNA fragments hybridized to the PERV *pol* probe as expected. The film was scanned

5

10

20

and the bands quantitated by Image Quant (Molecular Dynamics, Sunnyvale, CA). The blot was stripped and reprobed with ³²P-labeled human GAPDH sequences to normalize for the amount of DNA loaded per lane. After the band densities were normalized, the PERV *pol* level of the 1°-293/PERV DNA was set as 1 copy.

Figure 5. RT activity observed in cocultures of 293/2° virus producer cells and human hematopoietic cell lines. The values shown are of cpm of ³H-TTP incorporated in a reverse transcriptase assay measured in cell supernatants sampled at the times indicated post-coculture of 293/2° virus producer cells with each of the cell lines.

Figure 6. A comparison of the deduced amino acid sequences of the *env* genes of PERV-1.15 (SEQ ID NO:3 encoded by SEQ ID NO:18), PERV-A (Letissier et al., 1997; SEQ ID NO:4) and PERV-C (Akiyoshi et al., 1998; SEQ ID NO:5). A schematic representation of the regions of the PERV-1.15 gene homologous to PERV-A (hatched) and PERV-C (open) genes (top figure). The envelope surface (SU) and transmembrane (TM) glycoprotein regions are also shown. Identical amino acids are denoted by a (.); gaps are denoted (_).

Figure 7. Schematic representation of the cloned PERV sequences. A proposed model for a PERV provirus with estimated locations of the viral domains is shown at the top. The PERV sequences contained in the six unique lambda clones are shown compared to the model. The lamA8 *pol* gene contains a 86 nucleotide deletion (del), and the *env* gene contains a 101 nucleotide insertion (ins) compared to all of the comparable sequences.

Figure 8. A comparison of the nucleotide sequences of two PERV LTRs.

The LTR sequence of the published cDNA clone PERV-MSL (SEQ ID NO:7) is compared to the lamA1 LTR sequence (SEQ ID NO:6). Identical bases are denoted by (.); gaps are denoted (_). The estimated location of the major regions of the LTR, U3, R, and U5 are indicated. The sequence of the putative tRNA binding site is also indicated (18 of 18 nucleotides identical to the tRNA Pro

binding site).

Figure 9. Schematic representation of the construction of recombinant PERV molecular clones.

5

10

15

30

Figure 10. Analysis of PERV proteins by SDS-PAGE and Western blot.

(A) Virus was pelleted by ultracentrifugation through a 20% sucrose pad, separated by 12% SDS-PAGE, and visualized by staining with Coomassie brilliant blue. Lane 1, 5 ml 293 supernatant; lane 2, 5 ml 2°-293/PERV supernatant; lane 3, 15 ml 2°-293/PERV supernatant; lane 4, 5 ml DF-1 supernatant; and lane 5, 5 ml avian leukosis virus infected DF-1 supernatant. (B) Viral proteins were prepared and separated by SDS-PAGE as described above, and transferred to nitrocellulose. Lanes 1, 3, 5 and 7, 5 ml 293 supernatant; and lanes 2, 4, 6 and 8, 5 ml 2°-293/PERV supernatant. Lanes 1 and 2 were exposed to 1:1000 dilution of goat anti-GALV capsid antisera; lanes 3 and 4 to 1:1000 goat preimmune sera; lanes 5 and 6 to 1:1000 goat anti-SSAV capsid antisera; and lanes 7 and 8 to 1:1000 goat preimmune sera. The filters were washed and exposed to peroxidase-labeled rabbit anti-goat antibody. The antibody complexes were detected by chemiluminescence, and exposed to film for 30 seconds. The PERV CA protein migrates at about 25 kDa.

Figure 11. Schematic of sTva antiviral gene constructs and the ALV-based retroviral vectors.

Figure 12. General procedure for using replication-competent ALV-based retroviral vector system *in vitro* and *in vivo*.

Figure 13. sTva-mIgG receptor expression levels in DF-1 cells. The sTva-mIgG protein was immunoprecipitated with goat α-mouse-IgG agarose beads from supernatants (500 μl) of DF-1 cultures infected with the RCASBP(C), RCAS(C) or RCOSBP(C) vectors alone (V) or containing the stva-mIgG gene (sTva). The immunoprecipitates were denatured, separated by 12% SDS-PAGE, and analyzed by Western transfer. The filter was probed with peroxidase-conjugated goat anti-mouse IgG, and the bound protein-antibody complexes were visualized by chemiluminescence on Kodak X-Omat film. sTva-mIgG protein expressed transiently in human embryonic kidney 293 cells was included as a positive control (+).

Figure 14. sTva-mIgG expression in sera of chickens infected with RCASBP vectors. The sTva-mIgG protein was immunoprecipitated from chicken serum (500 µl) and analyzed as described (Figure 13 legend). Lane 1, uninfected control; Lane 2, a RCASBP(B) vector alone infected bird; Landes 3-

PCT/US00/14296 WO 00/71726

4. RCASBP(B)stva-mIgG infected birds; Lane 5, a RCASBP(C) vector alone infected bird; Lane 6, a RCASBP(C)stva-mIgG infected bird; Lane 7, RCASBP(C)stva-mIgG infected DF-1 cells as a positive control.

Figure 15. Analysis of viral and soluble receptor RNA levels in tissues of chickens infected with RCASBP(B). The figure shows autoradiograms of 6% polyacrylamide/7.6 M urea gels used to separate the protected RNA probe fragments produced in an RNase protection assay with RNA from a bird infected with RCASBP(B)stva or RCASBP(B)stva-mIgG. RNA was prepared from liver (L), heart (H), spleen (S), bursa (B), thymus (T), kidney (K), and muscle (M) tissues of each bird. RNA from DF-1 cells infected with the appropriate virus 10 was included as a positive control (+). RNA from bursa of an uninfected bird was the negative control (-). ALV(B) env RNA protects a 467-nt fragment from the 522-nt ³²P-labeled full-length probe [env(B)]; stva RNA protects a 388-nt fragment from the 498-nt probe (stva); and stva-mIgG RNA protects a 363-nt fragment from the 423-nt probe (stva-mIgG). Each assay contained a chicken GAPDH probe as a control for RNA quality and quantity. GAPDH RNA protects a 200-nt fragment from the 279-nt GAPDH probe.

15

Figure 16. Analysis of viral and soluble receptor RNA levels in the bursa of birds infected with RCASBP(B) vectors. The figure shows an autoradiogram of a 6% polyacrylamide/7.6 M urea gel used to separate the protected RNA 20 probe fragments produced in an RNase protection assay with RNA from the bursa of birds infected with RCASBP(B) alone (V), RCASBP(B)stva (S), RCASBP(B)stva-mIgG (I), or uninfected (U). RNA from DF-1 cells infected with the appropriate virus was included as a positive control (+). ALV(A) env RNA protects a 423-nt fragment from the 483-nt ³²P-labeled full-length probe 25 [env(A)]: ALB(B) env RNA protects a 467-nt fragment from the 522-nt probe [env(B)]; stva RNA protects a 388-nt fragment from the 498-nt probe (stva); and stva-mIgG RNA protects a 363-nt fragment from the 423-nt probe (stva-mIgG). Each assay contained a chicken GapDH probe as a control for RNA quality and quantity. GAPDH RNA protects a 200-nt fragment from the 279-nt GAPDH 30 probe. Full-length probes not treated with RNase: lane A, env(A); lane B, env(B); lane S, stva; lane I, stva-mIgG; and Lane G, GAPDH.

Figure 17. Analysis of viral and soluble receptor DNA in tissues of RCASBP(B) infected birds and challenged with RAV-1. Genomic DNA was isolated from liver (L), heart (H), spleen (S), bursa (B), thymus (T), kidney (K), and muscle (M) samples of birds infected with RCASBP(B) alone (I),

RCASBP(B)stva (II) and RCASBP(B)stva-mIgG (III) and challenged with RAV-1. DNA isolated from DF-1 cells infected with the appropriate virus was used as a positive control (+). DNA isolated from the bursa of an uninfected control bird was used as the negative control (-). DNA sequences were detected by PCR using specific primer pairs: RAV-1 challenge virus DNA [env(A)] was detected using primers specific for ALV(A) *env* yielding a 937 bp fragment; RCASBP(B) vector DNA [env(B)] was detected using primers specific for ALV(B) *env* yielding a 429 bp fragment; *stva* DNA (stva) and *stva-mIgG* DNA (stva-mIgG) were detected with specific primers yielding fragments of 314 bp and 589 bp respectively. The amplified DNA fragments were separated on 0.8% agarose gels and visualized with ethidium bromide. The molecular weight marker is the 1 Kb plus DNA ladder (Gibco).

Figure 18 A) Nucleotide sequence of env gene of PERV 1.15 (SEQ ID NO:18). B) Nucleotide sequence of PERV sequences in lambda A1 clone (SEQ ID NO:19). C) Nucleotide sequence of PERV sequences in lambda A10 clone (SEQ ID NO:20). D) Nucleotide sequence of PERV sequences in lambda A11 clone (SEQ ID NO:32). E) Nucleotide sequence of PERV sequences in lamba A3A clone (SEQ ID NO:21). F) Nucleotide sequence of PERV sequences in lambda A6 clone (SEQ ID NO:22). G) Nucleotide sequence of PERV sequences in lambda clone A8 (SEQ ID NO:23).

Detailed Description of the Invention

Definitions

5

10

15

20

25

As used herein, the terms "isolated and/or purified" refer to *in vitro* preparation, isolation and/or purification of a nucleic acid molecule or polypeptide (e.g., antibody) of the invention, so that it is not associated with *in vivo* substances. Thus, with respect to an "isolated nucleic acid molecule", which includes a polynucleotide of genomic, cDNA, or synthetic origin or some combination thereof, the "isolated nucleic acid molecule" (1) is not associated with all or a portion of a polynucleotide in which the "isolated nucleic acid

molecule" is found in nature, (2) is operably linked to a polynucleotide which it is not linked to in nature, or (3) does not occur in nature as part of a larger sequence. An isolated nucleic acid molecule means a polymeric form of nucleotides of at least 10 bases in length, either ribonucleotides or

- deoxynucleotides or a modified form of either type of nucleotide. The term includes single and double stranded forms of DNA. The term "oligonucleotide" referred to herein includes naturally occurring, and modified nucleotides linked together by naturally occurring, and non-naturally occurring oligonucleotide linkages. Oligonucleotides are a polynucleotide subset with 200 bases or fewer in length. Preferably, oligonucleotides are 10 to 60 bases in length, and most are the second process.
- in length. Preferably, oligonucleotides are 10 to 60 bases in length and most preferably 12, 13, 14, 15, 16, 17, 18, 19, or 20 to 40 bases in length.

 Oligonucleotides are usually single stranded, e.g., for probes; although oligonucleotides may be double stranded, e.g., for use in the construction of a variant. Oligonucleotides of the invention can be either sense or antisense
- oligonucleotides. The term "naturally occurring nucleotides" referred to herein includes deoxyribonucleotides and ribonucleotides. The term "modified nucleotides" referred to herein includes nucleotides with modified or substituted sugar groups and the like. The term "oligonucleotide linkages" referred to herein includes oligonucleotides linkages such as phosphorothioate.
- phosphorodithioate, phophoroselenoate, phosphorodiselenoate, phosphoroanilothioate, phosphoraniladate, phosphoroanidate, and the like. An oligonucleotide can include a label for detection, if desired.

25

30

The term "isolated polypeptide" means a polypeptide encoded by cDNA or recombinant RNA, or is synthetic origin, or some combination thereof, which isolated polypeptide (1) is not associated with proteins found in nature, (2) is free of other proteins from the same source, e.g., free of human proteins, (3) is expressed by a cell from a different species, or (4) does not occur in nature.

The term "sequence homology" means the proportion of base matches between two nucleic acid sequences or the proportion amino acid matches between two amino acid sequences. When sequence homology is expressed as a percentage, e.g., 50%, the percentage denotes the proportion of matches over the length of a sequence that is compared to some other sequence (the reference sequence). Gaps (in either of the two sequences) are permitted to maximize

matching; gap lengths of 15 bases or less are usually used, 6 bases or less are preferred with 2 bases or less more preferred. When using oligonucleotides as probes or treatments, the sequence homology between the target nucleic acid and the oligonucleotide sequence is generally not less than 17 target base matches out of 20 possible oligonucleotide base pair matches (85%); preferably not less than 9 matches out of 10 possible base pair matches (90%), and more preferably not less than 19 matches out of 20 possible base pair matches (95%).

The term "selectively hybridize" means to detectably and specifically bind. Polynucleotides, oligonucleotides and fragments of the invention selectively hybridize to nucleic acid strands under hybridization and wash conditions that minimize appreciable amounts of detectable binding to nonspecific nucleic acids. High stringency conditions can be used to achieve selective hybridization conditions as known in the art and discussed herein. Generally, the nucleic acid sequence homology between the polynucleotides, oligonucleotides, and fragments of the invention and a nucleic acid sequence of interest is at least 65%, and more typically with preferably increasing homologies of at least about 70%, about 90%, about 95%, about 98%, and 100%. Preferred PERV nucleic acid sequences for use in the methods of the invention are those which have at least about 80%, more preferably 90%, and even more preferably 95%, contiguous nucleotide sequence homology or identity to SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23 or SEQ ID NO:32. Preferably, PERV nucleic acid sequences are those which hybridize under moderate, preferably under stringent, hybridization conditions to SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23 or SEQ ID NO:32.

20

25

30

Two amino acid sequences are homologous if there is a partial or complete identity between their sequences. For example, 85% homology means that 85% of the amino acids are identical when the two sequences are aligned for maximum matching. Gaps (in either of the two sequences being matched) are allowed in maximizing matching; gap lengths of 5 or less are preferred with 2 or less being more preferred. Alternatively and preferably, two protein sequences (or polypeptide sequences derived from them of at least 30 amino acids in length) are homologous, as this term is used herein, if they have an alignment

score of at more than 5 (in standard deviation units) using the program ALIGN with the mutation data matrix and a gap penalty of 6 or greater. See Dayhoff, M. O., in Atlas of Protein Sequence and Structure, 1972, volume 5, National Biomedical Research Foundation, pp. 101-110, and Supplement 2 to this volume, pp. 1-10. The two sequences or parts thereof are more preferably homologous if their amino acids are greater than or equal to 50% identical when optimally aligned using the ALIGN program.

5

10

15

20

25

The term "corresponds to" is used herein to mean that a polynucleotide sequence is homologous (i.e., is identical, not strictly evolutionarily related) to all or a portion of a reference polynucleotide sequence, or that a polypeptide sequence is identical to a reference polypeptide sequence. In contradistinction, the term "complementary to" is used herein to mean that the complementary sequence is homologous to all or a portion of a reference polynucleotide sequence. For illustration, the nucleotide sequence "TATAC" corresponds to a reference sequence "TATAC" and is complementary to a reference sequence "GTATA".

The following terms are used to describe the sequence relationships between two or more polynucleotides: "reference sequence", "comparison window", "sequence identity", "percentage of sequence identity", and "substantial identity". A "reference sequence" is a defined sequence used as a basis for a sequence comparison; a reference sequence may be a subset of a larger sequence, for example, as a segment of a full-length cDNA or gene sequence given in a sequence listing, or may comprise a complete cDNA or gene sequence. Generally, a reference sequence is at least 20 nucleotides in length, frequently at least 25 nucleotides in length, and often at least 50 nucleotides in length. Since two polynucleotides may each (1) comprise a sequence (i.e., a portion of the complete polynucleotide sequence) that is similar between the two polynucleotides, and (2) may further comprise a sequence that is divergent between the two polynucleotides, sequence comparisons between two (or more) polynucleotides are typically performed by comparing sequences of the two polynucleotides over a "comparison window" to identify and compare local regions of sequence similarity.

PCT/US00/14296 WO 00/71726

A "comparison window", as used herein, refers to a conceptual segment of at least 20 contiguous nucleotides and wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) of 20 percent or less as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. Optimal alignment of sequences for aligning a comparison window may be conducted by the local homology algorithm of Smith and Waterman (1981) Adv. Appl. Math. 2: 482, by the homology alignment algorithm of Needleman and Wunsch (1970) J. Mol. Biol. 48: 443, by the search for similarity method of Pearson and Lipman (1988) Proc. Natl. Acad. Sci. (U.S.A.) 85: 2444, by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package Release 7.0, Genetics Computer Group, 575 Science Dr., Madison, Wis.), or by inspection, and the best alignment (i.e., resulting in the highest percentage of homology over the comparison window) generated by the various methods is 15 selected.

10

20

25

30

The term "sequence identity" means that two polynucleotide sequences are identical (i.e., on a nucleotide-by-nucleotide basis) over the window of comparison. The term "percentage of sequence identity" means that two polynucleotide sequences are identical (i.e., on a nucleotide-by-nucleotide basis) over the window of comparison. The term "percentage of sequence identity" is calculated by comparing two optimally aligned sequences over the window of comparison, determining the number of positions at which the identical nucleic acid base (e.g., A, T, C, G, U, or I) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison (i.e., the window size), and multiplying the result by 100 to yield the percentage of sequence identity. The terms "substantial identity" as used herein denote a characteristic of a polynucleotide sequence, wherein the polynucleotide comprises a sequence that has at least 85 percent sequence identity, preferably at least 90 to 95 percent sequence identity, more usually at least 99 percent sequence identity as compared to a reference sequence over a comparison window of at least 20 nucleotide positions, frequently over a window of at least 20-50 nucleotides,

wherein the percentage of sequence identity is calculated by comparing the reference sequence to the polynucleotide sequence which may include deletions or additions which total 20 percent or less of the reference sequence over the window of comparison.

As applied to polypeptides, the term "substantial identity" means that two peptide sequences, when optimally aligned, such as by the programs GAP or BESTFIT using default gap weights, share at least about 80 percent sequence identity, preferably at least about 90 percent sequence identity, more preferably at least about 95 percent sequence identity, and most preferably at least about 99 percent sequence identity.

Moderate and stringent hybridization conditions are well known to the art, see, for example sections 9.47-9.51 of Sambrook et al. (Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1989). For example, stringent conditions are those that (1) employ low ionic strength and high temperature for washing, for example, 0.015 M NaCl/0.0015 15 M sodium citrate (SSC); 0.1% sodium lauryl sulfate (SDS) at 50°C, or (2) employ a denaturing agent such as formamide during hybridization, e.g., 50% formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50 mM sodium phosphate buffer at pH 6.5 with 750 mM NaCl, 75 mM sodium citrate at 42°C. Another example is use of 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% sodium dodecylsulfate (SDS), and 10% dextran sulfate at 42°C, with washes at 42°C in 0.2 x SSC and 0.1% 25 SDS.

CTVI

30

5

10

The present invention describes a method of interfering with viral replication. For example, the fusion of an enzyme (such as a nuclease, lipase, or protease) to a viral capsid, glycoprotein or accessory polypeptide which is incorporated into virions can inactivate essential viral components, such as nucleic acid, protein or lipid, which are necessary for replication. This technique is referred to as capsid-targeted viral inactivation (CTVI), although it includes the targeting of non-capsid fusions, e.g., env fusions, to the virion as well.

The present invention uses nucleases, proteases, and/or lipases as antiviral therapeutic agents that degrade or modify essential viral components, such as nucleic acids, proteins, or lipids. The method of the present invention can be applied to viruses that have relatively flexible capsid structures such as retroviruses, lentiviruses, herpesviruses, poxviruses, togaviruses, hepadnaviruses, caulimoviruses, myxoviruses and paramyxoviruses, that readily allow the encapsidation of foreign proteins in an aqueous internal compartment containing the viral nucleic acid. Preferred viruses to be inhibited by the methods of the invention include porcine endogenous retroviruses (PERVs), herpesviruses such as EBV and CMV, and hepatitis viruses, e.g., hepatitis A, B, 10 and C. See Fields, Virology, Lippincourt-Raven, Philadelphia, PA (1996), which is specifically incorporated by reference herein. The enzyme portion of the fusion protein preferably is oriented so that it is internal to the assembled capsid structure. However, externally facing enzymes may be used in the case of proteases and lipases. The use of such constructs does not require the virus to 15 have an aqueous internal compartment. In the case of lipases, it is desirable to target the enzyme to the envelope of the virus. The enzyme portion can be either inside or outside the viral envelope. In this way, any enveloped virus can be targeted by CTVI.

Retroviruses, and the closely related retrotransposons, which are referred 20 to herein generically as retroviruses, are particularly susceptible to CTVI because of their assembly mechanism. It is possible to take advantage of the knowledge of how these capsids assemble to direct a destructive enzyme molecule to the inside surface of the capsid, where it can contract viral nucleic acids or proteins. Retroviral and retrotransposon reverse transcriptases are expressed as natural 25 fusion proteins, in which the polymerase protein forms the C-terminal portion of the primary translation product and the gag protein forms the N-terminal portion. The mechanisms by which Gag-Pol fusion proteins are formed in different retroviruses vary, sometimes involving frameshifting during translation, suppression of a nonsense codon, or splicing. Regardless of the mechanism of 30 synthesis, the reverse transcriptase protein sequence is found at the C-terminus of the Gag protein sequences. In addition, the reverse transcriptase is found internal to the capsid structure. Substitution of a destructive enzyme coding

region for the reverse transcriptase gene (i.e., inserting an open reading frame for a destructive enzyme into the *pol* gene in frame), leads to the assembly of the enzyme in retroviral particles. Further, the nuclease or protease is oriented relative to the capsid in such a way as to have access to the viral RNA or reverse transcriptase. Gag-degradative enzyme fusions, as well as gag-pol-degradative enzyme fusions and env-degradative enzyme fusions, are envisioned. Further, the amino acid linkage between the viral proteins and the degradative enzyme may include sequences that are susceptible to proteases present in the virion, and so are cleaved to yield a soluble degradative enzyme.

Thus, a dominant negative effect is obtained by fusion of a protein with a destructive or modifying enzymatic activity that can alter or destroy an essential component of the virus (e.g., DNA, RNA or protein). The destructive or modifying activity is targeted to the desired site of action (inside or closely adjacent to the virus particle) because it is covalently linked to a core protein of the viral particle, e.g., in the same manner as reverse transcriptase.

10

15

20

25

30

Besides its versatility, one of the main advantages of CTVI is that it is enzymatic. It does not rely on massive over-expression of a mutant protein that in some way interferes with the viral life cycle, as is the case in the strategies referenced above. Rather, a small number of enzyme molecules (e.g., nucleases) are incorporated into a viral particle, and they cleave or modify one or more essential components of the virus.

CTVI also has advantages over traditional chemotherapy. Viruses,

especially viruses with error-prone replicative mechanisms such as HIV and other retroviruses, are likely to become resistant to a chemotherapeutic agent. For example, HIV isolates resistant to AZT, a widely used anti-HIV drug, have been reported. Clearly, resistance to drugs is an important problem with viral infections just as it is for bacterial infections. Gene therapy or intracellular immunization strategies may also be subject to problems of resistance. For example, if a virus mutates such that the capsid protein no longer assembles with the capsid fusion protein of the present invention, the virus will become resistant to the particular CTVI therapy. However, the DNA of the mutant virus can be cloned and the mutant capsid gene can be fused to the gene encoding a nuclease or other enzyme, thereby restoring the efficacy of the therapy. This approach is

not possible with normal drug therapy; once a virus becomes resistant to a given drug, use of that drug has to be abandoned.

Fusion proteins according to the present invention comprise a destructive enzyme which is capable of destroying the replication ability of a virus or virus-like particle. As used in the present invention, "replication ability" encompasses infectivity of a virus particle or transposition of a retrotransposon. Typically, the destructive enzyme is a nuclease, lipase or protease. Protein-modifying enzymes, such as kinases, which diminish enzyme activity can also be used, for example, one which inactivates a viral encoded polymerase. Fusion proteins also comprise a capsid protein, such as Gag, which confers upon the fusion proteins the ability to co-assemble into a virus particle or envelope protein. As mentioned above, it may be desirable in the case of lipase fusions that the virus-encoded portion of the fusion protein be an envelope protein, rather than a capsid protein.

10

Nucleases which can be used in the practice of the present invention to make fusions include restriction endonucleases such as EcoRI, and less specific RNAses, such as T1 RNase, barnase, RNase H1, RNase III, retroviral RNase H domains from reverse transcriptase, and staphylococcal nuclease (SN). Other destructive enzymes which can be used in the practice of the present invention include proteases, e.g., nonhomologous retroviral proteases, and lipases. Any such enzymes known in the art can be used. These enzymes degrade proteins and lipids which may be essential for the replication (infectivity) of particular viruses. Other enzymes which modify protein structures may be used, if they result in the inactivation of proteins which are necessary for the viral life cycle.

25 such protein modification enzymes include: kinases, glycosylases, phosphatases, methyl transferases, acetylases, acylases, farnesyl transferases, and demyristylases.

High levels of expression of the fusion proteins are not required, as a single active enzyme molecule can be sufficient to inactivate a virus particle. However, it is desirable that each virus particle formed have at least a single molecule of the fusion protein. Thus, statistical considerations would dictate that expression at a rate greater than one molecule of fusion protein per virus particle be obtained.

conditionally active or marginally active enzymes, since the destructive activity of the enzyme may be deleterious to cells used to prepare the constructs of the present invention. Thus, for example, calcium dependent enzymes may be used, which require a threshold level of calcium for activity. Calcium-dependent destructive enzymes which may be used include calpain, phospholipase A2 and staphylococcal nuclease. It may also be desirable to non-conditionally diminish the activity of the destructive enzyme to a level which is not toxic to cells but which is toxic to viruses. Random mutagenesis can be used to accomplish this end.

DNA molecules are also provided according to the invention, which code for fusion proteins as described above. The DNA molecules comprise appropriate transcriptional and translational control signals which allow the DNA molecules to be expressed in the host cells of the viruses from which the capsid or envelope protein is derived. The expression of the fusion proteins may be dependent upon the presence of virus in the cell.

15

20

25

30

Recombinant virus is also provided by the present invention. These particles comprise a nucleic acid molecule encoding a fusion protein, as described above. Thus, infection of cells with these recombinant viruses will yield fusion proteins. The recombinant virus is desirably not related to the virus from which the capsid or envelope protein of the fusion protein is derived. The recombinant virus is a mere vehicle for introduction of the fusion constructs. When the cells are infected with viruses which are cognate to the fusion protein capsid or envelope portion, the normal infection cycle of the viruses proceeds. However, when the viruses encapsidate viral nucleic acid, wild-type capsid or envelope proteins co-assemble with fusion proteins. Thus, the newly assembled viruses are hybrids comprising wild-type and fusion capsid or envelope proteins. The newly assembled viruses are not infective because the enzyme portion of the fusion protein inactivates either the viral nucleic acid or the essential viral-packaged proteins or lipid comprising the viral envelope.

Exemplary nucleic acid molecules are those comprising PERV nucleic acid sequences. Thus, for fusion proteins of the invention, it is preferred that nucleic acid molecules encoding PERV capsid or env are employed. For

example, recombinant virus may be obtained from cells, e.g., helper cells, transfected or infected with a recombinant DNA molecule encoding a fusion protein comprising PERV capsid linked to a degradative enzyme. Preferably, once the fusion protein is packaged into virions, the degradative enzyme is proteolytically cleaved from the fusion protein, i.e., it is soluble within the virion.

A method of inhibiting virus replication or infectivity is also provided by the present invention. A nucleic acid molecule encoding a fusion protein is introduced into a cell susceptible to a virus. The fusion protein may comprise a capsid or envelope protein of the virus as well as a destructive enzyme as described above. The method of introducing the nucleic acid molecule into the cell is not critical to the invention, and many such methods are known in the art. These include, without limitation: infection, transformation, transfection, lipofection, tungsten microprojectiles, electroporation, cell fusion, and transduction.

10

15

20

25

30

Nucleic acid molecules encoding the fusion protein of the present invention can be introduced into solid organs, tissues or cells, e.g., stem cells of the hematopoietic lineage or B lymphocytes. T-lymphocytes and monocytes/macrophages, which are derived from hematopoietic stem cells, are the primary target of viruses such as HIV. Bone marrow cells can be taken from an individual, and if desired, the stem cells can be purified from the mixed population of marrow cells. The nucleic acid molecules of the present invention can be introduced into the stem cells by means of transfection, or any other means known in the art. The transfected stem cells can then be reinfused into the individual (autologous stem cell transplantation). To facilitate proliferation of the transplanted stem cells, the individual's bone marrow can be partially cleared by irradiation or chemotherapy.

In another embodiment, the nucleic acid molecule of the present invention is introduced into an animal, or gametic cells or embryos. Techniques such as microinjection or transfection can be used, as is well known to the art. The nucleic acid-treated cells can be used in artificial insemination (gametes) or can be reimplanted into a hormone-prepared female animal (embryos) as is known in the art.

10

20

25

30

In yet another method of introducing the nucleic acids of the present invention into cells or an animal, a recombinant virus can be used. The recombinant virus is desirably derived from a virus having the same cell-type tropism as the virus from which the capsid or envelope protein of the gene fusion is derived. Thus, for example, in order to introduce a gene fusion which employs a PERV capsid protein gene, a PERV viral vector may be used. The recombinant virus is then administered according to a route of viral infection for the particular virus of the vector.

An alternative way for introducing the fusion protein encoding construct of the present invention into cells and animals is by means of hybrid virus particles which contain the nucleic acid of the present invention but do not contain the fusion protein of the invention. Hybrid viruses can be prepared which contain the nucleic acid encoding the fusion protein of the present invention using a packaging cell line. Such cell lines may be propagated in culture according to techniques well known in the art. Such cell lines produce the proteins necessary for packaging viral genomes. The nucleic acids of the present invention can be introduced into such a packaging cell line. Desirably, the nucleic acids will contain the packaging signal which is recognized by the packaging proteins of the cell line. Preferably, the fusion protein gene is under the control of an inducible promoter, such as the metallothionine promoter. When it is desirable to produce viral particles carrying the nucleic acid of the present invention, the inducible promoter can be induced to turn on transcription of the fusion protein gene construct, and a protein synthesis inhibitor, such as cycloheximide, can be added so that the fusion protein transcript is not translated in the packaging cell line. Thus, the fusion protein gene construct is transcribed and packaged into the particles produced by the cell line, without interference from (co-assembly with) the fusion protein. When the packaged particles are used to infect cells, the nucleic acid encoding the fusion protein is expressed, and the fusion protein becomes co-assembled into the progeny particles, rendering them non-infectious.

Methods of producing animals, e.g., humans or other non-human animals, which are resistant to a virus are also provided by the present invention.

A nucleic acid molecule which encodes a fusion protein as described above is

introduced into the germ line of an animal or an embryo, or to the animal, e.g., via DNA immunization or by infection with recombinant virus. Methods for the introduction of nucleic acid molecules to animals are known in the art (see, for example, Salter et al., 1991; Salter et al., 1989; Henininghausen, 1990; Ebert et al., 1991; Wright et al., 1991; Jaenish, 1988; Wagner et al. (U.S. Patent No. 4,873,191); Donahue et al., 1992; Rogers et al., 1993). All somatic cells resulting from development of zygotes containing the nucleic acid molecule are resistant to the virus from which the fusion protein capsid or envelope portion is derived. Methods for making transgenic animals which carry foreign genes in each cell of the animal are known in the art.

Receptor Interference (RI)

10

15

20

25

Another antiviral strategy targets the first step in the virus life cycle, the interaction of the viral envelope glycoprotein(s) with a host cell receptor. For example, retroviral infection of cells can be significantly reduced by expressing soluble forms of the host receptor, which bind the glycoproteins of the incoming virions before the virion can bind to the host receptor, or the viral envelope glycoproteins (the product of the *env* gene), which block the host receptors, directly preventing virion binding.

There are at least two distinguishable steps in viral entry; the first involves recognition and binding of the viral envelope glycoprotein to the host receptor, the second involves steps that lead to the fusion of the host and viral membranes. The *env* gene of retroviruses and lentiviruses encodes a polyprotein precursor that is subsequently processed into two glycoproteins, the surface glycoprotein (SU) which contains the domains that interact with the host receptor, and the transmembrane glycoprotein (TM) that anchors the SU protein to the membrane and appears to be directly involved with the fusion between the virus and cell membranes. Following membrane fusion, the viral core is released into the cytoplasm where the single-stranded RNA genome found in the virion is copied into double-stranded DNA (dsDNA) by the viral enzyme reverse transcriptase (RT). This reverse transcription process produces a linear dsDNA that is longer than the RNA genome from which it derives. The ends of the linear DNA contain terminally redundant sequences known as long terminal repeats (LTRs). Retroviruses, and vectors that derive from them, appear to

require the breakdown of the nuclear membrane that occurs during mitosis for the viral DNA to gain access to the cellular genome. Once this occurs, the linear viral DNA is inserted into the host genome by the virally encoded enzyme integrase (IN). The integrated viral DNA is called a provirus. Efficient production of progeny viruses requires the integration of the viral genome into host DNA.

Cells and animals engineered to express envelope glycoproteins have been shown to be highly resistant to that strain of retrovirus and to have less virus-associated pathogenesis (Federspiel et al., 1989; Nihrane et al., 1996; 10 Salter et al., 1989). Based on the receptor interference of endogenous ALV(E), Salter and Crittenden (1989) proposed that insertion of the ALV(A) envelope gene into the germline of chickens and its subsequent expression would provide resistance to infection by ALV(A). Their initial germline studies produced 23 lines of transgenic chickens using ALV-based retroviral vectors and proved two major points. First, these studies showed that avian retroviruses can be 15 artificially inserted into the germline of chickens at a reasonable efficiency, stably inherited by subsequent generations, and that the viral genes were expressed (Crittenden et al., 1989; Salter et al., 1987). Second, these studies showed that expression of the inserted subgroup (A) envelope glycoproteins 20 could inhibit subsequent ALV(A) infection by subgroup-specific receptor interference, both in vitro and in vivo (Salter et al., 1989; Salter et al., 1991).

Two transgenic chicken lines, alv6 and alv11, that did not produce infectious virus were extensively characterized at the RNA level in different tissues and at different stags of development. Viral RNA was expressed by both lines in all tissues tested regardless of the stage of development; insert alv6 expressed moderate levels of env RNA, while insert alv11 expressed low levels of gag and env. Both lines, however, expressed much higher levels of viral RNA in the tissues associated with ALV pathogenesis, the bursa, thymus, and spleen. In vitro, both alv6 (about 4400-fold) and alv11 (about 11-fold) chicken embryo fibroblasts (CEF) were resistant to ALV(A) infection. The susceptibilities of the alv6 and alv11 transgenic lines to ALV(A) infection and associated pathogenesis were compared to other properties (e.g., resistance measured in vitro, env expression levels, etc.) to determine which characteristics were important

25

5

10

20

25

30

indicators of an antiviral effect in vivo. None of the alv6 birds were productively infected as determined by the absence of both infectious ALV(A) and ALV(A) neutralizing antibodies, while two-thirds of the alv11 birds were productively infected by ALV(A). However, neither the alv6 nor the alv11 chickens showed significant ALV-induced pathogenesis. Thus, the level of Env glycoprotein expressed in the target tissues for ALV pathogenesis determined the magnitude of ALV interference (Federspiel et al., 1991).

Reticuloendotheliosis virus (REV) is very cytotoxic to avian cells *in vitro* presumably due to the fusionogenic properties of the envelope glycoproteins, especially the TM glycoprotein. REV stocks are therefore produced in D17 cells, a canine osteosarcoma line tolerant to REV infection. D17 cell lines expressing the REV envelope glycoproteins were significantly more resistant to REV infection (up to 25,000-fold) compared to controls. However, when the same REV envelope expression cassette was delivered into chicken embryo fibroblasts (CEF), the REV envelope gene sustained large deletions and the envelope proteins were not expressed due to cytotoxicity of the REV envelope glycoproteins to CEF cells. Thus, to reduce potential cytotoxic effects, it is preferred that soluble forms of viral envelope glycoprotein are employed. The soluble protein differs from the wild-type protein in that it lacks the transmembrane (TM) domain which anchors the protein in the membrane, and are therefore secreted from the cell.

The invention will be further described by the following examples.

Example 1

Expression of a Mo-MLV Gag-E. coli RNase H Fusion Polyprotein

A gene encoding a Mo-MLV Gag-*E. coli* RNase HI fusion was constructed by linking the RNase HI coding region to the Mo-MLV Gag polyprotein six codons upstream of the Gag termination codon (Van Brocklin et al., 1997). The RNase HI coding sequence was isolated from *E. coli* DNA by PCR amplification with a 5' primer containing a BglII site (5'-GCG CAT GCA GAT CTG ATG CTT AAA CAG GTA GAA ATT TTC ACC GAT GG-3'; SEQ ID NO:1) and a 3' primer containing a SalI site (5'-GCT GCT GCG TCG ACT TAA ACT TCA ACT TGG TAG CCT GTA TCT TCC-3'; SEQ ID NO:2). The reagents and conditions used for the PCR were as described in Federspiel et al.

5

10

15

20

25

30

(1996). The *E. coli* RNase HI fragment was sequenced and contained one silent nucleotide difference from the sequence published by Kanaya and Crouch (1983), at nucleotide 470 (A-G). The Mo-MLV Gag-RNase H fusion construct was based on the design of the Mo-MLV Gag-SN fusion which has been used to introduce SN into virions (Natsoulis et al., 1995).

The genes encoding the CTVI constructs were delivered into CEF by the RCASBP(A) retroviral vector and expressed under the control of the viral promoter-enhancer in the long terminal repeat. CEF were cultured and passaged every 2 days when they reached confluence (Federspiel et al., 1994). ALV retroviral vector propagation was initiated by transfection of plasmid DNA that contained the retroviral vector in a proviral form by the calcium phosphate precipitation method (Kingston et al., 1989). The course of the retroviral infection was monitored by assaying culture supernatants from confluent cells for the viral Gag protein. Three plasmids were used for comparison: Mo-MLV Gag-SN plasmid pGN1600 (Natsoulis et al., 1995); Mo-MLV Gag plasmid pGN1601 (Natsoulis et al., 1995), which produces only the Mo-MLV Gag polyprotein; and Mo-MLV Gag-SN* plasmid pGS293 (Schumann et al., 1996), which differs from Mo-MLV Gag-SN in two missense mutations in the SN gene that result in an inactive SN enzyme. Maximum levels of Mo-MLV Gag, Mo-MLV Gag-RNase H, Mo-MLV Gag-SN, and Mo-MLV Gag-SN* were seen 10 to 14 days posttransfection, which coincided with maximum RCASBP(A) production. Immunoblot analysis indicated that the Mo-MLV Gag-RNase H polyprotein was expressed at a lower level than Mo-MLV Gag or Mo-MLV Gag-SN. Several viral proteins that were smaller than the expected full-length polyproteins were also observed. These proteins were the likely result of partial degradation of the Pr85 or Pr65 polyprotein associated with the virion isolation procedure, since smaller Gag-associated proteins were also seen in the Mo-MLV Gag control. Southern blot analysis of genomic DNA isolated from day 24 cultures detected only intact proviruses. An in situ RNase H assay showed that virus particles isolated from CEF expressing the Mo-MLV Gag-RNase H polyprotein contained high levels of RNase H activity. Little or no RNase H activity was detected in any other culture supernatants.

CEF Expressing the Mo-MLV Gag-RNase H Polyprotein Significantly

Inhibit Mo-MLV Spread. The antiviral effect of the CTVI fusions expressed by the CEF cultures was measured first in a prophylaxis assay. The cultures were challenged with a low dose (multiplicity of infection [MOI] of 0.05 to 0.1 focusforming units [FFU] per cell) of amphotropic Mo-MLV strain Mo(4070A) (Ott et al., 1990; Ott et al., 1992). Mo(4070A) is an engineered hybrid Mo-MLV that contains the env gene and a portion of the pol gene of the amphotropic 4070A virus. The Mo(4070A) virus stock was produced on NIH 3T3 cells (1×10^6 to 2 \times 106 FFU/ml). Infectious virus was quantitated by the S+L- focus assay on D56 cells (Bassin et al., 1971). The CEF cultures, which were dividing rapidly, 10 were passaged for 14 days after Mo(4070A) infection. Infectious Mo(4070A) was quantitated at day 14 by the S+L- focus assay (Table 1). CEF expressing the Mo-MLV Gag-RNase H polyprotein reduced the level of infectious Mo(4070A) produced by more than 1,500-fold compared to the Mo(4070A)-15 infected CEF control.

Table 1. Inhibition of Mo-MLV spread by CTVI^a

		Titer (FFU/ml) ^b (fold inhibition) ^c		
	Construct used for treatment	Expt 1	Expt 2	
0	None (mock treatment)	1.6×10^{5}	2.1×10^{5}	
	Vector alone	$2.4 \times 10^4 (6.6)$	ND^d	
	Mo-MLV Gag	$2.5 \times 10^4 (6.3)$	$4 \times 10^4 (5)$	
	Mo-MLV Gag-RNase H	$1 \times 10^2 (1,550)$	$9 \times 10^{1} (2,330)$	
	Mo-MLV Gag-SN	$5 \times 10^{1} (3,100)$	$1.6 \times 10^{1} (13,125)$	
5	Mo-MLV Gag-SN*	$1.9 \times 10^4 (8.4)$	ND	

[&]quot; In a prophylaxis assay, CEF cultures seven passages after transfection with plasmids containing the RCASBP(A) retroviral vector-CTVI constructs were challenged with Mo(4070A) (multiplicity of infection, 0.05 to 0.1 FFU per cell). Infectious Mo(4070A) levels were quantitated from supernatants 14 days postinfection by S+L- focus assays. This experiment was performed twice.

30

35

^b Titers were averages of duplicate assays.

^c Fold inhibition was determined by comparing the Mo(4070A) titers of the experimental constructs to the Mo(4070A) titer on mock-treated CEF.

^d ND, not done.

5

Several cultures were further characterized over the course of the 14-day Mo(4070A) infection by quantitating the levels of infectious Mo(4070A) by S+L- focus assay and analyzing the expression levels of the Mo-MLV Gag and RCASBP(A) Gag polyproteins by Western immunoblotting. The amount of

- infectious Mo(4070A) produced peaked 7 to 8 days after infection in the control cultures (CEF alone and CEF expressing Mo-MLV Gag), and the titer remained relatively constant over the next 6 days. In contrast, the amount of infectious Mo(4070A) produced by CEF expressing either the Mo-MLV Gag-RNase H or
- Mo-MLV Gag-SN polyprotein was significantly inhibited, although the levels of infectious virus slowly increased over the course of the experiment. Mo(4070A) virus production was also monitored by the appearance of the Mo-MLV CA protein (about 30 kDa) on immunoblots. The levels of expression of the Mo-MLV Gag, Mo-MLV Gag-RNase H, and Mo-MLV Gag-SN polyproteins
- remained relatively constant throughout the experiment. At late time points

 15 post-Mo(4070A) infection (days 12 to 14), the Mo-MLV Gag polyprotein levels
 may decrease slightly due to proteolytic processing by protease from the
 infectious Mo(4070A). The levels of the MA protein (about 19 kDa) from the
 RCASBP(A) retroviral vector remained relatively constant throughout the
 experiment.
- The Mo-MLV Gag-SN fusion polyprotein was tested in parallel with the Mo-MLV Gag-RNase H fusion to provide a direct comparison to a characterized CTVI construct known to have antiviral activity. Unexpectedly, the antiviral effect of Mo-MLV Gag-SN was 100-fold greater in the experiments reported herein (> 3,000-fold [Table 1]) than that previously observed (about 30-fold
- [Natsoulis et al., 1995]). The increase in the antiviral effect may be due to an increase in the efficiency of producing CEF cultures that express higher levels of the fusion polyprotein. The CEF themselves grew at a faster rate, presumably due to changes in the growth media including different serum lots. This difference in growth rate was especially apparent in CEF infected with
- 30 Mo(4070A). In previous experiments, Mo(4070A) infection dramatically slowed the growth of CEF and caused some cytotoxicity.

In the experiments reported herein, all CEF cultures grew at the same rate until natural senescence. A consistent six- to eight-fold inhibition of Mo(4070A)

titer was observed in cells expressing the control constructs, including cultures that expressed RCASBP(A) vector (Table 1), that was not detected previously (Natsoulis et al., 1995; Schuman et al., 1996). Although this may be related to the "steric effect" (three- to five-fold) of the Mo-MLV Gag-SN* polyprotein on Mo(4070A) virus production observed by Schumann et al. (1996), the decrease in the Mo(4070A) titers produced in cultures that expressed RCASBP(A) suggests that the RCASBP vector may alter cells in ways that interfere with Mo-MLV virus production, leading to a slight overestimate of the CTVI effect. While several CTVI constructs have significant antiviral effects when used prophylactically, the levels of infectious virus do increase during the experiment. 10 The production of a small number of viruses that do not contain CTVI-derived polyproteins most likely accounts for the small number of infectious viruses needed to spread the infection. Mutations in the gag gene capable of circumventing this antiviral strategy would likely be exceedingly rare since they would require the selection of Gag mutants that would still assemble normally 15 but be able to exclude Gag-degradative enzyme polyproteins from virions.

Delivery of the Mo-MLV Gag-RNase HI Fusion Polyprotein to Mo-MLV-Infected CEF has a Therapeutic Effect. To assess the therapeutic potential of the CTVI strategy, CEF chronically infected with Mo(4070A) were transfected with the RCASBP(A) retroviral vector plasmids containing the Mo-20 MLV Gag fusions and passaged for 16 days. For the therapy assay, CEF chronically infected with Mo-MLV were generated by infecting 4×10^6 cells with 2 x 10⁶ FFU of Mo(4070A) (1 ml of virus stock) and passaging the culture four to six times to allow maximum spread of the virus. The Mo(4070A)infected CEF cultures produced a maximum titer of 3×10^5 to 4×10^5 FFU/ml as 25 quantitated by the S+L- focus assay. The RCASBP virus spread through Mo(4070A)-infected CEFs with kinetics similar to the spread on CEF that had not been infected with Mo(4070A), reaching maximum levels after about 8 days as determined by Western analysis of ALV MA levels. The levels of infectious 30 Mo(4070A) were quantitated throughout the experiment. The Mo(4070A) titers at 14 days posttransfection in two experiments are shown in Table 2. The level of infectious Mo(4070A) was reduced 7.5- to 18-fold in CEF expressing the Mo-MLV Gag-RNase H polyprotein and 15- to 38-fold in CEF expressing the Mo-

MLV Gag-SN polyprotein. Infection with the RCASBP(A) vector alone or RCASBP(A) expressing the Mo-MLV Gag polyprotein lowered the titer of Mo(4070A) produced by a chronically infected culture two- to three-fold compared to Mo-MLV-infected CEF. This suggests that only a slight therapeutic effect is directly attributable to the CTVI constructs once the infection is established.

Table 2. Inhibition of Infectious Mo-MLV Produced after CTVI Treatment^a

		Titer (FFU/ml) ^b (fold inhibition) ^c		
10	Construct used for treatment	Expt 1	Expt 2	
	None (mock treatment)	3×10^{5}	3.8×10^{5}	
	Vector alone	$1.5 \times 10^5 (2)$	ND^d	
	Mo-MLV Gag	$1 \times 10^5 (3)$	$1.6 \times 10^{5} (2.4)$	
	Mo-MLV Gag-RNase H	$4 \times 10^4 (7.5)$	$2.1 \times 10^4 (18)$	
15	Mo-MLV Gag-SN	$2 \times 10^4 (15)$	$1.3 \times 10^4 (29)$	

^a In a therapy assay, CEF chronically infected with Mo(4070A) were transfected with plasmids containing the RCASBP(A) retroviral vector-CTVI constructs. Levels of infectious Mo-MLV in supernatants were quantitated 14 days posttransfection by S+L- focus assays. The experiment was performed twice.

^d ND, not done.

20

25

30

35

Cleavage of the Mo-MLV Gag-Nuclease Fusion Junction by the Incoming Mo-MLV Protease. An Mo-MLV Gag polyprotein fusion junction was designed that could be cleaved by the protease of an incoming infectious Mo-MLV by including the region six amino acids on either side of the normal Mo-MLV protease cleavage site (PRCS). The new Gag fusion plasmid moves the BamHI site to a position just downstream of the Gag termination codon (pMGagPRCS). The termination codon was changed to a codon for glutamine, the same amino acid that Mo-MLV inserts during suppression of the stop codon when the Gag-Pol polyprotein is produced. Genes encoding the Mo-MLV Gag PRCS-RNase H and Mo-MLV Gag PRCS-SN polyproteins were prepared to determine if cleavage of the RNase HI or SN domain from the polyprotein would

^b Titers were averages of duplicate assays.

^c Fold inhibition was determined by comparing the mo-MLV tiers of the experimental constructs to the Mo-MLV titer on mock-treated CEF.

increase the antiviral effect. A prophylaxis assay was used to measure

Mo(4070A) production in cells expressing the Mo-MLV Gag PRCS fusions and the original Mo-MLV Gag fusions. Addition of the PRCS did not alter the level of inhibition by the RNase HI or SN polyprotein, nor did inclusion of the PRCS alter the therapeutic antiviral effect of either the Mo-MLV Gag-RNase H or the SN polyprotein. Virion proteins isolated from day 16 of the therapy assay of Mo-MLV-infected CEF and CEF expressing Mo-MLV Gag, Mo-MLV Gag-RNase H, and Mo-MLV Gag PRCS-RNase H were assayed for RNase H activity by *in situ* RNase H assay. Only virions from Mo-MLV Gag-RNase H and Mo-

- 10 MLV Gag PRCS-RNase H cultures contained measurable RNase H activity. A new protein with RNase H activity about 22 kDa was observed in virions obtained from cultures expressing the Mo-MLV Gag PRCS-RNase H polyprotein. Presumably, this protein, which corresponds in size to *E. coli* RNase HI, is the RNase HI domain cleaved from the Mo-MLV Gag PRCS-
- 15 RNase H polyprotein by the Mo-MLV protease. The ability to release a degradative enzyme from the polyprotein should make it possible to design

future CTVI fusions in which activation of the degradative enzyme is controlled by protease cleavage (e.g., a zymogen), which would allow the use of degradative enzymes whose expression would otherwise be too toxic for the cell.

CTVI Inhibition of Mo-MLV Replication in Avian DF-1 Cells. Line 0 CEF have been the cells of choice for the study of ALV, since line 0 does not 5 contain endogenous loci related to ALV and produces high ALV titers. However, CEF are primary cells and have a limited life span (i.e., about 25 passages). Recently, a permanent cell line was derived from line 0 CEF, named DF-1 (Dr. Douglas Foster, University of Minnesota). Previously, permanent 10 avian cell lines have been unsatisfactory for many ALV studies due to the poor growth of ALV or the presence of interfering exogenous and/or endogenous viruses. The growth of ALV-based retroviral vectors and Mo-MLV was compared in DF-1 and CEF. The DF-1 and CEF cells grow at similar rates and support vigorous growth of ALV and Mo-MLV. The rate at which both ALV and Mo-MLV reach maximum titers, and the level of viral protein produced, is 15 approximately the same in either cell type. The titer of the ALV-based retroviral vectors is 5-10-fold higher when produced in DF-1 cells compared to CEF. A slight increase in the titer of Mo(4070A) viral stocks was obtained when grown in DF-1 cells compared to CEF. These data demonstrate that the DF-1 permanent cell line can be used to test antiviral strategies in vitro, and also to 20 generate clonal cell lines that support vigorous ALV replication. Table 3 summarizes the CTVI antiviral effects on Mo-MLV replication in DF-1 cells both prophylaxis and therapy assays.

In animal model systems, it is possible to deliver genes by using either DNA microinjection or retroviral infection into an embryo. This eliminates the host's immune response, and such methods can be used to deliver genes efficiently. Thus, the CTVI strategy may be useful *in vivo*, e.g., in animal systems in which viral infections have important economic consequences (e.g., agriculture), or in non-human transplant tissues.

Table 3. CTVI inhibition of Mo-MLV replication in avian DF-1 cells

	Titer	Titer (FFU/ml) ^a (fold inhibition) ^b			
Construct	Prophylaxis Assay ^c	Therapy Assay ^d			
		1st Delivery ^e	2nd Delivery ^f		
None (mock)	3×10^5	2 × 10 ⁵	2.1×10^{5}		
Mo-MLV Gag	$1.2 \times 10^5 (2.5)$	$1.4 \times 10^5 (1.4)$	$1.4 \times 10^5 (1.5)$		
Mo-MLV Gag-R	cH <10 (>30,000)	$1 \times 10^2 \ (2,000)$	<10 (>21,000)		
Mo-MLV Gag-S	SN <10 (>30,000)	$1 \times 10^2 (2,000)$	<10 (>21,000)		

10

15

20

25

Example 2

Human Tropic PERVs Produced by Pig Primary PBMC

Methods

30 <u>Cells and Cell Culture</u>. The cell lines used were either obtained from the ATCC unless otherwise indicated. 293 human embryonic kidney (ATCC CRL-1573), MMK *Mus musculus molossinus* kidney (ATCC CRL-6439), SC-1 mouse embryo epithelial cells (ATCC CRL-6450), ST-IOWA (pig fetal testis), 10 PT (pig testis), MDTF *Mus dunni* tail fibroblast (kindly provided by Olivier Danos), NRK normal rat kidney cells (ATCC CRL-6509), Rat2 rat embryo fibroblasts

(ATCC CRL-1764), SIRC rabbit corneal fibroblast (ATCC CCL-60), D17

^a Titers were averages of duplicate assays.

^b Fold inhibition was determined by comparing the Mo(4070A) titers of the construct to mock-treated DF-1 cells.

 ^c DF-1 cells chronically infected with RCASBP(A)-CTVI construct were challenged with Mo(4070A) (multiplicity of infection, 0.05-0.1 FFU per cell). Infectious Mo(4070A) levels were quantitated from supernatants by S+L- assay 30 days post-infection.

^d DF-1 cells chronically infected with Mo(4070A) were infected with RACSBP(A)-CTVI or RCASBP(C)-CTVI construct viruses. The levels of infectious Mo(4070A) in supernatants were quantitated by S+L- assay 30 days post-infection.

^e RCASBP(A)-CTVI viruses were used to initiate the therapy assay with the maximum antiviral effect achieved at 30 days.

f The antiviral constructs were delivered a second time by the RCASBP(C) vectors.

canine osteosarcoma (ATCC CRL-6248), MDBK bovine kidney epithelium (ATCC CCL-22), FRHK4 rhesus monkey kidney cells (ATCC CRL-1688), Mv1Lu mink lung epithelial cells (ATCC CCL-64), MiCL.1 (S+L-) mink lung (ATCC CCL-64.1), Fc3TG feline tongue fibroblast (ATCC CCL 176), AK-D feline lung epithelial cells (ATCC CCL-150), CRFK feline kidney epithelial cells (ATCC CCL-94), CaKi-1 clear cell kidney carcinoma (ATCC HTB-46), HeLa cervical adenocarcinoma (ATCC CCL-2), HOS human osteosarcoma (ATCC CRL-1543), CaCo-2 colorectal adenocarcinoma (ATCC HTB-37) were maintained in Dulbecco's modified Eagle's medium supplemented with 10% 10 fetal bovine serum (FBS), 2 mM glutamine, 1 mM sodium pyruvate, 100 U of penicillin per m1, and 100 µg of streptomycin per ml. PG-4 feline glial and astrocytes (ATCC CRL 2032) were maintained in McCoy's 5A medium supplemented with 15% FBS, 2 mM glutamine, 1 mM sodium pyruvate, 100 U of penicillin per ml, and 100 μg of streptomycin per ml. HepG-2 hepatoblastoma (ATCC HB-8065) and HT1080 fibrosarcoma (ATCC CCL-121) 15 were maintained in Eagle's Modified Essential Medium supplemented in a similar manner. The following cell lines were maintained in RPMI 1640 supplemented with 10% FBS, 2 mM glutamine, 0.05 mg gentamicin sulfate per ml, 1X non-essential amino acids (Biofluids, Rockville, MD): Molt 4 acute 20 lymphoblastic (ATCC CRL-1582), Daudi Burkitt's lymphoma (ATCC CCL-213), Raji Burkitts lymphoma (ATCC CCL-86), U937 histiocytic lymphoma (ATCC CRL 1593.2) UCLA-SO-M14 T cells (Ramsdell et al., 1988), and the human natural killer cell line YTN10 (Yodoi et al., 1985). PBMCs were obtained from NIH and Yucatan minipigs.

Primary pig aortic endothelial cells are isolated by incubating the inner layer of pig aortas with 0.2% collagenase type I for 20 minutes at 37°C.

Detached cells are washed with phosphate-buffered saline, and cultured in M199 medium supplemented with 10% fetal bovine serum, 25 mM HEPES, 2 mM glutamine, 100 U per ml of penicillin, and 100 µg streptomycin. Cultures are fed three times per week and subpassaged every 4-7 days.

Pig hepatocytes are isolated by a two-step perfusion technique as described by Sielaff et al. (1995). The liver is treated by systemic heparinization before isolation. The liver is then treated with an EDTA and collagenase D

solution to free the hepatocytes. The hepatocytes are isolated by gently combing the liver, washing the dislodged cells in Williams E medium supplemented with 5% calf serum, 2 mM L-glutamine, 15 mM HEPES, 1.5 mg/ml insulin, 100 U penicillin per ml and 100 µg streptomycin per ml. The freshly harvested hepatocytes are cultured in collagen gel.

Derivation of viral pseudotypes. Cells productively infected with wild-type PERV were superinfected with retroviral vector-containing supernatant harvested form PA317/G1BgSvN (McLaughlin et al., 1993), in a manner similar to that previously described (Leverett et al., 1998), to generate PERV pseudotypes carrying the murine retrovirus genome coated by PERV core and envelope proteins. G1BgSvN-containing cells were selected in 187 μ g/ml G418 (active component). After 10-14 days selection, G418-resistant colonies were pooled and used to generate viral pseudotypes. Since the G1BgSvN vector genome contains the coding sequence for both neomycin phosphotransferase and for β -galactosidase, pseudotype infection is monitored by the acquisition of target cell resistance to G418 or by immunohistochemical detection of target cells expressing β -galactosidase.

10

15

Infectivity assays. Supernatant infections were used to determine the infectivity of cell-free virus derived from different producer cell lines. Comparisons between fresh and frozen virus-containing supernatants 20 demonstrated that storage of viral supernatants at -70° C resulted in > 100-fold decrease in infectious titer. Therefore, all experiments are performed using freshly harvested supernatant. On the day prior to initiating an infectivity assay, target cells were seeded at 3-5 x 10⁴ cells per well in 12-well dishes. On the next day, supernatant from post-confluent cultures of virus producer cells was 25 harvested, filtered through a 0.45 μm filter, adjusted to 6 μg/m1 polybrene and used to replace the medium on target cells. On the following day, the cells were fed with fresh medium. When β -galactosidase expression was used to monitor infectivity, the cells were fixed and histochemically stained 48-72 hours postexposure to viral supernatant and observed microscopically to enumerate blue-30 staining foci of cells (BFU/m1), as described elsewhere (Wilson et al., 1991). If the experiment was performed to assess productive infection of the target cell,

PCT/US00/14296 WO 00/71726

5

10

15

20

the cells would be passaged 1-2 times per week as needed during the course of the experiment and monitored as described below.

The coculture of target cells with virus-producer cells was used to assess the ability of the virus to be transmitted to non-adherent cells. Since several different cell types were used as a source of virus-producer cells in these experiments, the dosage of irradiation required to ensure the virus-producer cells would die within 5 days after irradiation was first determined. Three to four x 106 cells for each virus-producer cell line were subjected to 2,000,5,000 or 10,000 rads (using a 137Cs source, Nordion GammaCell 100) and seeded into a 96-well plate at 2-3 x 10⁵ cells/well. Irradiated and control non-irradiated cells were monitored for proliferation daily for 5 days by measuring [3H]TTP uptake after cells were cultured for 8 h with 1.0 µCi [3H]TTP (6.7 Ci/mmol; Dupont NEN, Boston, MA), harvested (Skatron, Sterling, VA) and processed for scintillation counting. Based on these findings, the following conditions were used: Daudi cells received 2000 rad, Molt4 and U937 received 5000 rads.

When the target cells in the coculture were human peripheral blood mononuclear cells (hPBMC), the mononuclear layer was collected from a buffy coat from a human donor separated by lymphocyte separation medium (Organon Teknika, Durham, NC). Cells were stimulated in 1X PHA (according to manufacturer's instructions, Life Technologies, Gaithersburg, MD) for 3 days and then mixed in a 1:1 ratio with 3 x 106 lethally irradiated virus-producer cells in 12-well dishes. Thereafter, the cocultures were maintained in RPMI 1640 medium supplemented with 10% FBS, 10% purified human IL-2 (Boerhinger Manheim), 10 ng/m1 recombinant human IL-2 (Chiron Corporation, Emeryville, CA) 2 mM glutamine, 100 U penicillin, and 100 µg per m1 of streptomycin. 25 Control wells containing irradiated virus-producer cells only were monitored for cell viability by visualizing samples microscopically for trypan blue exclusion. Cocultures containing 293 cells as target cells were carried out in parallel as a control for the infectivity of the virus-producer cells. After initiation of the coculture, the next day the medium was replaced with fresh medium and cells 30 were sub-cultured 1-2 times/week as needed. Cocultures containing the hPBMC as targets were reinfused every two weeks with freshly-stimulated hPBMC from the same buffy coat of cells. Methods to monitor virus infection of target cells.

Infection of target cells was assessed by one of three different methods. In some cases, the titer of infectious virus pseudotypes was determined by β -galactosidase expression. Cells were histochemically stained 48-72 hours after exposure to virus-containing supernatant, and foci of β -galactosidase expressing blue cells were counted (BFU/ml). Those titers were then normalized to the titer observed on 293 cells exposed to the same viral supernatant and are reported as % of control. The titer observed on 293 cells ranged 10-fold from several hundred BFU/ml to 2-3,000 BFU/ml.

Virion isolation and purification. Culture supernatants are harvested

from infected cells, cleared of floating cells and debris by centrifugation at 3000 × g for 10 minutes, and the virus stocks stored at -80°C. Virions are isolated for RT assays and Western analysis by ultracentrifugation of 10 ml virus stock underlaid with 1 ml 20% sucrose at 35,000 rpm in a Beckman SW40 rotor for 1 hour at 4°C to pellet the virus (Federspiel et al., 1994). The viral pellet is then resuspended in either RT assay buffer or Laemmli load buffer.

PERV virions are further purified for antibody production either by sucrose equilibrium density centrifugation or by a sucrose step gradient to concentrate the virions and remove proteins, nucleic acids and ribosomes from the cell supernatants. For equilibrium gradients, the supernatants are mixed with sucrose to a density slightly higher than 1.16 g/ml, the average buoyant density of retroviral particles, and the gradient formed by ultracentrifugation. Viral supernatants are layered onto a two-step sucrose gradient, 20% and 50%, and the virions banded at the boundary between the sucrose steps. In both methods, the banded virions are harvested by removing the band with a syringe through the side of the tube to prevent contamination.

20

25

30

PCR. Standard PCR reactions contains 1.25 μ l of 10X PCR buffer (50 mM Tris, pH 8.3, 50 mM KCl, 7 mM MgCl₂), 1.25 μ l of 1.7 mg/ml BSA, 0.5 μ l of each dNTP at 25 mM, 0.56 μ l of each oligonucleotide primer (A₂₆₀ = 5), 0-8% dimethyl sulfoxide, 4.2 μ l water, and 1 μ l genomic DNA (about 100 μ g/ml) (Federspiel et al., 1996). The reactions are heated to 90°C and initiated with 1.5 μ l Taq DNA polymerase (0.75 units). Thirty cycles of PCR are carried out as follows: 90°C for 20 seconds, then 59°C for 80 seconds. Alternations of this basic protocol follow the recommendations of the polymerase manufacturer.

Three sets of primers specifically amplify the three PERV envelope

subgroups in the SU glycoprotein region: (all primers listed 5'-3') PERV-A: TGGAAAGATTGGCAACAGCG and AGTGATGTTAGGCTCAGTGG (SEQ ID NO:8 and SEQ ID NO:9, respectively); PERV-B:

TTCTCCTTTGTCAATTCCGG and TACTTTATCGGGTCCCACTG (SEQ ID NO:10 and SEQ ID NO:11, respectively); PERV-C:
CTGACCTGGATTAGAACTGG and ATGTTAGAGGATGGTCCTGG (SEQ ID NO:12 and SEQ ID NO:13, respectively). PCR products were separated on a 1.0% agarose gel and DNA fragments of 2-2.5 kB were purified using Qiaex gel purification kit (Qiagen, Valencia, CA). Purified PCR products were then cloned into the TA vector using the TA Cloning Kit (In Vitrogen, Carlsbad, CA). Restriction analysis of clones derived from the TA vector was used to ensure an insert of the appropriate size was present. A minimum of 4 representative clones was chosen for sequence analysis. All deoxynucleotide sequencing was performed by the Mayo Clinic Molecular Biology Core on a Perkin Elmer ABIPRISM™ 377 DNA Sequencer (with XL Upgrade) with the PE applied

ABIPRISM 377 DNA Sequencer (with XL Opgrade) with the PE applied
Biosystems ABI PRISM dRhodamine Terminator Cycle Sequencing Ready
Reaction Kit with AmpliTaq DNA Polymerase (PE Applied Biosystems, Foster
City, CA). The ClustalW Multiple Sequence alignment Program of Mac Vector
6.0.1 (Oxford molecular Group) was used to compare the deduced envelope
amino acid sequences.

RT-PCR. RNA is isolated from virions with RNA STAT 50-LS or STAT-60 (Tel-Test, Inc., Friendswood, TX), and converted to cDNA with 2.5 μ M random hexanucleotide primers and Superscript II (Life Technologies,

- Gaithersburg, MD) (Wilson et al., 1998). cDNA templates are amplified with primers corresponding to the PERV *pol/env* gene boundary (ACCTCGAGACTCGGTGGAAGGG; SEQ ID NO:14) and the untranslated region 3' of the PERV *env* gene (CTGGGTTCTGGGAGGGTTAGGTTG; SEQ ID NO:15), or amplified with PB906 (5'
- ACGTACTGGAGGAGGGTCACCTGA 3'; SEQ ID NO:16) and PB908 ('5
 GTCCCGAACCCTTATAACCTCTTG 3'; SEQ ID NO:17) or PERVenv1
 (sense) (5' ACCTCGAGACTCGGTGGAG; SEQ ID NO:11) and PERVenv2
 (anti-sense) (5' CTGGGTTCTGGGAGGGTTAGGTTG; SEQ ID NO:12) for

30 cycles at 94°C for 30 seconds, at 60°C for 30 seconds, and at 72°C for 1 minute. The amplified products are cloned into the PCRII T-A vector (Invitrogen Corp., San Diego, CA).

RT assays. The RT assays are conducted as described in Wilson et al. (1994) and Wilson et al. (1998) and are optimized for the detection of PERV RT 5 as reported by Phan-Thanh et al. (1992). Pelleted virions or virions in cleared supernatant are treated in solubilization buffer (25 mM NaCl, 0.20% Triton X-100, 10 mM Tris, pH 7.5) for 15 minutes at ambient temperature. The solubilized samples are incubated for 3 hours at 37°C in 2X substrate buffer (50 mM Tris, pH 7.5, 10 mM dithiothreitol, 0.6 mM MnCl₂, 10 µg of poly (rA)-poly 10 $(dT)_{12-18}$ (Pharmacia, Piscataway, NJ) per ml, and 10 μ Ci of $^3H[TTP]$ per 25 μ l (22 Ci/mmol). The incubation is terminated with EDTA followed by blotting the synthesized DNA on a DEAE filtermat (LKB-Wallac, Gaithersburg, MD) through use of a harvesting apparatus (Skatron, Sterling, VA) with a rinse of 1% Na₂HPO₄ (pH 7.0) containing 1 mM EDTA. Radioactivity is then determined 15 via liquid scintillation. All samples are assayed in triplicate.

RNase protection assays (RPA). RNA from viral particles or cells is extracted using RNAzolTM B (Tel-Test, Inc., Friendswood, TX). RPA are done with the RPA II Ribonuclease Protection Assay Kit (Ambion, Inc., Austin, TX).

20 Antisense riboprobes are synthesized in the presence ³²P[UTP] using the RNA Transcription Kit (Stratagene, La Jolla, CA). In standard RPA conditions, labeled riboprobe (5 × 10⁵ cpm) is incubated with the RNA at 50°C overnight. The annealed reaction is digested with an RNase A/T1 mix at 30°C for 45 minutes. The samples are cleaned up and fractionated on 6% acrylamide-7.6 M urea gels, dried and exposed to Kodak X-OMAT film.

Western analysis. Virion proteins were isolated and separated on 12% SDS polyacrylamide gels. The proteins were transferred to nitrocellulose filters with an Enprotech blotting apparatus (Integrated Separation Systems, Hyde Park, MA). The filters were blocked in phosphate-buffered saline with 10% nonfat dry milk (wt/vol) for 1 hour at ambient temperature. The blocked filters were drained and incubated with a 1:1000 of goat anti-GALV CA or goat anti-SSAV CA antisera in rinse buffer (100 mM NaCl, 10 mM Tris, pH 8, 1 mM EDTA, 0.1% Tween 20) containing 1% nonfat dry milk for 1 hour at ambient

30

temperature (Biological Carcinogenesis Branch Repository of the National Cancer Institute). The filters were then washed extensively in rinse buffer, and incubated with 50 ng/ml peroxidase-labeled rabbit anti-goat IgG (H + L) (Kirkegaard and Perry, Gaithersburg, MD) in rinse buffer with 1% nonfat dry milk for 1 hour at ambient temperature. The filters were washed and the protein-antibody-peroxidase complexes detected with the Western Blot Chemiluminescence Reagent (DuPont NEN, Boston, MA). The filters were exposed to Kodak X-OMAT film.

Results

15

20

30

10 Host Range of PERV Isolates

Primary PBMC were isolated from an NIH miniature pig and exposed to two different combinations of mitogens: phytohemagglutinin (PHA) and phorbol myristate acetate (PMA) or PMA and the calcium ionophore A23187. Mitogenic activation by either treatment resulted in a sharp increase in reverse transcriptase (RT) activity levels five days after stimulation (Figure 1A). To determine if the increase in RT activity correlated with the production of infectious virus, stimulated PBMC from NIH and Yucatan minipigs were cocultured with the pig ST-IOWA cell line, a line that does not produce significant endogenous RT activity levels in the supernatants but is susceptible to PERV infection. Both cultures produced detectable RT activity 14 days after exposure to activated PBMC (Figure 1B). These results indicated that mitogenic activation of the pig PMBC released PERV capable of productive infection of pig cells. To determine if the produced PERV can infect human cells, stimulated PBMCs were cocultured with the human embryonic kidney 293 cell line. RT activity was detected in 293 cells at day 31 (Figure 1C) and continued to increase to levels comparable to RT activity levels produced by infected ST-IOWA cells. These results clearly demonstrated that the stimulated pig PBMC produce PERV that can productively infect human cells. These findings suggest that the stimulation of porcine lymphocytes, transferred to the patient in a xenograft, may result in the production of replication-competent human-tropic PERV capable of infecting the patient.

In order to test if the PERVs of this primary 293 infection (1°-293/PERV) could improve their infection efficiency in human cells, supernatant

from the 1°-293/PERV culture was passaged into a new culture of human 293 cells (2°-293/PERV) and allowed to spread through the culture (Figure 2). After the culture reached maximum RT activity, supernatant from the 2°-293/PERV culture was passaged into fresh 293 cells and allowed to spread (3°-293/PERV). The titer of the PERVs produced by these three cultures was quantitated by generating MLV- β gal pseudotypes.

The titers of all three PERV cultures were approximately the same on the control ST-IOWA cells (Figure 3). However, the titers of the MLV-βgal/PERV pseudotypes produced from the 2°-293 and 3°-293 cultures were 5-fold higher compared to the 1°-293 culture. The relative copy number of integrated PERV proviruses in these three PERV infected 293 cultures was quantitated by Southern analysis. As shown in Figure 4, the DNA isolated from the 2°-293/PERV and 3°-293/PERV cultures contained more integrated PERV proviruses per cell relative to the DNA of the 1°-293/PERV cultures.

10

15

Overall, the host range of the PERVs produced from the 293/PERV cultures appears to coincide with a PERV-A related isolate (see Table 4 compared to Takeuchi et al., 1998). One notable exception is the inability of PERV-1.15 MLV-βgal pseudotypes to infect D17 cells, suggesting that the changes in the envelope glycoproteins may have altered the host range.

Table 4. Host range of PBMC PERV produced by 293 cells

			Titer of LacZ pseudotype ^a		
	Species	Cell	1°	2°	3°
	Human	HeLa	+	+	+
5		HOS	+	+	+
		Caki-1	-	-	+
		293	+	++	++
	Primate	FRhK-4	-		-
10	Pig	ST-IOWA	++	++	++
		10 PT	+	+	+
	Dog	D17	-	_	_
15	Rabbit	SIRC	-	-	-
	Hamster	E36	-	-	_
20	Mink	Mv-1-Lu	++	++	++
		MiCl1	+	++	++
	Cat	Fc3Tg	+	+	++
		AK-D	+	+	++
25		CRFK	++	++	++
		PG4	++	++	++

 $[^]a$ MLV-βgal pseudotypes were produced in the PERV infected human 293 cultures: negative (-); 1-100 BFU/ml (+); >100 BFU/ml (++).

30

42

Table 5. Productive infection of human and animal cell lines by 293/PERV

	Cell Line	Virusª	$LacZ^{\flat}$	RT°	RT-PCR ^d
	ST	1°	++	neg	nd
5	(pig testis)	2°	++	d .9	nd
		3°	++	d .9	nd
	293	1°	+	d.18	nd
	(human kidney)	2°	++	d.10	nd
	· · · · · · · · · · · · · · · · · · ·	3°	++	d.10	nd
10	Caco-2	1°	?	neg	_
	(human colon)	2°	?	neg	_
	,	3°	?	d.18	_
15	HepG2	1°	++	neg	_
	(human liver)	2°	++	d.26	+
	,	3°	++	d.26	+
	HT1080	1°	+	neg	_
	(human fibrosarcoma)	2°	++	d.26	+
		3°	++ .	d.13	+
20	Mv-1-Lu	1°	+	neg	+
	(mink lung)	2°	++	d.21	+
		3°	++	d.38	+
	CRFK	1°	+	neg	+
	(cat kidney)	2°	+	d.14	+
		3°	++	d.14	+

²⁵

nd Not determined.

35

30

PERV Produced by the 293/PERV Cultures Can Productively Infect Other Human, Mink and Cat Cell Lines. A number of different cell lines were employed to determine the tissue and species specificity of PERV produced by 293/PERV cultures (Table 5). All the cell lines tested except Caco-2 were

productively infected by the virus populations produced by the 2° and 3°-293/PERV cultures. The 3°-293/PERV infected Caco-2 culture produced RT

^a Culture supernatant from 1°, 2°, or 3° -293/PERV cells.

^b Titer of MLV-βgal pseudotype: negative (-); 1-100 BFU/ml (+); >100 BFU/ml (++).

^c The day when the RT activity was greater than 2-fold over background activity from uninfected cells. The assay was terminated after 8 weeks.

d At 8 weeks postinfection, RNA was isolated from the culture supernatant and assayed for *pol* sequences by RT-PCR.

[?] High background staining made the data uninterpretable.

activity at day 18 and detectable *pol* RNA by RT-PCR indicating a productive infection. The 1°-293/PERV culture produces 5-fold lower PERV titers (see Figure 3) which may account for the lower replication efficiency in most lines. The transfer of significant levels of replication-defective PERV genomes into the primary 293 cells may account for the lower titers. The ability of the PERV produced by the 2° and 3°-293/PERV cultures to efficiently replicate in most cultures may be the result of the loss of defective interfering PERV genomes or the adaptation of the virus, or both.

An expanded analysis of species tropism indicated that cell lines derived from mouse, rat, rabbit, dog, cow, and rhesus monkey were resistant to infection by 293/2° pseudotypes, while cell lines from cat or mink cell lines were susceptible. This latter susceptibility was low compared to 293 cells, with the exception of MiCL and PG4 cell lines. Mink and cat cell lines were also infected with PERV infected 293/2° supernatant. Only low levels of RT activity were observed in MiCL and CRFK cultures. PG-4 cultures had RT activity of 933 cpm [³H-TTP] at 3 weeks, which was reduced to 2344 cpm by 5 weeks, post infection.

10

20

25

Analysis of the Susceptibility of Hematopoietic Cells to Infection by PERV. In order to assess whether primary human PBMC (hPBMC) were susceptible to infection by PERV, PHA-activated hPBMC were cocultured with irradiated 293/2° virus-producer cells and maintained in IL-2-containing medium for 8 weeks. No RT activity above background levels or viral RNA as measured by RT-PCR were produced during the course of the experiment. Since maintenance of the hPBMC in IL-2 during the course of the experiment biases the culture conditions towards the proliferation of T cells, other hematopoietic lineages that may be susceptible to infection may not have been represented. To investigate the possibility that other hematopoietic lineages may be permissive for infection, a number of human hematopoietic cell lines representing the T cell, B cell, myeloid, and NK cell lineages were analyzed for their susceptibility to PERV infection by coculture with irradiated 293/2° virus producer cells. Supernatants from the cocultures containing either M14 (T cell lineage), Jurkat (T cell lineage), K562 (myeloid lineage), or the NK cell line YTN10, remained negative for RT activity during the 8 week course of the experiment, although

shows the data from the RT activity for the cocultures containing the T cell line Molt 4, the B cell lines Daudi and Raji, and the myeloid cell line U937. By 4 weeks post-coculture, the RT activity in the Molt 4- and Daudi-containing cocultures was positive and continued progressively to increase throughout the period of the coculture. In contrast, the RT activity for both the Raji and U937 sets of cocultures plateaued by 4 weeks post-culture. The results from this experiment suggest that the Molt 4, Daudi, Raji, and U937 cell lines were permissive for productive infection by PERV.

5

10

15

20

25

The RT positive hematopoietic cell lines were then co-cultured with primary hPBMC, as a virus population may have been selected in the susceptible hematopoietic cell lines that could more efficiently infect primary hematopoietic cells. The irradiation conditions optimal for lethal irradiation were determined for each of the RT positive Daudi, Molt 4 and U937 cell lines. Then lethallyirradiated RT-positive Daudi, Molt 4 and U937 were each employed as virusproducer cells in cocultures with primary hPBMC activated with PHA or with human 293 cells as positive controls. By 2 weeks post-coculture, each of the cocultures containing 293 cells as target cells became significantly RT positive (>7,000 cpm), and by 3 weeks, the RT activity in each of these cultures increased to >20,000 cpm. By contrast, none of the cocultures containing the hPBMC demonstrated RT activity higher than negative control cultures over the course of the experiment. Although viral RNA was detected by RT-PCR in the supernatant of cultures sampled 1-2 weeks post-exposure, the supernatant was negative for viral RNA by RT-PCR by three weeks and remained so out to the 8week time point. The positive RT-PCR results obtained at the early time points most likely reflect presence of residual irradiated virus-producer cells which disappear from the culture at the later time points, rather than infected cells.

Table 6. Comparison of the nucleotide and deduced amino acid sequence of PERV-1.15 with the three known PERV env subgroups.

	1.15	А	В	С
PERV-1.15	_	96	85	73
PERV-A	97	-	82	76
PERV-B	87	84	_	70
PERV-C	72	76	72	-

Nucleotide Identity (%)

Amino Acid Identity (%)

Env Subgroups of PERV in PERV- infected 293 Cultures

5

10

15

20

25

30

To determine what PERV envelope subgroups were present in virions produced by the 293/PERV populations, the envelope regions was cloned by RT-PCR amplification of virion RNA isolated from 1°, 2°, and 3°-293/PERV supernatants. The primers used for RT-PCR were designed from the PERV-A and PERV-B envelope sequences. Oligonucleotide primers were synthesized homologous to a conserved region at the 3′-end of the *pol* gene (5′ of the PERV *env* gene) and a conserved untranslated region 3′ of the *env* gene. The amplified products were cloned into the pCRII T-A cloning vector (Invitrogen) and representative clones sequenced from each culture. Unexpectedly, only one PERV envelope gene species was detected in virions produced from all three cultures. The deduced amino acid sequence shows that the surface glycoprotein region of the gene is almost identical to PERV-A, but the transmembrane glycoprotein region of the gene is almost identical to PERV-C (Figure 6). The nucleotide and amino acid sequence homologies of PERV-1.15 with PERV-A, -B, and -C are summarized in Table 6.

While the PERV-1.15 envelope gene is clearly in the PERV-A receptor subgroup family, the gene contains a unique transmembrane glycoprotein that may be important for efficient PERV replication in human cells. The variation in the PERV env genes may be significant since variant MLV env genes have

5

10

15

20

25

30

been shown to not only broaden the host range of the esotropic MLV but increase the pathogenicity (Rosenberg et al., 1997). These viruses arise by recombination between different endogenous MLVs. Recombination between *env* sequences of exogenous and endogenous feline leukemia viruses (FeLV) also occurs in FeLV-induced thymic tumors.

All 293/PERV cultures contained PERV-A envelope sequences as determined by PCR using subgroup-specific envelope primers. However, PERV-B envelope sequences were not detected in the 293/PERV culture DNAs. The PERV-C envelope sequence was detected in the 1°-293/PERV culture, but not in the subsequent virus passages. PERV-C does not infect most human cells efficiently.

Isolation of Other Human-Tropic PERVs. To identify other human-tropic PERV, primary cell cultures, e.g., pig aortic endothelial cells (PAEC from, for example, NIH and Yucatan minipig breeds), pig PBMC, and primary pig hepatocytes, from normal pig tissues are assayed for the production of human-tropic PERVs on human 293 cells. The cultures are passaged and monitored for PERV production by RT-PCR and RT activity. The PERV populations are passaged onto fresh human cells to generate secondary cultures to determine if a particular PERV species dominates, and if the overall replication efficiency improves. Env subgroups and LTR-U3 sequences are determined as described above.

New preparations of PERV cDNA are synthesized by RT-PCR from virion RNA isolated from PERV sources, and 1°, 2°, and 3°-293/PERV culture supernatants. The region from the virion mRNA polyA tail to the PERV pol gene is amplified by RT-PCR. The RT-PCR reactions use a pol primer that hybridizes just 5′ of the env gene, and a second reaction using a pol primer homologous to the RT active site region. Two reactions that use different pol primers increase the chance of amplifying all PERV variants. ProSTAR Ultra High Fidelity (HF) RT-PCR System (Stratagene) is employed for RT-PCR as it combines high fidelity with long target amplification using the PfuTurbo DNA polymerase. PfuTurbo DNA polymerase efficiently generates blunt ends. The product is cloned into PCR-Script (Stratagene) designed for blunt-end cloning. This approach does not rely on specific restriction sites to be present in the

PERV sequence, and produces cDNA clones that contain both the envelope gene and the U3 LTR region.

The resulting bacterial colonies are grouped first by envelope subgroups by probing colony lifts on nitrocellulose with ³²P-labeled DNA probes specific for each PERV envelope subgroup. These regions only detect the variable region of the envelope surface glycoprotein. The probing colony lifts with a specific ³²P-labeled probe (DNA region or oligonucleotide), by PCR with specific primers designed to identify the variant sequence, and/or by Southern blots of digested plasmid clone DNA probed with a specific ³²P-labeled probe. 10 In this way, thousands of colonies are easily probed for specific PERV sequences, thereby increasing the detection of rare PERV RNAs. For example, PERV-B was not detected in the DNA in any of the 293/PERV cultures, and PERV-C was barely detected in the DNA of 1°-293/PERV. Thus, PERV-B and PERV-C envelope sequences may be represented in virions, especially in virions produced by PBMC and the 1°-293/PERV culture supernatants. RT-PCR 15 reactions designed to amplify only PERV-B and PERV-C mRNAs are also done. Finally, the expression profile of PERV envelope and U3-LTR sequences in PBMC is analyzed by RNAse protection assays. Specific PERV envelope and U3-LTR sequences are cloned into pBluescript KS (Stratagene), and used to produce ³²P-labeled antisense RNA probes. Pig and human GPDH probes are 20 also used to normalize RNA levels in the assays for quantitation of PERV RNA levels in PBMC. These assays determine if PERV-B and PERV-C loci are expressed in PBMC. The nucleotide sequence is then determined for 10-15

Cellular DNA is probed, by Southern analysis and PCR, for the presence of any new or variant PERV sequence identified by these experiments. This analysis verifies that the PERV sequence is present in the genome, and not an artifact of the cloning or analysis procedure, although the sequence could be the result of adaptation. The biological properties of all PERV sequences is characterized in an infectious molecular clone. RNase protection assays are used to identify and quantitate the expression levels of the unique PERV elements in RNA isolated from a variety of tissues: brain, pancreas, spleen, liver, heart, kidney, muscle, lung, and PBMC.

representative clones of each unique PERV group.

25

30

Example 3

Molecular Cloning and Characterization of PERVs

A lambda genomic library was prepared from Sau3A-partially digested DNA isolated from 2°-293/PERV cells in the Lambda Dash II vector

(Stratagene). The library was probed with a ³²P-labeled PERV pol gene fragment, and ten genomic clones isolated. Six of the clones were unique and contained partial PERV proviruses.

10

15

All of the cloned PERV regions were nearly identical, with at most two amino acid changes in the coding regions, except the PERV sequences in lamA8 (Figure 7). The lamA8 pol gene contains an 86 nucleotide deletion that results in the truncation of the open reading frame at amino acid 887, a loss of 308 amino acids. The lamA8 env gene contains 101 nucleotide insertion that results in the truncation of the open reading frame at amino acid 599, a loss of 60 amino acids which includes the putative transmembrane region of the transmembrane glycoprotein. It is unlikely, therefore, that lamA8 contains a functional pol or env gene.

The PERV regions cloned from the 2°-293/PERV culture were compared to the putative complete PERV cDNA, PERV-MSL (Akiyoshi et al., 1998), the PERV-A (Letissier et al., 1997) and PERV-1.15 envelope genes, and gibbon ape leukemia virus (GALV) (Table 7). The 293/PERV gag and pol genes, as well as the 5' and 3' untranslated regions were nearly identical to the PERV-MSL nucleotide and deduced amino acid sequence.

PCT/US00/14296 WO 00/71726

Table 7. Comparison of the 2°-293/PERV regions contained within the lambda clones with other PERV isolates

			Percent Identity			
	Region	Sequence ^a	PERV- MSL ^b	PERV-A	PERV- 1.15	GALV°
5	LTR	N	78			
	5'-ut	N	97			
	gag	N	98.9			
		Α	98.5			62.6
	pol	N	99			
С		A	99			68
	env	N	86.7	97	99.8	
		Α	85	96	99.7	47
	3'-ut	N	100			

^a N, nucleotide sequence; A, amino acid sequence.

^c GALV, gibbon ape leukemia virus. ut untranslated region.

20

30

15

The env gene in lamA6 has 4 nucleotide and 2 amino acid differences with PERV-1.15, the 293/PERV env gene cloned by RT-PCR from virion RNA (Figure 18). Upon comparing the nucleotide sequence of the 293/PERV and PERV-MSL LTRs, a large divergence was observed in the U3 region (Figure 8). 25 The 293/PERV LTR contains 71 more nucleotides than the PERV-MSL LTR. The sequence of the PERV sequences in Lambda clones A1, A11, A3A, A10, and A8 is also shown in Figure 18.

Two recombinant PERV proviruses were prepared to test if they produce infectious virus in 293 cells. The constructs, an A3A/A6 chimera and an A1/A6 chimera, are outlined in Figure 9. All constructs were built in the pBluescript KS vector (Stratagene), and used the unique BspEI site in the pol gene to link the 3'-half of the genome contained in lam A6 to the 5'-half of the genome contained in either lamA1 or lamA3A. A minimum of genomic sequence was cloned with the PERV region: A1, about 1000 bp; A3A, about 30 bp; A6, about 35 700 bp.

b PERV-MSL, a putative full-length PERV cDNA clone with the C-type envelope gene.

Example 4

Preparation and Detection of Anti-PERV Antibodies

Rabbit polyclonal antisera is produced against 2°-293/PERV virion proteins and against 2°-293/PERV capsid proteins. The 2°-293/PERV cultures produce a relatively large amount of viral particles as demonstrated in Figure 10A. Goat anti-SSAV CA and goat anti-GALV CA antisera easily detect PERV Gag proteins on a Western blot with only minor cross-reaction to control 293 cell supernatant proteins (Figure 10B). These antisera are used as positive controls for the characterization of the anti-PERV antisera and monoclonal antibodies. The 2°-293/PERV cultures produce approximately equal amounts of the p30 capsid (CA) protein in 5 ml of supernatant as ALV infected cultures produce ALV p27 CA protein (lanes 2 and 5). The high-titer ALV stock contained 10⁷ iu/ml. The 2°-293/PERV virions are collected and purified by equilibrium density centrifugation on sucrose gradients. The banded virions are collected, dialyzed, and used for antibody production. The PERV CA protein is isolated from SDS-PAGE gels (Figure 10A) and also used for polyclonal antisera production.

10

15

20

25

30

Two rabbits are used for each antigen preparation. The antigen (100 µg) mixed in Complete Freund's Adjuvant is used for the initial inoculation. The animal is boosted with 50 µg antigen in Incomplete Freund's Adjuvant on day 14, 21 and 49 after the initial inoculation. The route of injection is normally subcutaneous and/or intramuscular at multiple sites. Test bleeds are drawn on day 35 and 56 and screened against PERV proteins by Western blot and immunoprecipitation assays.

Purified PERV CA protein is used to produce mouse hybridomas that secrete monoclonal antibodies against PERV CA. The PERV capsid protein, purified from polyacrylamide gels, is used as the antigen. Five Balb/c mice are injected intraperitoneally or subcutaneously with 400 µg antigen (per mouse) in Complete Freund's Adjuvant and boosted two weeks later with 250-500 µg of the antigen in Incomplete Freund's Adjuvant. All five mice are then tested for the presence of PERV anti-capsid antibodies by Western blot. One of the positive mice is used for hybridoma preparation. Spleenocytes and/or lymph node cells are removed and fused to non-immunoglobulin secreting myeloma

5

10

15

20

25

30

cell line F/O. Hybridomas are screened and positive hybrids are cloned and then subcloned to obtain monoclonal antibody (MoAb) producing cell lines. The hybridomas are screened by both Western blot and immunoprecipitation of PERV capsid protein. The MoAb cell lines are isotyped and cryopreserved. MoAbs are purified from the cell line supernatant by protein A/G column chromatography.

The new anti-PERV antibodies, as well as the anti-SSAV and anti-GALV CA antisera, are tested for their ability to detect PERV proteins in live and fixed PERV infected cells. An immunofluorescence and/or immunohistochemical assay for PERV infection simplifies the detection of PERV infection and allows the direct quantitation of PERV titers.

To detect anti-PERV antibodies in sera, a Western-based assay system is employed. Protein preparations made from PERV virions produced by the 2°-293/PERV culture are separated by SDS-PAGE and transferred to nitrocellulose filters. Virion protein extracts are used for two reasons. First, PERV produced by human cells does not display the α-Gal xenoreactive carbohydrate that may bind to the α-Gal antibodies in human sera. Second, the 2°-293/PERV culture produces high levels of PERV virions as shown by analysis of PERV virion protein extracts. Goat anti-SSAV and anti-GALV CA antisera that cross-reacts with the PERV CA protein is employed. The 2°-293/PERV virions are purified by equilibrium centrifugation on sucrose gradients. In one embodiment, the assay employs peroxidase-conjugated secondary antibodies and enhanced chemiluminescence (Amersham, Pharmacia Biotech, Piscataway, NJ) to detect antibodies bound to the PERV proteins. To determine what effect human sera may have on the performance of the assay (e.g., alter the antibody sensitivity or cross-reaction with common human antibodies), the anti-CA antisera is added to dilutions of control human sera (1:10, 1:50, 1:100) and tested.

Serum samples from at least 300 pig slaughterhouse workers, plus controls, are screened. Control samples, including sera from a patient with little or no exposure to pigs, and sera from two populations of retrovirus-infected patient populations (50 HIV-1; 20 HTLV-1) is used to evaluate the specificity of antibody binding to PERV and non-PERV proteins. Preparative single well SDS-PAGE is used to separate the PERV virion proteins, and the proteins

transferred to a nitrocellulose filter. The Miniblotter II apparatus (Immunetics, Cambridge, MA) permits the screening of 45 different samples by dividing the Western blot into 45 sample cells. A negative control serum and a positive control serum (human sera with SSAV or GALV anti-capsid antisera) is assayed with each blot. A 1:50 dilution of the serum samples is screened initially. Human blood samples are also tested for PERV nucleic acid and anti-PERV antibodies.

Example 5

Soluble Forms of the Subgroup A Avian Leukosis Virus Receptor Tva
Significantly Inhibit ALV(A) Infection *in vitro* and *in vivo*

Three cell surface proteins have been identified as ALV receptors: Tva, the receptor for ALV(A) (Bates et al., 1998; Bates et al., 1993; Young et al., 1993); CAR1, the receptor for ALV(B) and ALV(D) (Brojatsch et al., 1996; Smith et al., 1998); and SEAR, the receptor for ALV(E) (Adkins et al., 1997). To aid in the characterization of the interactions between Tva and ALV(A) envelope glycoproteins, soluble forms of the 83 amino acid extracellular domain of the Tva receptor protein (sTva) were constructed by Young and colleagues (Connolly et al., 1995), who reported that pre-incubation of the sTva proteins with different envelope subgroup ALVs caused a specific block to infection of susceptible chicken cells by ALV(A), but had no effect on ALV(B) or ALV(C) infection.

To determine if cells and chickens expressing sTva proteins are resistant to ALV(A) infection, ALV-based replication-competent retroviral vectors were used to efficiently deliver and express stva genes (Federspiel et al., 1997). The vectors are available with five different envelope subgroups (A-E) which enables multiple genes to be delivered and expressed in virtually every cell.

Materials and Methods

15

20

25

30

Soluble receptor and retroviral vector constructs. The two soluble receptor gene constructs, contained in the plasmids pLC126 and pKZ457, were gifts of John Young (Harvard Medical School). The pLC126 stva gene (Connolly et al., 1994; Hughes et al., 1987), encoding the 83-amino-acid Tva extracellular domain fused to a 9-amino-acid antibody epitope tag derived from

influenza virus hemagglutinin, followed by six histidine residues, was isolated as a NcoI-PstI fragment and cloned into the NcoI and PstI sites of the CLA12NCO adaptor plasmid (Federspiel et al., 1997; Hughes et al., 1987). The pKZ457 stvamIgG gene, encoding the 83-amino-acid Tva extracellular domain fused to the constant region of the mouse IgG heavy chain (nucleotides 353-1072) (Tucker et al., 1979), was isolated as a NcoI-BlpI fragment. The BlpI site was made blunt and the modified fragment was cloned into the NcoI and SmaI sites of CLA12NCO. Both the stva and stva-mlgG genes had been modified to contain NcoI sites at their initiator ATGs. The soluble receptor gene cassettes were 10 isolated as ClaI fragments from the adaptor plasmids and cloned into the unique ClaI site of the RCASBP, RCAS, and RCOSBP retroviral vectors with subgroup (B) and subgroup (C) envelope genes. The RCAS family of replicationcompetent retroviral vectors have been described (Federspiel et al., 1994; Federspiel et al., 1997; Hughes et al., 1987; Petropoulous et al., 1991; Petropoulous et al., 1992). 15

The stva-mlgG gene isolated as a ClaI fragment from the CLA12NCO adaptor plasmid was subcloned into the TFANEO expression vector (Federspiel et al., 1989). TFANEO is a companion expression vector to the RCAS family of retroviral vectors. The expression cassette of TFANEO consists of two LTRs derived from the RCAS vector that provide strong promoter, enhancer, and polyadenylation sites flanking a unique ClaI insertion site. The TFANEO plasmid also contains a neo resistance gene expressed under the control of the chicken β-actin promoter, and an ampicillin resistance gene for selection in E. coli.

The RCASBP(A)AP, RCASBP(C)AP and RCASBP(C)AP retroviral vectors which contain the heat-stable human placental alkaline phosphatase gene (AP) have been described (Federspiel et al., 1997; Field-Berry et al., 1992; Fekete et al., 1993). The AP gene, contained on a SalI fragment, was cloned into the SalI site of the CLA12 adaptor plasmid, and then subcloned into the RCASBP vectors as a ClaI fragment (gift of Constance Cepko).

Cell culture and virus propagation. Chicken embryo fibroblasts (CEFs) derived from 10-day, line 0 embryos (C/E) (Astrin et al., 1979) were grown in DMEM (GIBCO/BRL) supplemented with 10% tryptose phosphate broth

(GIBCO/BRL), 5% fetal bovine serum (GIBCO/BRL), 5% newborn calf serum (GIBCO/BRL), 100 units of penicillin per ml, and 100 μg of streptomycin per ml (Quality Biological, Inc., Gaithersburg, MD) as previously described (Federspiel et al., 1997). DF-1 cells were grown in DMEM supplemented with 10% fetal bovine serum, 100 units of penicillin per ml, and 100 μg of streptomycin per ml (Himly et al., 1998; Schaefer-Klein et al., 1998). Both CEF and DF-1 cultures were passages 1:3 when confluent.

Virus propagation was initiated by calcium phosphate transfection of plasmid DNA that contained the retroviral vector in proviral form (Federspiel et al., 1997). In standard transfections, 5 μg of purified plasmid DNA was introduced into DF-1 cells or early passage CEF by the calcium phosphate precipitation method (Kingston et al., 1989). Viral spread was monitored by assaying culture supernatants for ALV capsid protein by either Western transfer analysis or ELISA (Smith et al., 1979). Virus stocks were generated from the cell supernatants. The supernatants were cleared of cellular debris by centrifugation at 2000 × g for 10 minutes at 4°C and stored in aliquots at -80°C. DF-1 cells transfected with the TFANEO plasmid were grown in 500 μg/ml G418 (Gibco/BRL) to select for neomycin-resistant cells. Clones were isolated using cloning cylinders (Bellco Glass Inc., Vineland, NJ), expanded, and maintained with standard medium supplemented with 250 μg/ml G418.

10

15

20

25

30

ALV alkaline phosphatase challenge assay. In a direct AP challenge assay, CEF or DF-1 cell cultures (about 30% confluent) were incubated with 10-fold serial dilutions of the RCASBP/AP virus stocks for 36-48 hours at 39°C. In a pre-absorption AP challenge assay, the 10-fold viral serial dilutions were first mixed with 2 ml of supernatant containing sTva-mIgG for 3 hours at 4°C, and then assayed as above. The assay for alkaline phosphatase activity was modified from procedures of Cepko and co-workers (Federspiel et al., 1994; Fekete et al., 1993; Fields-Berry et al., 1995). Cells were fixed in 4% paraformaldehyde in Dulbecco's phosphate-buffered saline (PBS) for 30 minutes at 25°C, washed twice in PBS for 5 minutes each, and incubated for 1 hour at 65°C to inactivate endogenous AP activity. The cells were then washed twice with AP detection buffer (100 mM Tris•Cl, pH 9.5/100 mM NaCl/50 mM MgCl₂) for 10 minutes and exposed to the AP chromogenic substrates nitroblue tetrazolium (300 μg/ml)

and 5-bromo-4-chloro-3-indoyl phosphate (170 μg/ml) (GIBCO/BRL).

5

30

Enzymatically active AP produces an insoluble purple precipitate. The reaction was stopped by the addition of 20 mM EDTA, pH 8.0 in PBS.

Immunoprecipitation and Western transfer analysis of stva-mIgG proteins. A 500 µl aliquot of culture supernatant or serum was incubated with 50 µl of anti-mouse IgG-agarose beads (Sigma) for ≥ 1 hour at 4°C. The sTvamIgG agarose bead complexes were collected by centrifugation and washed twice in dilution buffer [50 mM Tris-buffered saline (TBS), 1% Triton X-100, 1 mg/ml BSA], once in 50 mM TBS and once in 0.05 M Tris•Cl, pH 6.8. The washed complexes were collected by centrifugation, resuspended in 50 µl 1X Laemmli buffer (2% SDS, 10% glycerol, 0.05 M Tris•Cl, pH 6.8, 0.1% bromophenol blue) without β -mercaptoethanol, and heated for 5 minutes at 100°C. The agarose in the samples was collected by centrifugation for 2 minutes and the supernatants were transferred to new tubes. Prior to gel electrophoresis, $1.0 \mu l$ β -mercaptoethanol was added to each $50 \mu l$ sample and the samples were 15 heated for 5 minutes at 100°C. The denatured immunoprecipitates were separated by 12% SDS-PAGE, and transferred to a nitrocellulose membrane. The filters were blocked with 10% non-fat dry milk (NFDM) in PBS, probed with 0.05 µg/ml peroxidase-conjugated goat anti-mouse IgG antibodies (Kirkegaard and Perry laboratories, Gaithersburg, MD) in rinse buffer (100 mM 20 NaCl, 10 mM Tris•Cl, pH 8, 1 mM EDTA, 0.1% Tween 20) and 1% NFDM, and washed in rinse buffer. Protein/antibody complexes were detected with the Western Blot Chemiluminescence Reagent (NEN) according to the manufacturer's instructions. The immunoblot was then exposed to Kodak X-25 Omat film.

In vivo ALV challenge assay. Line 0 embryos were somatically infected with RCASBP(B), RCASBP(B)stva, or RCASBP(B)stva-mIgG by injecting unincubated eggs near the blastoderm with 100 μ l containing 1 × 10⁶ CEF or DF-1 cells producing the virus. Line 0 is a White Leghorn line that is genetically susceptible to all ALV subgroups except subgroup (E) and is free of endogenous proviruses that are closely related with ALV (Astrin et al., 1979). Viremic chicks were identified at hatch by ELISA for the ALV capsid protein p27. Viremic and uninfected control chicks were infected intra-abdominally

5

10

15

with 10⁵ infectious units of either RAV-1 [an ALV(A) isolate] or RAV-49 [an ALV(C) isolate]. Blood was collected at 2, 4 or 9 weeks post-challenge, and the serum assayed for infectious subgroup (A), (B), or (C) ALV by the *in vitro* ALV Assay (see below).

In vitro ALV assay. The presence of infectious ALV in chickens was determined by assaying the serum samples on a panel of cell lines with different ALV envelope subgroup susceptibilities. The panel of indicator cell lines included: line 0 CEF (C/E) which supports the replication of ALV(A), ALV(B), and ALV(C); line alv6 CEF (C/A) which supports ALV(B) and ALV(C) replication; line RP30B-cell line (C/B) which supports ALV(A) and ALV(C) replication; and line 15.C-12 CEF (C/C) which supports ALV(A) and ALV(B) replication. Serum samples (100 μl) were added to the cells and the cells were incubated for 9 days in media (5% serum) to allow ALV to spread. The media was changed after 3 days to avoid detection of ALV proteins in the original serum sample. The cells were then solubilized by 2 cycles of rapid freeze-thaw to release ALV Gag antigens. The ALV capsid protein was detected by ELISA. A positive sample was defined as having an optical density reading of > 0.200. The *in vitro* ALV Assay can detect infectious ALV titers as low as 10 IFU/ml.

RNase protection assay. Total RNA was isolated from cells in culture, or 20 homogenized tissues of experimental birds, by the RNazol B method (Tel-Test, Inc., Friendswood, TX). Sequence-specific RNA probes were cloned into pBluescript KS as follows: the RAV-1 envelope sequences were cloned as an XhoI to XbaI fragment (Genbank accession # M19113; nucleotides 248-676) (Bora et al., 1988); the RAV-2 envelope sequences were cloned as a BamHI to Sall fragment (Genbank accession # M14902; nucleotides 612-1080) (Bora et 25 al., 1986); the stva probe was generated from an EcoRI to PstI fragment from the plasmid pLC126 (Genbank accession # L22752; nucleotides 9-386) (Bates et al., 1993; Connolly et al., 1994); and the stva-mIgG probe was generated from a ClaI to BamHI fragment derived from the adaptor plasmid construct which contains the 5'-ClaI site and transcription leader from the CLA12NCO adaptor 30 plasmid and a synthetic sequence encoding the 83-amino acid Tva extracellular domain from pKZ457 that is different from the stva gene. A fragment of the chicken glyceraldehyde-3-phosphate dehydrogenase (GAPDH) gene (Genbank

accession # K01458; nucleotides 163-361) (Panabieres et al., 1984) was used as a control for the quantity and quality of the RNA. The constructs were linearized by restriction endonuclease digestion and gel purified. ³²P-labeled antisense RNA probes were synthesized using the RNA Transcription Kit

5 (Stratagene, La Jolla, CA). The probes were hybridized with 20 μg of total RNA in 20 μl hybridization solution (80% formamide, 10 mM sodium citrate pH 6.4, 300 mM sodium acetate pH 6.4, 1 mM EDTA) overnight at 42°C. RNase protection assays were performed using the RPA II Ribonuclease Protection Kit (Ambion, Austin, TX). The RNA samples were digested with the RNase A/T1 mixture diluted 1:75. The protected RNA probe fragments were separated on a 6% acrylamide/7.6 M urea gel and exposed to Kodak X-Omat film.

PCR assays. DNA was isolated from cells in culture or tissues of experimental birds using the QIAmp Tissue Kit (Qiagen). Each PCR contained 1.25 µl 10X PCR buffer (final concentration, 50 mM Tris•Cl, pH 8.3, 50 mM KCl, 7 mM MgCl₂, 1.1 mM β -mercaptoethanol), 1.25 μ l of 1.7 mg/ml BSA, 0.5 15 μ l of each dNTP at 25 mM, 0.5 μ l of each primer (A₂₆₀ = 5), 6.0 μ l H₂O, and 1.0 μl of DNA (genomic DNA about 100 ng/μl; plasmid DNA about 2 ng/μl). The reactions were heated to 90°C for 1 minute and initiated by the addition of 1.5 μl of Taq DNA polymerase (Promega, Madison, WI) diluted 1:10 v/v (0.75 units). Thirty cycles of PCR were carried out as follows: 90°C for 40 seconds, then 20 59°C for 80 seconds. Diagnostic primers used to detect ALV(A) env (Bora et al., 1988) were 5'-GGGACGAGGTTATGCCGCTG-3' (SEQ ID NO:24; about 50 bp upstream of KpnI site) and 5'-GGGCGTGCGCGCATTACCAC-3' (SEQ ID NO:25; nucleotides 871-851), yielding a 937 bp fragment. The PCR extension temperature was increased to 62°C for amplifying ALV(A) env. 25 Diagnostic primers used to detect ALV(B) env (Bora et al., 1986) were 5'-GACCGACCCAGGGAACAATC-3' (SEQ ID NO:26; nucleotides 713-732) and 5-ATGAGGAAAATTGCGGGTGG-3' (SEQ ID NO:27; nucleotides 1141-1122), yielding a 429 bp fragment. Diagnostic primers used to detect stva (Bates et al., 1993; Connolly et al., 1994) were 5'-GGAATGTGACTGGTAATGGA-3' 30 (SEQ ID NO:28; nucleotides 56-75) and 5'-GCCTTAGTGATGGTGATGGT-3' (SEO ID NO:29; nucleotides 369-350), yielding a 314 bp fragment. Diagnostic primers used to detect stva-mIgG were 5'-CCATCCGTCTTCATCTTCCCT-3'

(SEQ ID NO:30; nucleotides 974-994) and 5'-

TGGTGCGGTGTCCTTGTAGTT-3' (SEQ ID NO:31; nucleotides 1562-1542), yielding a 589 bp fragment of the mouse IgG gene (Tucker et al., 1979). The amplified DNA fragments were separated on 0.8% agarose gels and visualized with ethidium bromide.

Results

30

Experimental approach. The stva and stva-mIgG receptor gene fusions were subcloned into the CLA12NCO adaptor plasmid which contains a transcriptional leader sequence and has a consensus ATG start site contained in a Ncol site. These sequences work very efficiently with the promoter/enhancer elements of the ALV-based retroviral vectors to express experimental genes at high levels (Hughes et al., 1987). The RCAS family of retroviral vectors were derived from the Schmidt-Ruppin A strain of Rous sarcoma virus (RSV) and were present in proviral form on pBR-based plasmids (Federspiel et al., 1997). Experimental genes were inserted into the vectors in the unique ClaI site (which 15 replaces the src gene in RSV) and so are translated from a spliced mRNA. Retroviral vectors that carry and express the stva and stva-mIgG genes are shown schematically in Figure 11. Virus propagation was initiated by transfection of plasmid DNA containing the retroviral vector into avian cells (Figure 12). The culture was then passaged until a maximum viral titer was achieved (6-10 cell 20 passages depending on the vector (Federspiel et al., 1997). Because vectors that use different receptors are available, this system can be used to deliver multiple genes to virtually all cells in the culture (Givol et al., 1994). Cell cultures that express sTva or sTva-mIgG from a subgroup (B) or (C) vector were subsequently challenged with ALV(A) to quantitate the antiviral effect of the 25 sTva proteins.

Antiviral effect of sTva in vitro and in vivo. The initial experiments testing the effects of soluble receptors on viral replication were done with the sTva protein. The stva gene was introduced into the RCASBP vector, which produces the highest titer viral stocks and the highest level of expression of an experimental. To quantitate the antiviral effect of sTva on ALV(A) infection, CEF fully infected with the vector alone [either RCASBP(B) or RCASBP(C)], vectors that express the stva gene [RCASBP(B)stva or RCASBP(C)stva] or

uninfected CEF were challenged with RCASBP(A)AP, the subgroup (A)

RCASBP vector containing the human placental alkaline phosphatase reporter gene (AP). The results of a representative assay are shown in Table 8. CEF cultures producing sTva, either from RCASBP(B) or RCASBP(C), were > 100-fold more resistant to infection by RCASBP(A)AP than cultures infected with the vector alone. CEF cultures infected by the RCASBP(B) or RCASBP(C) vector alone were 3-5-fold less susceptible to RCASBP(A)AP. The antiviral effect of sTva was specific for RCASBP(A)AP infection, since infection by viruses with other envelope subgroups were not inhibited (Table 8).

10

Table 8. Relative resistance of CEF expressing sTVA or sTVA-mIgG to ALV infection.

		Challenge Virus		
		RCASBP(A)AP ^a	RCASBP(B)AP	RCASBP(C)AP
15	Uninfected CEF	1.5 x 10 ⁶	2.1 x 10 ⁵	8.4 x 10 ⁴
	RCASBP(B)	$4.3 \times 10^5 (4)^b$	ND^c	$2.0 \times 10^4 (4.2)$
	RCASBP(B)stva	$2.9 \times 10^3 (517)$	ND	2.1 x 10 ⁴ (4.0)
	RCASBP(C)	$6.0 \times 10^5 (3)$	8.3 x 10 ⁴ (2.5)	ND
	RCASBP(C)stva	$4.0 \times 10^3 (375)$	$1.2 \times 10^5 (1.8)$	ND
20	RCASBP(C)stva-mlgG	1.5 x 10 ³ (1000)	8.0 x 10 ⁴ (2.6)	ND

^aAP-human placental alkaline phosphatase reporter gene.

25

30

The CEF cultures infected with RCASBP(B) and RCASBP(B)stva were used to inoculate unincubated line 0 eggs to produce chicks viremic with RCASBP(B) or RCASBP(B)stva. Viremic chicks produced in this manner are tolerant to most ALV antigens since the early embryo was infected. The chicks were challenged with 10⁵ infectious units of RAV-1, an aggressive ALV(A) strain, to quantitate the antiviral effect of sTva. Blood samples were collected from representative birds of each group at 2 weeks post-challenge, and from all birds 9 weeks post-challenge. The sera were assayed for ALV(A) and ALV(B) by the *in vitro* ALV Assay. The results of the challenges are summarized in

^bThe resistance of experimental and control CEF to ALV infection relative to uninfected CEF is given in parentheses.

^{&#}x27;ND-not done

Table 9. Ninety-six percent of the birds infects with the RCASBP(B) vector alone and then challenged with RAV-1 produced ALV(A) at both experimental time points as expected. However, 95% of the birds infected with RCASBP(B)stva did not produce detectable levels of ALV(A). These results demonstrate that sTva has a strong antiviral effect on ALV(A) infection both *in vitro* and *in vivo*. These results also demonstrate the utility of using vectors with different subgroups *in vivo* since experimental birds could be infected with both the RCASBP(B) vector and RAV-1. However, the level of sTva expression could not be quantitated since neither the hemagglutinin nor the histidine epitope tags included on the sTva protein allowed efficient immunoprecipitation of the protein.

Table 9. Chickens expressing sTva are resistant to ALV(A) infection.

		Virus Subgroup Detected					
		2 Weeks ^a			9 Weeks		
15		A&B	В	Α	A&B	В	A
	Uninfected	0/10	0/10	0/10	0/9	0/9	0/9
	RCASBP(B)	9/9	9/9	0/9	6/6	6/6	0/6
	RCASBP(B)+RAV-1	10/10	10/10	9/10	15/15	15/15	15/15
	RCASBP(B)stva	10/10	10/10	1/10	11/12	11/12	0/12
20	RCASBP(B)stva+RAV-1	10/10	10/10	0/10	20/20	20/20	1/20

^aAll of the birds in each group were not necessarily assayed at both time points.

Antiviral effect of sTva-mIgG in vitro. Although the tagged version of sTva could not be immunoprecipitated efficiently, an sTva immunoadhesin sTva-mIgG can be immunoprecipitated and quantitated. sTva-mIgG consists of the 83-amino acid Tva extracellular domain fused to the constant region of the mouse IgG heavy chain. The stva-mIgG gene was introduced into the RCASBP(C) vector. To quantitate the antiviral effect of sTva-mIgG compared to sTva, CEF cultures infected with RCASBP(C)stva-mIgG, RCASBP(C)stva, or RCASBP(C) were challenged with either RCASBP(A)AP or RCASBP(B)AP. CEF expressing sTva-mIgG were about 300-fold more resistant to

5

10

15

20

25

30

RCASBP(A)AP infection compared to cells infected with the vector alone, and 2- to 3-fold more resistant than cells expressing sTva (Table 8). The antiviral effect was specific for ALV(A) since no significant change in susceptibility was observed when the cultures were challenged with RCASBP(B)AP.

ALV replication in a permanent, non-transformed cell line derived from line 0 CEF called DF-1 has been described (Himly et al., 1998; Schaefer-Klein et al., 1998). ALV and ALV-based retroviral vectors replicate and express inserted genes in DF-1 cells at levels similar to CEF, and DF-1 can be used to generate clonal cell lines. The antiviral effect of sTva-mIgG produced in DF-1 cultures infected with RCASBP(C)stva-mIgG (Table 10) was similar to that seen in CEF cultures (Table 8). The sTva-mIgG protein was immunoprecipitated from cell culture supernatants with anti-mouse IgG antibody conjugated to agarose beads and analyzed by Western transfer of SDS-PAGE gels (Figure 13). The immunoprecipitated sTva-mIgG protein migrates as a broad band (50-60 kDa) due to post-translational modification and as a minor about 38 kDa band. The about 38 kDa band is probably a degradation product of sTva-mIgG since both bands appear after immunoprecipitation with an ALV(A) surface glycoprotein immunoadhesin, and the amount of the about 38 kDa band increases after repeated freeze-thaw cycles of the viral supernatants. Stable clonal DF-1 cell lines were generated that express different levels of sTva-mIgG under the control of the TFANEO expression vector. These cell lines do not produce infectious ALV and are resistant to RCASBP(A)AP infection at levels similar to cultures expressing sTva-mIgG from the retroviral vectors. Therefore, chronic ALV infection does not make a major contribution to the antiviral effect obtained.

Relationship between sTva-mIgG expression level and the antiviral effect. The stva-mIgG gene was subcloned into the RCAS(C) and RCOSBP(C) retroviral vectors. Previously, it has been reported that the RCAS vector replicates to a 5-15-fold lower titer (depending on envelope subgroup and inserted gene) compared to RCASBP and produces protein at an equivalently reduced level (Federspiel et al., 1994). The RCOSBP vector, which lacks a strong transcription enhancer in the LTR, replicates to about 100-fold lower titer compared to RCASBP and produces lower levels of protein. DF-1 cultures infected with RCASBP(C)stva-mIgG, RCAS(C)stva-mIgG, or RCOSBP(C)stva-mIgG, or

mIgG were challenged with RCSABP(A)AP to determine the antiviral effect of different levels of sTva-mIgG on ALV(A) infection. The results of a representative assay are shown in Table 10. The sTva-mIgG produced by these DF-1 cultures was detected by immunoprecipitation of sTva-mIgG followed by the analysis of the proteins by Western transfer (Figure 13). As expected, cultures infected with RCASBP produced the highest level of sTva-mIgG and the greatest antiviral effect (about 200-fold). Cultures infected with RCAS produced slightly lower levels of sTva-mIgG protein and a lower antiviral effect (about 100-fold). Finally, cultures infected with RCOSBP produced the lowest level of sTva-mIgG protein and a modest antiviral effect (about 15-fold).

Table 10.

Relative resistance of DF-1 cells expressing sTva-mIgG to ALV(A) infection.

		RCASBP(A)AP	Resistance
	Uninfected DF-1	3.3×10^6	
15	RCASBP(C)	1.3×10^6	3
	RCASBP(C)stva-mIgG	5.3×10^3	622
	RCAS(C)	6.4×10^5	5
	RCAS(C)stva-mIgG	5.7×10^3	579
	RCOSBP(C)	9.0×10^{5}	4
20	RCOSBP(C)stva-mIgG	5.2 x 10 ⁴	63

^aThe resistance of the cells to ALV(A) infection was determined by dividing the titer obtained on the control uninfected DF-1 cells by the titer obtained for each experimental group.

25

30

The antiviral effect of sTva and sTva-mIgG on ALV(A) infection may represent the minimum antiviral effect attainable *in vitro* as measured by the direct ALV AP challenge assay. The assays were done on subconfluent cell cultures (30%) where the levels of the soluble receptor protein had not accumulated to the levels expressed by a confluent culture. To determine the antiviral effect of higher levels of sTva-mIgG, RCASBP(A)AP was pretreated with supernatants collected from confluent DF-1 cultures infected with RCASBP(B), RCASBP(B)stva-mIgG, RCOSBP(B), or RCOSBP(B)stva-mIgG, and then assayed as before. Preabsorption of RCASBP(A)AP with high levels of

5

15

20

sTva-mIgG significantly increased the antiviral effect compared to a direct assay: RCASBP(B)stva-mIgG pretreatment increased the antiviral effect of the direct assay about 500-fold; and RCOSBP(B)stva-mIgG pretreatment increased the direct antiviral effect about 60-fold.

Delivery and expression of sTva and sTva-mlgG in vivo. To characterize the efficiency of RCASBP delivery and expression of the sTva proteins in chickens, unincubated line 0 eggs were injected with CEF producing the RCASBP(B) or RCASBP(C) vectors alone, the vectors with the stva gene, or the vectors with the stva-mIgG gene. Viremic chicks were identified on the day of hatch by an ELISA assay for ALV capsid protein. The sTva-mIgG protein was immunoprecipitated from serum samples of both RCASBP(B)stva-mIgG and RCASBP(C)stva-mIgG infected birds and visualized by Western analysis (Figure 14). The stva, stva-mIgG, and RCASBP(B) env RNA expression levels in liver, heart, spleen, bursa, thymus, kidney, and muscle tissues of infected birds were analyzed by RNase protection assay. An RNase protection analysis of a representative bird infected with RCASBP(B)stva and a representative bird infected with RCASBP(B)stva-mIgG are shown in Figure 15. Relatively high levels of the stva or stva-mlgG and ALV(B) env RNAs were detected in all tissues assayed, indicating that the inserted genes were delivered and expressed efficiently by the RCASBP(B) vector.

Table 11. Chickens expressing either sTva or sTva-mIgG are resistant to ALV(A) infection but not ALV(C) infection.

		Virus Subgroup Detected ^a		
5		В	Α	C
	RCASBP(B)+RAV-1	4/4	4/4	
	RCASBP(B)stva+RAV-1	5/5	0/5	
	RCASBP(B)stva-mIgG+RAV-1	9/9	0/9	
10	RCASBP(B)+RAV-49	4/4		3/4
	RCASBP(B)stva+RAV-49	4/4		4/4
	RCASBP(B)stva-mIgG+RAV-49	7/8		6/8

^aAssays were done 4 weeks after challenge.

15

20

25

30

Antiviral effect of sTva-mIgG in vivo. Chicks infected with the RCASBP(B) vector alone, RCASBP(B)stva, or RCASBP(B)stva-mIgG were split into two groups and challenged with 10⁵ infectious units of either RAV-1 (subgroup A) or RAV-49 (subgroup C). Blood was collected from each bird four weeks after challenge, and the serum was assayed for ALV(A), ALV(B) and ALV(C) by the in vitro ALV Assay (Table 11). As expected, ALV(B) was detected in virtually all of the birds since the RCASBP(B) vector was used for gene delivery. ALV(A) was not detected in the serum of RAV-1 challenged birds containing the stva or the stva-mlgG genes. However, ALV(A) was detected in the serum of the birds infected with the RCASBP(B) vector alone and challenged with RAV-1. In contrast, the birds of all three experimental groups were equally susceptible to RAV-49 challenge as shown by the presence of ALV(C) in the majority of the birds. Since 19% of the birds challenged with RAV-49 did not produce detectable levels of ALV(C), the titer of the RAV-49 stock may have been lower than expected. The antiviral effect of sTva and sTvamIgG was specific for ALV(A), consistent with the proposed mechanism of antiviral action, receptor interference.

Representative birds from each RAV-1 challenged experimental group were analyzed for the presence of ALV(A) and ALV(B) env, stva and stva-mIgG

sequences in RNA and DNA isolated from a variety of tissue samples of each bird. A low level of RAV-1 infection and replication in a subset of tissues may go undetected by the in vitro ALV assay due to virus inactivation by the sTva or sTva-mIgG proteins in the serum. Antisense riboprobes and primer sets were developed to specifically detect each target sequence by RNase protection assay and PCR. RNA and DNA were isolated from liver, heart, spleen, bursa, thymus, kidney, and muscle tissue of each bird and analyzed by RNase protection assay and PCR assay. A representative RNase protection assay of RNA of one tissue (bursa) from a bird of each experimental group, and an uninfected control bird, is 10 shown in Figure 16. A representative PCR analysis of DNA isolated form tissues of a RAV-1 challenged bird from each experimental group is shown in Figure 17. RAV-1 RNA and DNA were only detected in tissues of birds infected with the RCASBP(B) vector challenged with RAV-1. Therefore, the expression of sTva and sTva-mIgG significantly reduces, if not eliminates, infection by the ALV(A) strain RAV-1 in chickens. 15

Cells expressing the sTva proteins showed significant resistance to

Discussion

ALV(A) infection, presumably due to the secreted receptor proteins binding the glycoproteins of the invading virion, blocking the interactions of the virus and 20 the membrane bound Tva, a form of receptor interference. Tva has been hypothesized to be necessary and sufficient to mediate ALV(A) entry (Balliet et al., 1998; Bates et al., 1993). Several possible mechanisms could account for the sTva inhibition of ALV(A) entry sTva binding of an ALV(A) surface glycoprotein may lead to an irreversible conformational change in SU and TM. 25 Several studies have shown that sTva binding to purified ALV(A) envelope glycoproteins induces a temperature-dependent conformational change in the glycoproteins, and appears to convert the envelope glycoproteins to a membranebinding state (Balliet et al., 1999; Damico et al., 1999; Gilbert et al., 1995; Hernandez et al., 1994). Binding of sTva or sTva-mIgG to the envelope glycoproteins on the surface of the virus induces a conformational change in 30 both SU and TM similar to the events leading to fusion of the viral and host cell membranes, and converts SU and TM into a form that is unable to bind Tva on the surface of the cell. A conformational change may also lead to the loss of

some of the SU subunits (Orloff et al., 1993). Finally, sTva may inhibit ALV(A) entry by simply binding to SU and physically blocking the access of membrane bound Tva to the virion. By whatever mechanism(s), the sTva proteins block the entry of ALV(A) into cultured cells and cells and tissues of chickens.

5 The replication-competent ALV-based retroviral vector experimental system enabled the efficient delivery and expression of the stva and stva-mIgG genes both in cultured cells and in virtually all the cells and tissues of the chicken. RCASBP(A) and (RCASBP(B), as well as other combinations of ALV retroviral vectors [ALV(B) followed by ALV(A); ALV(C) followed by 10 AVL(A)], can be used in CEF and DF-1 cells in vitro and in vivo. As reported previously, the replication of some RCASBP(B) viruses in CEF and DF-1 cells, and RCASBP(C) viruses on DF-1 cells, were somewhat cytopathic (Himly et al., 1998; Schaefer-Klein et al., 1998). The cytopathic effect manifests itself as a pause in growth rate (2-6 days), after which the cells recover, divide at a normal rate, and express the viral and experimental proteins. It appears that only the 15 cells that produce high levels of the RCASBP(B) or RCASBP(C) envelope glycoproteins show the cytopathic effects. The replication of subgroup (B) and(C) RCAS and RCOSBP viruses, which replicate to lower titers compared to RCASBP, do not cause detectable cytopathic effects. Chronic infection of CEF and DF-1 cells with an ALV vector results in a low level of resistance to 20 infection by other ALV env subgroup vectors (2-5-fold) compared to uninfected cells. This may indicate that while the ALV glycoproteins specifically and efficiently interact with the appropriate receptor resulting in receptor interference, the high level expression of one type of envelope glycoprotein on the cell surface may interfere either directly or indirectly with the ability of other 25 ALVs to interact with their host receptors.

Both the RCASBP(B) and RCASBP(C) retroviral vectors were efficient in generating viremic chicks without detectable pathologic effects in short term infections, and the infected chicks expressed relatively high levels of sTva-mIgG protein in their serum. Chicks infected with RCASBP(B) were also efficiently infected with ALV(A). The RCASBP(B) vector efficiently delivered and expressed the *stv-a* and *stva-mIgG* genes in all tissues tested, and resulted in a significant antiviral effect on ALV(A) infection and replication. By delivering

30

the stva genes in the early embryo, the immune system of the chicken can be evaded. The birds are tolerized to the sTva and sTva-mIgG proteins, and to most of the ALV antigens since the viral vector used to deliver stva and stva-mIgG and the challenge viruses are virtually identical except for small regions of SU.

The expression of the sTva and sTva-mIgG proteins *in vivo* allowed the examination of the effects of the viral glycoprotein-soluble receptor interactions in a wide variety of cells. One RAV-1 challenged RCASBP(B)stva infected bird did contain infectious ALV(A) in its serum. Unfortunately the bird died of non-viral causes before tissues could be obtained.

CD4, an important cell-surface protein of the T-lymphocytes, is the 10 primary receptor for HIV-1. Several groups developed and expressed soluble forms of CD4 (sCD4) and demonstrated that recombinant sCD4 proteins could bind specifically to HIV-1 envelope glycoproteins and inhibit HIV-1 infection in vitro (Daar et al., 1990; Harbison et al., 1990; Klasse et al., 1993; Orloff et al., 1993: Schacker et al., 1995; Weiss, 1992). For injection of a recombinant 15 antiviral protein to be effective against cell to cell transmission of the virus, it may be necessary to use a gene therapy approach in which target cells actively express the soluble receptor. The gene therapy approach has been tested for HIV-1: a sCD4 gene construct was expressed by a murine leukemia virus-based retroviral vector in human T-cell lines and in primary peripheral blood 20 lymphocytes (Morgan et al., 1994). In cell culture populations engineered to express sCD4 (30-50% of the cells contained the sCD4 gene) HIV-1 replication was inhibited 50-70% indicating that a sCD4 antiviral approach against HIV-1 might be more effective. Since the initial sCD4 studies were published, the chemokine receptors have been identified as co-receptors necessary for efficient 25 HIV-1 entry into cells (Hunter, 1997). Since both CD4 and a chemokine receptor are required for efficient HIV-1 entry into cells, sCD4 alone may not be an effective inhibitor of HIV-1 entry.

The results clearly indicate that a soluble receptor interference antiviral strategy can effectively block the replication of at least some retroviruses, and that this approach may be applicable to other virus groups that require specific viral glycoprotein-host receptor interactions for entry into the cell. The application of this strategy to protect animals against specific viral diseases is

30

relatively straightforward since transgenic technology can be used to introduce genes into animals and the transgenes will produce the desired protein without provoking an immune response.

Example 6

5

10

15

20

25

30

Selection of Subgroup A Avian Leukosis Virus Mutants Resistant to the Receptor

Interference Imposed by Soluble TVA and SUA

Genes encoding soluble forms of the subgroup A avian leukosis virus (ALV) receptor Tva (sTva) and soluble envelope surface glycoprotein (SUA), delivered by ALV-based retroviral vectors to cells, significantly inhibit ALV(A) infection of those cells. The antiviral effects of sTva and SUA are consistent with a receptor interference (RI) mechanism; the antiviral effect is specific for ALV(A) since the susceptibility to infection by a neutral subgroup remains unchanged. A virus-free stable cell line was developed which expressed either sTva or SUA and so was significantly resistant to ALV(A) infection, 50-350-fold and 1,000 to 200,000-fold, respectively. Since the SU region of Env determines receptor usage, mutations capable of overcoming the RI may occur in the SU region of the *env* gene of ALV(A) and would affect interactions between Tva and Env(A).

ALV(A) was passaged on the sTva and SUA expressing cell lines. Variant viruses capable of more rapid growth on the sTva cell lines were identified at passage 7 and on the SUA cell lines at passage 9 post-infection. The SU region of the *env* gene was PCR amplified from these cells, cloned and analyzed by sequencing. Selection on sTva expressing cells produced mutant viruses which were identical in sequence to ALV(A) except that 50% of the clones screened had an amino acid change at codon 142, while the other 50% had an amino acid change at codon 149 of *env*. All mutants selected on SUA expressing cell lines contained a 6 amino acid deletion at codons 155-160 within *env*. To determine if these mutations were sufficient to confer enhanced growth on the resistant cell lines, the SU region of *env* containing the mutation, from Kpn I to Sal I, was introduced into a wild-type ALV(A) molecular clone.

Transfection of these mutants showed an enhanced rate of growth as compared to wild-type.

5

15

30

References

Achong, G. G., P. A. Trumper, and B. C. Giovanella. C-type virus particles in human tumors transplanted into nude mice. British Journal of Cancer, 34:203-206 (1976).

Adkins, H. B., J. Brojatsch, J. Naughton, M. M. Rolls, J. M. Pesola, and J.A.T. Young. Identification of a cellular receptor for subgroup E avian leukosis virus. PNAS. 94:11617-11622 (1997).

Akiyoshi, E. E., M. Denaro, H. Zhu, J. L. Greenstein, P. Banerjee, and J. A. Fishman. Identification of a full-length cDNA for an endogenous retrovirus of miniature swine. <u>Journal of Virology</u>, 72:4503-4507 (1998).

Armstrong, J. A., J. S. Porterfield, and A. T. de Madrid. C-type virus particles in pig kidney cell lines. <u>Journal of General Virology</u>, 10:195-198 (1971).

Astrin, S. M., E. G. Buss, and W. S. Hayward. Endogenous viral genes are non-essential in the chicken. <u>Nature</u>. 282:339-341 (1979).

Balliet, J. W., and P. Bates. Efficient infection mediated by viral receptors incorporated into retroviral particles. <u>Journal of Virology</u>. 72:671-676 (1998).

Balliet, J. W., J. Berson, C. M., D'Cruz, J. Huang, J. Crane, J. M. Gilbert, and P. Bates. Production and characterization of a soluble, active form of Tva, the subgroup A avian sarcoma and leukosis virus receptor. Journal of Virology. 73:3054-3061 (1999).

Bassin, R. H., N. Tuttle, and J. R. Fischinger. Rapid cell culture assay technique for murine leukemia viruses. <u>Nature</u> (London), <u>229</u>:564-566 (1971).

Bates, P., L. Rong, H. E. Varmus, J. A. T. Young, and L. B. Crittenden. Genetic mapping of the cloned subgroup A avian sarcoma and leukosis virus receptor gene to the TVA locus. <u>Journal of Virology</u>. 72:2505-2508 (1998).

Bates, P., J. A. T. Young, and H. E. Varmus. A receptor for subgroup A. Rous sarcoma virus is related to the low density lipoprotein receptor. <u>Cell</u>. 74:1043-1051 (1993).

Belanger, C., K. Zingler, and J. A. Young. Importance of cysteines in the LDLR-related domain of the subgroup A avian leukosis and sarcoma virus receptor for viral entry. <u>Journal of Virology</u>. <u>69</u>:1019-1024 (1995).

Bergelson, J. J., J. G. Mohanty, R. L. Crowell, N. F. St. John, D. M. Lublin, and R. W. Finberg. Coxsackievirus B3 adapted to growth in RD cells binds to decay-accelerating factor (CD55). <u>Journal of Virology</u>, 69:1903-1906 (1995).

10

25

30

Boeke, J. D., and J. P. Stoye. Retrotransposons, endogenous retroviruses, and the evolution of retroelements. *In J. M. Coffin and S. H. Hughes and H. E. Varms* (ed.), Retroviruses, p. 343-435, Cold Spring Harbor Laboratory Press.

Bostock, D. E., and L. N. Owen. Porcine and ovine lymphosarcoma: A review. <u>Journal of the National Cancer Institute</u>, <u>50</u>:9330939 (1973).

Bova, C. A., J. P. Manfredi, and R. Swanstrom. Env genes of avian retroviruses: nucleotide sequence and molecular recombinants define host range determinants. <u>Virology</u>. <u>152</u>:343-354 (1986).

Bova, C. A., J. C. Olsen, and R. Swanstrom. The avian retrovirus env gene family: molecular analysis of host range and antigenic variants. <u>Journal of Virology</u>. 62:75-83 (1988).

Breimer, M. E., S. Bjorck, S. C. T., A. Bengtsson, L. Rydberg, K. Lie-Karlsen, P.-O. Attman, M. Aurell, and B. E. Samuelsson. Extracorporeal ("ex vivo") connection of pig kidneys to humans. <u>I. Clinical data and studies of platelet destruction</u>. Xenotransplantation, 3:328-339 (1996).

Brojatsch, J., J. Naughton, M. M. Rolls, K. Zingler, and J. A. T. Young. CAR1, a TNFR-related protein, is a cellular receptor for cytopathic avian leukosis-sarcoma viruses and mediates apoptosis. <u>Cell.</u> <u>87</u>:845-855 (1996).

Chong, H., W. Starkey, and R. G. Vile. A replication-competent retrovirus arising from a split-function packaging cell line was generated by recombination events between the vector, one of the packaging constructs, and endogenous retroviral sequences. <u>Journal of Virology</u>, 72:2663-2670 (1998).

Connolly, L., K. Zingler, and J. A. T. Young. A soluble form of a receptor for subgroup A avian leukosis and sarcoma viruses (ALSV-A) blocks infection and binds directly to ALSV-A. Journal of Virology. 68:2760-2764 (1994).

- Crawford, D. H., B. G. Achong, N. M. Reich, S. Finerty, J. L. Thompson, M. A. Epstein, and B. C. Giovanella. Identification of murine endogenous xenotropic retrovirus in cultured multicellular tumor spheroids from nudemouse-passaged nasopharyngeal carcinoma. International Journal of Cancer, 23:1-7 (1979).
- 10 Crittenden, L. B., D. W. Salter, and M. J. Federspiel. Segregation, viral phenotype, and proviral structure of 23 avian leukosis viruses inserted in the germ line of chickens. Theoretical Applied Genetics, 77:505-515 (1989).
- Daar, E. S., X. L. Li, T. Moudgil, and D. D. Ho. High concentrations of recombinant soluble CD4 are required to neutralize primary human immunodeficiency virus type 1 isolates. Proceedings of the National Academy of Sciences, USA. 87:6574-6578 (1990).
 - Damico, R., L. Rong, and P. Bates. Substitutions in the receptor-binding domain of the avian sarcoma and leukosis virus envelope uncouple receptor-triggered structural rearrangements in the surface and transmembrane subunits. Journal of Virology. 73:3087-3094 (1999).
 - Deacon, T., J. Schumacher, J. Dinsmore, C. Thomas, P. Palmer, S. Kott, A. Edge, D. Penney, S. Kassissieh, P. Dempsey, and O. Isacson. Histological evidence of fetal pig neural cell survival after transplantation into a patient with Parkinson's disease. Nature Medicine, 3:350-353 (1997).
- Diamond, L. E. K. R. McCurry, M. J. Martin, S. B. McClellan, E. R. Oldham, J. L. Platt, and J. S. Logan. Characterization of transgenic pigs expressing functionally active human CD59 on cardiac endothelium.

 Transplantation, 61:1241-1249 (1996).

20

Donahue, R. E., S. W. Kessler, D. Bodine, K. McDonagh, C. Dunbar, S. Goodman, B. Agricola, E. Byrne, M. Raffeld, R. Moen, J. Bacher, K. M. Zsebo, and A. W. Nienhuis. Helper virus induced T cell lymphoma in nonhuman primates after retroviral mediated gene transfer. The Journal of Experimental Medicine, 176:1125-1135 (1992).

Dorig, R. E., A. Marcil, A. Chopra, and C. D. Richardson. The human

CD46 molecule is a receptor for measles virus (Edmonston Strain). Cell, 75:295-305 (1993).

Ebert et al., Biotech, 9, 835 (1991).

10

30

Federspiel, M. J., L. B. Crittenden, and S. H. Hughes. Expression of avian reticuloendotheliosis virus envelope confers host resistance. <u>Virology</u>, 173:167-177 (1989).

Federspiel, M. J., L. B. Crittenden, L. P. Provencher, and S. H. Hughes. Experimentally introduced defective endogenous proviruses are highly expressed in chickens. Journal of Virology, 65:313-319 (1991).

Federspiel, M.J., and S.H. Hughes. Effects of the *gag* region on genome stability: avian retroviral vectors that contain sequences from the Bryan strain of Rous sarcoma virus. <u>Virology</u>, 203:211-220 (1994).

Federspiel, M.J., D.A. Swing, B. Eagleson, S.W. Reid, and S.H. Hughes.

Expression of transduced genes in mice generated by infecting blastocysts with avian leukosis virus-based retroviral vectors. Proc. Natl. Acad. Sci. USA, 93:4931-4936 (1996).

Federspiel, M. J., P. Bates, J. A. T. Young, and H. E. H. Varmus, S.H. A system for tissue-specific gene targeting: Transgenic mice susceptible to subgroup A avian leukosis virus-based retroviral vectors. Proc. Natl. Acad. Sci. USA, 91:11241-11245 (1994).

Federspiel, M. J., and S. H. Hughes. Retroviral gene delivery. Methods in <u>Cell Biology</u>. 52:179-214 (1997).

Fekete, D. M., and C. L. Cepko. Retroviral infection coupled with tissue transplantation limits gene transfer in the chicken embryo. Proc. Natl. Acad. Sci. USA. 90:2350-2354 (1993).

Fields-Berry, S. C., A. L. Halliday, and C. L. Cepko. A recombinant retrovirus encoding alkaline phosphatase confirms clonal boundary assignment in lineage analysis of murine retina. <u>Proceedings of the National Academy of Sciences</u>. USA. 89:693-697 (1992).

Fishman, J. A., Miniature swine as organ donors for man: strategies for prevention of xenotransplant-associated infections. Xenotransplantation, 1:47-57 (1994).

Fodor, W. L., B. L. Williams, L. A. Matis, J. A. Madri, S. A. Rollins, J.

W. Knights, W. Velander, and S. P. Squinto. Expression of a functional human complement inhibitor in a transgenic pig as a model for the prevention of xenogeneic hyperacute organ rejection. Proceedings of the National Academy of Sciences, 91:11153-11157 (1994).

Frazier, M. E. Evidence for retrovirus in miniature swine with radiation-induced leukemia or metaplasia. Archives of Virology, 83:83-97 (1985).

Gautsch, J. W., A. F. Knowles, F. C. Jensen, and N. O. Kaplan. Highly efficient induction of type C retroviruses by a human tumor in athymic mice.

10 Proceedings of the National Academy of Sciences, 77:2247-2250 (1980).

5

25

30

Gilbert, J. M., L. D. Hernandez, J. W. Balliet, P. Bates, and J. M. White. Receptor-induced conformational changes in the subgroup A avian leukosis and sarcoma virus envelope glycoprotein. <u>Journal of Virology</u>. 69:7410-7415 (1995).

Givol, I., I. Tsarfaty, J. Resau, S. Rulong, P. P. da Silva, G. Nasioulas, J. DuHadaway, S. H. Hughes, and D. L. Ewert. Bcl-2 expressed using a retroviral vector is localized primarily in the nuclear membrane and the endoplasmic reticulum of chicken embryo fibroblasts. Cell Growth and Differentiation. 5:419-429 (1994).

20 Groth, C. G., O. Korsgren, A. Tibell, J. Tollemar, E. Moller, J. Bolinder, J. Ostman, F. P. Reinholt, C. Hellerstrom, and A. Andersson. Transplantation of porcine fetal pancreas to diabetic patients. <u>Lancet</u>, 344:1402-1404 (1994).

Harbison, M. A., J. M. Gillis, P. Pinkston, R. A. Byrn, R. M. Rose, and S. M. Hammer. Effects of recombinant soluble CD4 (rCD4) on HIV-1 infection of monocyte/macrophages. <u>Journal of Infectious Diseases</u>. <u>161</u>:1-6 (1990).

Heneine, W., A. Tibell, W. M. Switzer, P. Sandstrom, G. Vazques Rosales, A. Mathews, O. Korsgren, L. E. Chapman, T. M. Folks, and C. G. Groth. No evidence of infection with porcine endogenous retrovirus in recipients of porcine isle-cell xenografts. Lancet, 352:695-699 (1998).

Henninghausen, Protein Expression and Purification, 1:3 (1990).

Hernandez, L. D., R. J. Peters, S. E. Delos, J. A. T. Young, D. A. Agard, and J. M. White. Activation of a retroviral membrane fusion protein: soluble

receptor-induced liposome binding of the ALSV envelope glycoprotein. Journal of Cell Biology. 139:1455-1464 (1997).

Himly, M., D. N. Foster, I. Bottoli, J. S. Iacovoni, and P. K. Vogt. The DF-1 chicken fibroblast cell line: transformation induced by diverse oncogenes and cell death resulting from infection by avian leukosis viruses. <u>Virology</u>. 248:295-304 (1998).

Hirsch, M. S., P. H. Black, G. S. Tracy, S. Leibowitz, and R. S. Schwartz. Leukemia virus activation in chronic allogeneic disease. <u>Proceedings of the National Academy of Sciences</u>, <u>67</u>:1914-1917 (1970).

Hirsch, M. S., S. M. Phillips, C. Solnik, P.H. Black, R.S. Schwartz, and C.B. Carpenter. Activation of leukemia viruses by graft-versus-host and mixed lymphocyte reactions *in vitro*. Proceedings of the National Academy of Sciences, 69:1069-1072 (1972).

Hughes, S. H., J. J. Greenhouse, C. J. Petropoulos, and P. Sutrave.

Adaptor plasmids simplify the insertion of foreign DNA into helper-independent retroviral vectors. <u>Journal of Virology</u>. 61:3004-3012 (1987).

Hunter, E. Viral entry and receptors, p. 71-119. *In* J. M. Coffin and S. H. Hughes and H. E. Varmus (ed.), <u>Retroviruses</u>. Cold Spring Harbor Laboratory Press (1997).

20 Jaenish, <u>Science</u>, <u>240</u>:468 (1988).

30

Kanaya, S., and R.J. Crouch. Low levels of RNase H activity in *Escherichia coli* FB2 *mh* result from a single-base change in the structural gene of RNase H. J. Bacteriol., 154:1021-1026 (1983).

Kawakami, T.G., G.V. Kollias, C. Holmberg. Oncogenicity of Gibbon type-C myelogenous leukemia virus. International Journal of Cancer, 25:641-646 (1980).

Kingston, R. E., C. A. Chen, and H. Okayama. Introduction of DNA into eukaryotic cells, p. 911-919. *In* F. M. Ausubel and R. Brent and R. E. Kingston and D. D. Moore and J. G. Seidman and J. A. Smith and K. Struhl (ed.), Current Protocols in Molecular Biology, vol. 1. Wiley, New York (1989).

Klasse, P. J., and J. A. McKeating. Soluble CD4 and CD4 immunoglobulin-selected HIV-1 variants: a phenotypic characterization. AIDS Research and Human Retroviruses. 9:595-604 (1993).

Le Tissier, P., J.P. Stoye, Y. Takeuchi, C. Patience, and R.A. Weiss. Two sets of human-tropic pig retrovirus. Nature, 389:681-682 (1997).

Leverett, B.D., K.B. Farrell, M.V. Eiden, and C.A. Wilson. Entry of amphotropic murine leukemia virus is influenced by residues in the putative second extracellular domain of the human form of its receptor, Pit2, <u>L. Virol.</u>, 72, (6) 4956-4961 (1998).

Levy, J.A., S.K. Datta, and R.S. Schwartz. Recovery of xenotropic virus but not ecotropic virus during graft-versus-host reaction in mice. Clinical Immunology and Immunopatholog, 7:262-268 (1977).

Lieber, M.M., C.J. Sherr, G.J. Todaro, R.E. Benveniste, R. Callahan, and H.G. Coon. Isolation from the Asian mouse Mus caroli of an endogenous type C virus related to infectious primate type C viruses. <u>Proceedings of the National Academy of Sciences</u>, 72:2315-2319 (1975).

Lin, S.S., and J.L. Platt. Immunologic barriers to xenotransplantation. The Journal of Heart and Lung Transplantation, 15:547-555 (1996).

Makowka, L., D.V. Cramer, A. Hoffman, M. Breda, L. Sher, G. Eiras-Hreha, P.J. Tuso, C. Yasunaga, C.A. Cosenza, G. Du Wu, F.A. Chapman, and L. Podesta. The use of a pig liver xenograft for temporary support of a patient with fulminant hepatic failure. <u>Transplantation</u>, 59:1654-1659 (1995).

20 Marshall et al., Cell & Mol. Neuro., 14, 523 (1994).

10

15

Martin, U., V. Kiessig, J.H. Blusch, A. Haverich, K. von der Helm, T. Herden, and G. Steinhoff. Expression of pig endogenous retrovirus by primary porcine endothelial cell and infection of human cells. Lancet, 352:692-694 (1998).

McCurry, K.R., D.L. Kooyman, C.G. Alvarado, A.H. Cotterell, M.J. Martin, J.S. Logan, and J.L. Platt. Human complement regulatory proteins protect swine-to-primate cardiac xenografts from humoral injury. Nature Medicine. 1:423-427 (1995).

McLachlin, J.R., N. Mittereder, M.B. Daucer, M. Kadan, and M.A.

Eglitis. Factors affecting retroviral vector function and structural integrity.

Virology, 195:1-5 (1993).

Moennig, V., H. Frank, G. Hunsmann, P. Ohms, H. Schwarz, W. Schafer, and H. Strandstrom. C-type particles produced by a permanent cell line from a

leukemic pig II. Physical, chemical, and serological characterization of the particles. Virology, 57:179-188 (1974).

Morgan, R. A., G. Baler-Bitterlich, J. A. Ragheb, F. Wong-Staal, R. C. Gallo, and W. F. Anderson. Further evaluation of soluble CD4 as an anti-HIV type 1 gene therapy: demonstration of protection of primary human peripheral blood lymphocytes form infection by HIV type 1. AIDS Research and Human Retroviruses. 10:1507-1515 (1994).

Natsoulis, G., and J.D. Boecke. New antiviral strategy using capsid-nuclease fusion proteins. <u>Nature</u> (London), <u>352</u>:632-635 (1991).

10

20

25

Natsoulis, G., P. Seshaiah, M.J. Federspiel, A. Rein, S.H. Hughes, and J.D. Boeke. Targeting a nuclease to murine leukemia virus capsids inhibits viral multiplication. Proceedings of the National Academy of Sciences, USA, 92:364-368 (1995).

Nihrane, A., K. Fujita, R. Willey, M. S. Lyu, and J. Silver. Murine leukemia virus envelope protein in transgenic-mouse serum blocks infection *in vitro*. Journal of Virology, 70:1882-9 (1996).

Oldham, E.R., M.A. Velardo, J.L. Platt, J.S. Logan, and L. Diamond. High-level tissue specific expression of human CD59, MCP, and DAF proteins from genomic clones in transgenic mice. Transplantation Proceedings, 28:693 (1996).

Orloff, S. L., M. S. Kennedy, A. A. Belperron, P. J. Maddon, and J. S. McDougal. Two mechanisms of soluble CD4 (sCD4)-mediated inhibition of human immunodeficiency virus type 1 (HIV-1) infectivity and their relation to primary HIV-1 isolated with reduced sensitivity to sCD4. Journal of Virology. 67:1461-1471 (1993).

Ott, D., R. Fredrich, and A. Rein. Sequence analysis of amphotropic and 10A1 murine leukemia viruses: close relationship to mink cell focus-inducing viruses. J. Virol., 64:757-766 (1990).

Ott, D.E., J. Keller, K. Sill, and A. Rein. Phenotypes of murine leukemia virus-induced tumors: influence of 3' viral coding sequences. <u>J. Virol.</u>, <u>66</u>:6107-6116 (1992).

Panabieres, F., M. Piechaczyk, B. Rainer, C. Dani, P. Fort, S. Riaad, L. Marty, J. L. Imbach P. Jeanteur, and J. Blanchard. Complete nucleotide

sequence of the messenger RNA coding for chicken muscle glyceraldehyde-3-phosphate dehydrogenase. Biochem. Biophys. Res. Comm. 118:767-773 (1984).

Patience, C., G.S. Patton, Y. Takeuchi, R.A. Weiss, M.O. McClure, L. Rydberg, and M.E. Breimer. No evidence of pig DNA or retroviral infection in patients with short-term extracorporeal connection to pig kidneys. Lancet, 352:699-701 (1998).

Patience, C., G.R. Simpson, A.A. Colletta, H.M. Welch, R.A. Weiss, and M.T. Boyd. Human endogenous retrovirus expression and reverse transcriptase activity in the T47D mammary carcinoma cell line. <u>Journal of Virology</u>, 70:2654-2657 (1996).

Patience, C., Y. Takeuchi, and R.A. Weiss. Infection of human cells by an endogenous retrovirus of pigs. Nature Medicine, 3:282-286 (1997).

Petropoulos, C. J., and S. H. Hughes. Replication-competent retrovirus vectors for the transfer and expression of gene cassettes in avian cells. <u>Journal of Virology</u>. 65:3728-3737 (1991).

Petropoulos, C. J., W. Payne, D. W. Salter, and S. H. Hughes.

Appropriate *in vivo* expression of a muscle-specific promoter by using avian retroviral vectors for gene transfer. <u>Journal of Virology</u>. <u>66</u>:3391-3397 (1992).

Phan-Thanh, L., B. Kaeffer, and E. Bottreau. Porcine retrovirus: optimal conditions for its biochemical detection. <u>Archives of Virology</u>, 123:255-265 (1992).

20

Policy, C.o.X.T.E.I.a.P. Xenotransplantation science, ethics, and public policy. Institute of Medicine (1996).

Purcell, D.F., C.M. Broscius, E.F. Vanin, C.E. Buckler, A.W. Nienhuis, and M.A. Martin. An array of murine leukemia virus-related elements is transmitted and expressed in a primate recipient of retroviral gene transfer.

Journal of Virology, 70:887-897 (1996).

Ramsdell, F.J., H. Shau, and S.H. Golub. Role of proliferation in LAK cell development, <u>Cancer Immunol. Immunother.</u>, 26, 139-144 (1988).

Robinson, H. L., S. M. Astrin, A. M. Senior, and F. H. Salazar. Host susceptibility to endogenous viruses: defective, glycoprotein-expressing proviruses interfere with infections. <u>Journal of Virology</u>. <u>40</u>:745-751 (1981).

25

30

- Rong, L., and P. Bates. Analysis of the subgroup A avian sarcoma and leukosis virus receptor: the 40-residue, cysteine-rich, low-density lipoprotein receptor repeat motif of Tva is sufficient to mediate viral entry. Journal of Virology. 69:4847-4853 (1995).
- Rong, L., K. Gendron, and P. Bates. Conversion of a human low-density lipoprotein receptor ligand-binding repeat to a virus receptor: identification of residues important for ligand specificity. <u>PNAS</u>. <u>95</u>:8467-8472 (1998).
- Rosenberg, N., and P. Jolicoeur. Retroviral pathogenesis, p. 475-585. *In* J.M. Coffin and S.H. Hughes and H.E. Varmus (ed.), Retroviruses, Cold Spring Harbor Laboratory Press (1997).
 - Ryan, U.S. Complement inhibitory therapeutics and xenotransplantation.

 Nature Medicine, 1:967-968 (1995).
- Salter, D. W., E. J. Smith, S. H. Hughes, S. E. Wright, and L. B. Crittenden. Transgenic chickens: Insertion of retroviral genes into the chicken germline. <u>Virology</u>, 157:236-240 (1987).
 - Salter, D. W., and L. B. Crittenden. Artificial insertion of a dominant gene for resistance to avian leukosis virus into the germ line of the chicken.
- 20 Theoretical Applied Genetics, 77:457-461 (1989).
 - Salter, D. W., and L. B. Crittenden. Insertion of a disease resistance gene into the chicken germline. Biotechnology, 16:125-131 (1991).
 - Sambrook, J., E.F. Fritsch, and T. Maniatis. Molecular cloning: a laboratory manual, Second edition ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1989).
 - Sandrin, M.S., W.L. Fodor, E. Mouhtouris, N. Osman, S. Cohney, S.A. Rollins, E.R. Guilmette, E. Setter, S.P. Squinto, and I.F.C. McKenzie. Enzymatic remodeling of the carbohydrate surface of a xenogeneic cell substantially reduces human antibody binding and complement-mediated cytolysis. Nature Medicine, 1:1261-1267 (1995).
 - Schacker, T., A. C. Collier, R. Coombs, J. D. Unadkat, I. Fox, J. Alam, J.-P. Wang, E. Eggert, and L. Corey. Phase I study of high-dose, intravenous

rsCD4 in subjects with advanced HIV-1 infection. Journal of Acquired Immune Deficiency Syndromes. 9:145-152 (1995).

Schaefer-Klein, J., I. Givol, E. V. Barsov, J. M. Whitcomb, M. VanBrocklin, D. N. Foster, M. J. Federspiel, and S. H. Hughes. The EV-0-derived cell line DF-1 supports efficient replication of avian leukosis-sarcoma viruses and vectors. <u>Virology</u>. 248:305-311 (1998).

Schumann, G., L. Qin, A. Rein, G. Natsoulis, and J.D. Boecke. Therapeutic Effect of *Gag*-nuclease fusion protein on retrovirus-infected cell cultures. <u>J. Virol.</u>, 70:4329-4339 (1996).

Seifarth, W., C. Baust, A. Murr, H. Skladny, F. Krieg-Schneider, J. Blusch, T. Werner, R. Hehlmann, and C. Mosch. Proviral structure, chromosomal location, and expression of HERV-K-T47D, a novel human endogenous retrovirus derived from T47D particles. <u>Journal of Virology</u>, 72:8384-8391 (1998).

Sherr, C.J., L.A. Fedele, R.E. Benveniste, and G.J. Todaro. Interspecies antigenic determinants of the reverse transcriptases and p30 proteins of mammalian type C viruses. <u>Journal of Virology</u>. <u>15</u>:1440-1448 (1975).

20

Sherr, C.J. M.M. Lieber, and G.J. Todaro. Mixed splenocyte cultures and graft versus host reactions selectively induce an "S-tropic" murine type C virus. Cell, 1:55-58 (1974).

Sielaff, T.D., M.Y. Hu, S. Rao, K. Groehler, D. Olson, H.J. Mann, R.P. Remmel, R.A.D. Shatford, B. Amiot, W.-S. Hu, and F.B. Cerra. A technique for porcine hepatocyte harvest and description of differentiated metabolic functions in static culture. <u>Transplantation</u>, <u>59</u>:1459-1463 (1995).

Simpson, G.R., C. Patience, R. Lower, R.R. Tonjes, H.D.M. Moore, R.A. Weiss, and M.T. Boyd. Endogenous D-type (HERV-K) related sequences are packaged into retroviral particles in the placenta and possess open reading frames for reverse transcriptase. <u>Virology</u>, 222:451-456 (1996).

Smith, E. J., J. Brojatsch, J. Naughton, and J. A. T. Young. The CARI gene encoding a cellular receptor specific for the subgroup B and D avian leukosis viruses maps to the chicken tvb locus. <u>Journal of Virology</u>. 72:3501-3503 (1998).

Smith, E. J., A. M. Fadly, and W. Okazaki. An enzyme-linked

immunoabsorbant assay for detecting avian leukosis sarcoma viruses. Avian Diseases. 23:698-707 (1979).

Strandstrom, H., P. Veijalainen, V. Moening, G. Hunsmann, H. Schwarz, and W. Schafer. C-type particles produced by a permanent cell line from a leukemia pig I. Origin and properties of the host cells and some evidence for the occurrence of c-type-like particles. <u>Virology</u>, 57:175-178 (1974).

Suzuka, I., K. Sekiguchi, and M. Kodama. Some characteristics of a porcine retrovirus from a cell line derived from swine malignant lymphomas. FEBS Letters, 183:124-128 (1985).

10

25

Suzuka, I., N. Shimizu, K. Sekiguchi, H. Hoshino, M. Kodama, and K. Shimotohno. Molecular cloning of unintegrated closed circular DNA of porcine retrovirus. FEBS Letters, 198:339-343 (1986).

Suzuki, T., K. Yanagihara, K. Yoshida, T. Seido, N. Kuga, Y. Shimosato, and S. Oboshi. Infectious murine type-c viruses released from human cancer cells transplanted into nude mice. Gann, 68:96-106 (1977).

Takeuchi, Y., C. Patience, S. Magre, R.A. Weiss, P.T. Banerjee, P. LeTissier, and J.P. Stoye. Host range and interference studies of three classes of pig endogenous retrovirus. <u>Journal of Virology</u>, 72:9986-9991 (1998).

Todaro, G.J., R.E. Benveniste, M.M. Lieber, and C.J. Sherr.
Characterization of a type C virus released from the porcine cell line PK (15).
Virology, 58:65-74 (1974).

Tralka, T.S., C.L. Yee, A.B. Rabson, N.A. Wivel, K.J. Stromberg, A.S. Rabson, and J.C. Costa. Murine type C retroviruses and intracisternal Aparticles in human tumors serially passaged in nude mice. <u>Journal of the National Cancer Institute</u>, 71:591-599 (1983).

Tucker, P. W., K. B. Marcu, J. L. Slightom, and F. R. Blattner. Structure of the cloned gene for the constant region of murine gamma 2b immunoglobulin heavy chain. Science. 206:1299-1306 (1979).

VanBrocklin, M., A.L. Ferris, S.H. Hughes, and M.J. Federspiel.

Expression of a murine leukemia virus Gag-Escherichia coli RNase HI fusion polyprotein significantly inhibits virus spread. <u>Journal of Virology</u>, 71:3312-3318 (1997).

Vanin, E.F., M. Kaloss, C. Broscius, and A.W. Nienhuis.

5

30

Characterization of replication-competent retroviruses from nonhuman primates with virus-induced T-cell lymphomas and observations regarding the mechanism of oncogenesis. <u>Journal of Virology</u>, <u>68</u>:4241-4250 (1994).

Ward, T., P.A. Pipkin, N.A. Clarkson, D.M. Stone, P.D. Minor, and J.W. Almond. Decay-accelerating factors CD55 is identified as the receptor for echovirus 7 using CELICS, a rapid immuno-focal cloning method. The EMBO Journal, 13:5070-5074 (1994).

Weiss, R.A. Transgenic pigs and virus adaptation. <u>Nature</u>, <u>391</u>:327-328 10 (1998).

Weiss, R. Experimental biology and assay of RNA tumor viruses, p. 209-260. *In* R. Weiss and N. Teich and H. Varmus and J. coffin (ed.), RNA Tumor Viruses, vol. 1. Cold Spring Harbor Laboratory Press, New York (1982).

Weiss, R. A. Cellular receptors and viral glycoproteins involved in retrovirus entry, p. 1-108. *In J. A. Levy* (ed.), The Retroviruses, vol. 2. Plenum Press, New York (1992).

Wilson, C., and M. Eiden. Viral and cellular factors governing hamster cell infection by murine and gibbon ape leukemia viruses, <u>J. Virol.</u>, <u>65</u>(11) 5975-5982 (1991).

Wilson, C.A., M.V. Eiden, and J.W. Marsh. Quantitative micro P30 and reverse transcriptase assay s for Moloney murine leukemia virus. Journal of Virology Methods, 48:109-118 (1994).

Wilson, C.A., S. Wong, J. Muller, C.E. Davidson, T.M. Rose and P. Burd. Type C retrovirus released from porcine primary peripheral blood mononuclear cells infects human cells. <u>Journal of Virology</u>, 72:3082-3087 (1998).

Wright et al., Biotech 9:830 (1991).

Wu, X., H. Liu, H. Xiao, J. A. Conway, and J. C. Kappes. Inhibition of human and simian immunodeficiency virus protease function by targeting Vpx-protease-mutation fusion protein into viral particles. *J. Virol.*, 10:3378-3384 (1996).

Wunderli, H., D.D. Mickey, and D.F. Paulson. C-type virus particles in human urogenital tumors after heterotransplantation into nude mice. British Journal of Cancer, 39:35-42 (1979).

Yodoi, J.K., Teshigawara, T. Nikaido, F. Fukui, T. Noma, T. Hongo, M. Takigawa, M. Sasaki, N. Minato, M. Tsudo, T. Uchiyama, AND M. Maeda.

TCGR (IL-2)-receptor inducing factor(s). I. Regulation of IL-2 receptor on natural killer-like cell line (YT cells). J. Immunol., 134, 1623-1630 (1985).

Young, J. A. T., P. Bates, and H. E. Varmus. Isolation of a chicken gene that confers susceptibility to infection by subgroup A avian leukosis and sarcoma viruses. <u>Journal of Virology</u>. <u>67</u>:1811-1816 (1993).

10

15

All publications, patents and patent applications are incorporated herein by reference. While in the foregoing specification this invention has been described in relation to certain preferred embodiments thereof, and many details have been set forth for purposes of illustration, it will be apparent to those skilled in the art that the invention is susceptible to additional embodiments and that certain of the detailed herein may be varied considerably without departing from the basic principles of the invention.

WHAT IS CLAIMED IS:

 A method to inhibit or prevent infectious agent transmission in a mammalian transplant recipient, comprising:

- a) introducing to donor swine cells a recombinant DNA comprising a promoter operably linked to a DNA segment encoding a protein comprising at least a portion of a polypeptide of the infectious agent that is present in the extracellular form of the agent so as to yield transformed swine cells; and
- b) introducing the transformed swine cells to the recipient.
 - 2. A method to inhibit or prevent infectious agent transmission in a mammalian transplant recipient, comprising:
 - a) introducing to donor human blood cells a recombinant DNA comprising a promoter operably linked to a DNA segment encoding a protein comprising at least a portion of a polypeptide of the infectious agent that is present in the extracellular form of the agent so as to yield transformed human blood cells; and
 - b) introducing the transformed human blood cells to the recipient.

20

25

15

- 3. A method to inhibit or prevent infectious agent transmission to a mammalian transplant recipient, comprising:
 - a) introducing to a donor organ a recombinant DNA comprising a promoter operably linked to a DNA segment encoding a protein comprising at least a portion of a polypeptide of the infectious agent that is present in the extracellular form of the agent so as to yield a transformed organ; and
 - b) introducing the transformed organ to the recipient.
- 30 4. The method of claim 1, 2 or 3 wherein the DNA segment encodes a fusion protein comprising at least a portion of a polypeptide of the infectious agent that is present in the extracellular form of the agent and a degradative enzyme.

5. The method of claim 4 wherein the degradative enzyme is a nuclease or protease.

6. The method of claim 1, 2 or 3 wherein the infectious agent is a virus.

5

- 7. The method of claim 6 wherein the polypeptide of the infectious agent is a viral capsid protein, viral glycoprotein or an accessory protein.
- 8. The method of claim 6 wherein the virus is a lentivirus, retrovirus, hepatitis virus or a herpesvirus.
 - 9. The method of claim 1 wherein the donor cells are embryonic stem cells, blood cells, neuronal cells, liver cells, pancreatic cells, kidney cells or islet cells.

- 10. The method of claim 3 wherein the organ is a heart, liver or kidney.
- 11. The method of claim 3 wherein the organ is a human or pig organ.
- 20 12. The method of claim 4 wherein the DNA segment encodes a fusion protein encoding a polypeptide of a pig endogenous retrovirus.
- The method of claim 5 wherein the enzyme is barnase, staphylococcal nuclease, RNase H1, RNase T1, retroviral protease, RNase III, RNaseL,
 or a ribozyme.
 - 14. The method of claim 7 wherein the polypeptide of the infectious agent is Vpr, Vpx, Vif or Nef.
- 30 15. An isolated and purified nucleic acid molecule comprising a nucleic acid segment which comprises at least a portion of a pig endogenous retrovirus, wherein the nucleic acid segment hybridizes under hybridizing

conditions to SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, or SEQ ID NO:32.

- An isolated and purified polypeptide encoded by a nucleic acid molecule comprising a nucleic acid segment comprising SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, or SEQ ID NO:32.
- 17. A method to detect human tropic pig endogenous retroviruses, 10 comprising:

15

- a) contacting a mammalian sample suspected of being infected with a pig endogenous retrovirus with a probe comprising at least a portion of SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, or SEQ ID NO:32 so as to form complexes; and
- b) detecting or determining the presence of the complexes.
- 18. A method of using an isolated and purified nucleic acid molecule comprising the genome of a pig endogenous retrovirus comprising:

 20 introducing to a host cell a recombinant DNA molecule comprising a promoter operably linked to a DNA segment comprising SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, or SEQ ID NO:32 so as to yield a transformed host cell, and identifying the transformed host cell.
 - 19. The method of claim 1, 2, 3 or 18 wherein the recombinant DNA molecule further comprises transcriptional termination sequences 3' to the DNA segment.
- 30 20. A host cell, the genome of which is augmented with a recombinant DNA molecule comprising a promoter operably linked to a DNA segment encoding a fusion protein comprising at least a portion of a polypeptide of a pig endogenous retrovirus and a degradative enzyme.

A fusion protein, comprising: a capsid or envelope protein of a porcine 21. endogenous virus and a degradative enzyme. 22. The fusion protein of claim 21 wherein the enzyme is a nuclease. 5 The fusion protein of claim 21 wherein the enzyme is a protease. 23. 24. The fusion protein of claim 21 wherein the enzyme is a lipase. The fusion protein of claim 21 wherein the activity of said enzyme is 10 25. calcium-dependent. The fusion protein of claim 25 wherein the enzyme is staphylococcal 26. nuclease. 15 An isolated and purified DNA molecule encoding the fusion protein of 27. claim 21. A recombinant virus comprising a nucleic acid molecule encoding the 28. 20 fusion protein of claim 21. An antibody that specifically binds pig endogenous retrovirus. 29. The antibody of claim 29 which binds a viral capsid protein. 30. 25 The antibody of claim 29 which binds the viral envelope glycoprotein. 31. The method of claim 8 wherein the virus is Epstein Barr virus. 32. The method of claim 8 wherein the virus is cytomegalovirus. 30 33. The method of claim 8 wherein the virus is human immunodeficiency 34.

virus.

The method of claim 1, 2, 3 or 18 wherein the recombinant DNA is introduced to the recipient by infection with a recombinant virus.


36. A method to inhibit or prevent infectious agent transmission in a mammalian transplant recipient, comprising:


5

10

20

- a) introducing to donor mammalian cells a recombinant DNA comprising a promoter operably linked to a DNA segment encoding a protein comprising at least a portion of a polypeptide of the infectious agent that is present in the extracellular form of the agent so as to yield transformed mammalian cells; and
- b) introducing the transformed mammalian cells to the recipient.
- 37. A method to inhibit or prevent infectious agent transmission in a mammalian transplant recipient, comprising:
- a) introducing to donor human cells a recombinant DNA encoding a protein comprising at least a portion of a polypeptide of the infectious agent that is present in the extracellular form of the agent so as to yield transformed human cells; and
 - b) introducing the transformed human cells to the recipient.
 - 38. The method of claim 36 or 37 wherein the infectious agent is a virus.
 - 39. The method of claim 38 wherein the polypeptide of the infectious agent is a viral capsid protein, viral glycoprotein or an accessory protein.





SUBSTITUTE SHEET (RULE 26)




FIG. 1C




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6A

				/28 00 4	000	000	000
50 50 50	100 100 100	150 150 147	200 200 185	250 250 234	300 300 279	350 350 329	400 400 379
VNGKRLVDSP TM.IGL	LRSVIPGLND	TSNDGNWKWP	NKQISCHSLD TGSPK.SPS.	VLTIRLRIETK.N-	PTEPNITIKTS.S.T.M	ARGGKFNVTK 	EAFNRTSESQ Q
WFLTLSITPQ	GTWWPELYVC	DFFCKQWSCV I	WQQRVQKDVR LTWI.	YRGSGRKKGS .G	SDYNTTSGSV	ASGPPYYEGM	PSHQHLCNHT
KIPLSFASIA	INSTQGEAPL	NNEEYCGNPQ GKHR	FNYGHGRWKD	VNGMSWGIVY I	IQEQRPSPNP TG.G.	EATSSCWLCL	GKGTCIGMVP
PIRGGKPKRL	LLTDSGTGIN	YGFYVCPGPP	PVNNPTSYNQ	KGKQENIQKW	NKGLAEQGPP 	AFQALNSTTP	QNKLTLTEVS
MHPTLSRRHL	NSHKPLSLTW	QATPPDVLRA	VSQQDRVSYS TF.	LDYLKISFTE	QMEPPVAIGP	GAKLFNLIQG	EHRDQCTWGS
PERV-1.15 PERV-A	PERV-C PERV-1.15 PERV-A	PERV-C PERV-1.15 PERV-A	PERV-1.15 PERV-A PERV-C	PERV-1.15 PERV-A	PERV-C PERV-1.15 PERV-A	PERV-C PERV-1.15 PERV-A	PERV-1.15 PERV-A PERV-C

			8/28		
450 450 429	500 500 479	550 550 529	600	650 650 629	659 660 638
VYYYPEKAVL	GPQQLEKGLS 	LFLKEGGLCV	FEGWFNRSPW	QIMVLRQQYQ	
FCVMVQIVPR	VGTGTAALIT	VLQNRRGLDL	RREKETTQGWR.AD	AFIRERISAVVV	
STLVFNQTKD	VMLGLGVAAG	EESLTSLSEV	NKLRERLEKR SR.	VGPCI INKLIL.RFV	
ACNTGLTPCV	KREPISLTLA	OALEKSVSNL	DHSGAIRDSM	PLIVLLLLT 	
YLVPGYDRWW	DEYDYRYNRP 	NLHRIVTENLD.	ALKEECCFYV	LATLLSALTG MT	SPSSREAGR GLL.QGETDL
PERV-1.15 PERV-A PERV-C	PERV-1.15 PERV-A PERV-C	PERV-1.15 PERV-A PERV-C	PERV-1.15 PERV-A PERV-C	PERV-1.15 PERV-A PERV-C	PERV-1.15 PERV-A PERV-C

FIG. 6B (CONTINUED)




FIG. 7

]	L0/28				
50 31 100 81		150 113	200	250	300	350	400	450	500 429
AGGAAGTTAA C	CTATAGGTAA T	CTGTACAAA	CTATGCTTGA GA	TAAACCTGCT .CT.A	CCCCAGTAAG	ACAACATGTC	CAATTTAAAT	GACTGGTTTG	TTGTGAACCC
CCCAGAACCC	ACCCTGTTCC .GTT	GCTTTCTGCT GTATGTGCCTCAT		TTGGCTTCTG AAA.GGAG.T	GCCTAACGGA GCC.TAA.	TATCTTGGTG	AACTCTAAAA	GTTTTAAATT	GCGGGCTTTG
ACCTGACTCT	CGAATTCCAG TTTAAAATAT AGGCCCTGA		AGGCCCCTGA	AACTGCTTGT G.TAA.AA.A	GAAGTCAATT AACTGG	TTGAAACACA	GCAAATGTGT	GGCTCTCGAA	TTTGTAAAGC
TGAAAATGCA A TAAATGCCCT		TTTTTGCTGT	GGGGAAAAAC . AA.TT	AACTGCTCCT	AAAAAGAGGA CT.T	CCACAAAATG TG	CGAAACATGC	ACGAAGCGCG	AAATGATTGG
→ U3 AATGAAAGGA	TAAGAAGCTC GT AAGATCATAC		CTTTGTGGAA	AACTTCTTGA	AACTTCTTGATGCATAAGAT		TCCCCCACCC	TAATTGGTCC	•
<u>-</u>	PERV-MSL LamAl	PERV-MSL LamAl	PEKV-MSD LamAl	PERV-MSL LamAl	LamA1	LamA1	LamA1	LamAl	LamA1 PERV-MSL

11/	′28
-----	-----

LamA1	CATAAAAGCT	GTCCCGACTC	CACACTCGGG	GCCGCAGTCC	TCTACCCCTG	550 479
PERV-MSL					C E C C E	
LamA1	CGTGGTGTAC	GACTGTGGGC	CCCAGCGCGC	TTGGAATAAA	AAICCICIIG	529
PERV-MSL	:					
LamA1	CTGTTTGCAT	R←I→us CAAGACCGCT	TCTCGTGAGT	GATTAAGGGG	AGICGCCTIT	650
PERV-MSL	:			ord Pro		\ -)
LamA1	TCCGAGCCTG	GAGGTTCTTT	TIGCIAGICI	TACATTTGGG	GGCTCGTCCG	700
PERV-MSL			g)
LamA1	← GGAT					704
PERV-MSL	:					

FIG. 8 (CONTINUED)




FIG. 9A







FIG. 9B





SUBSTITUTE SHEET (RULE 26)



SUBSTITUTE SHEET (RULE 26)




FIG. 13







FIG. 14







SUBSTITUTE SHEET (RULE 26)

Figure 18

PERV 1.15

ATGCATCCCACGTTAAGCCGGCGCCACCTCCCGATTCGGGGTGGAAAGCCGAAAAGACTGAAAATCCCCTTAAGCTTCGC CTCCATCGCGTGGTTCCTTACTCTGTCAATAACTCCTCAAGTTAATGGTAAACGCCTTGTGGACAGCCCGAACTCCCATA AACCCTTATCTCTCACCTGGTTACTTACTGACTCCGGTACAGGTATTAATATTAACAGCACTCAAGGGGAGGCTCCCTTG GGGACCTGGTGGCCTGAATTATATGTCTGCCTTCGATCAGTAATCCCTGGTCTCAATGACCAGGCCACACCCCCCGATGT ACTCCGTGCTTACGGGTTTTACGTTTGCCCAGGGCCCCCAAATAATGAAGAATATTGTGGAAATCCTCAGGATTTCTTTT GCAAGCAATGGAGCTGCGTAACTTCTAATGATGGGAATTGGAAATGGCCAGTCTCTCAGCAAGACAGAGTAAGTTACTCT TTTGTTAACAATCCTACCAGTTATAATCAATTTAATTATGGCCATGGGAGATGGAAAGATTGGCAACAGCGGGTACAAAA GTTCTGACTATTCGCCTCAGAATAGAAACTCAGATGGAACCTCCGGTTGCTATAGGACCAAATAAGGGTTTGGCCGAACAAGGACCTCCAATCCAAGAACAGAGGCCATCTCCTAACCCCTCTGATTACAATACAACCTCTGGATCAGTCCCCACTGAGC CTAACATCACTATTAAAACAGGGGCGAAACTTTTTAACCTCATCAGGGGGGCTTTTCAAGCTCTTAACTCCACGACTCCA GAGGCTACCTCTTCTTGTTGGCTTAGCTTCGGGCCCACCTTACTATGAGGGAATGGCTAGAGGAGGGAAATTCAA TGTGACAAAGGAACATAGAGACCAATGTACATGGGGATCCCAAAATAAGCTTACCCTTACTGAGGTTTCTGGAAAAGGCA CCTGCATAGGGATGGTTCCCCCATCCCACCAACACCTTTGTAACCACCTGAAGCCTTTAATCGAACCTCTGAGAGTCAG
TATCTGGTACCTGGTTATGACAGGTGGTGGGCATGTAATACTGGATTAACCCCTTTGTGTTTCCACCTTTGGTTTTCAACCA AACTAAAGACTTTTGCGTTATGGTCCAAATTGTCCCCCGGGTGTACTACTACCCCTTGGTTTTCACCCAAAAAGACTTTTGCGTTATGATGAATATGACCACTTGATGAATATGACTATAAGATATAATCGGCCAAAAAAGAGACCCCAATCCCCTGACACAACAGCTGCAAAAAGCAGTCCTTGATGAATATGACCTGACAGAAAAAGCAGACCGCAAAAAAGCAGCTGGAGAAAAAGCAGACTGCAGGACTGCAGCGCGCAACAGCTGGAGAAAAAGCACTTAGTAACCTACATCGAATTGTAAC GGAAAATCTCCAAGCCCTAGAAAAATCTGTCAGTAACCTGGAGGAATCCCTAACCTCCTTATCTGAAGTGGTTCTACAGA ACAGAAGGGGGTTAGATCTGTTATTTCTAAAAGAAGAAGGAGGATTATGTGTAGCCTTAAAAGAAGAAGAAGGATTTTTTATGTG GATCATTCAGGGGCCATCAGAGACTCCATGAACAAGCTTAGAGAAAGGTTGGAGAAGCGTCGAAGGGAAAAGGAAACTAC TCCTCCTCCTGTTACTCACAGTTGGGCCATGTATTATTAACAAGTTAATTGCCTTCATTAGAGAACGAATAAGTGCAGTC CAGATCATGGTACTTAGACAACAGTACCAAAGCCCGTCTAGCAGAGAAGCTGGCCGCTAG

LamA

TACTTCTTGGGGAAGACCCTGGGCTGCTAACTGGGTCTTGGCTGGTCCTAGTGAAAGGATGAAAATGCAACCTGACTCTC CCAGAACCCAGGAAGTTAATAAGAAGCTCTAAATAATGAAAGGATGAAAATGCAACCTGACTCTCCCAGAACCCAGGAAG TTAATAAGAAGCTCTAAATGCCCTCGAATTCCAGACCCTGTTCCCTATAGGTAAAAGATCATACTTTTTGCTGTTTTAGG GCTTGCTTTCTGCTCTGTACAAAACTTTGTGGAAGGGGAAAAACAGGCCCCTGAGTATGTGCCTCTATGCTTGAAACTTC TTGAAACTGCTCCTAACTGCTTGTTTGGCTTCTGTAAACCTGCTTGCATAAGATAAAAAGAGGAGAAGTCAATTGCCTAA CGGACCCCAGTAAGATCGGGTGTACCACAAAATGTTGAAACACATATCTTGGTGACAACATGTCTCCCCCACCCCGAAAC ATGCGCAAATGTGTAACTCTAAAACAATTTAAATTAATTGGTCCACGAAGCGCGGGCTCTCGAAGTTTTAAATTGACTGG TTTGTGATATTTTGAAATGATTGGTTTGTAAAGCGCGGGCTTTGTTGTGAACCCCATAAAAGCTGTCCCGACTCCACACT CGGGGCCGCAGTCCTCTACCCCTGCGTGGTGTACGACTGTGGGCCCCAGCGCGCTTGGAATAAAAATCCTCTTGCTGTTT GCATCAAGACCGCTTCTCGTGAGTGATTAAGGGGAGTCGCCTTTTCCGAGCCTGGAGGTTCTTTTTGCTAGTCTTACATT TGGGGGCTCGTCCGGGATCTGTCGCGGCCACCCCTAACACCCGAGAACCGACTTGGAGGTAAAAAGGATCCTCTTTTTAA CGTGTATGCATGTACCGGCCGGCGTCTCTGTTCTGAGTGTCTGTTTTCAGTGGTGCGCGCTTTCGGTTTGCAGCTGTCCT TCCCCCTCCGCGGCCGTCCGACTCTTTTGCCTGCTTGTGGAAGACGCGGACGGGTCGCGTGTGTCTGGATCTGTTGGTTT CTGTTTTGTGTGTCTTTGTCTTGTCCTTGTCTACAGTTTTAATATGGGACAGACGGTGACGACCCCTCTTAGTTTG ACTCTCGACCATTGGACTGAAGTTAAATCCAGGGCTCATAATTTGTCAGTTCAGGTTAAGAAGGGACCTTGGCAGACTTT CTGTGTCTCTGAATGGCCGACATTCGATGTTGGATGGCCATCAGAGGGGACCTTTAATTCTGAGATTATCCTGGCTGTTA AAGCAATTATTTTCAGACTGGACCCGGCTCTCATCCCAATCAGAGGCCCTATATCCTTACGTGGCAGATTTGGCAGAG GATCTCCCGCCATGGGTTAAACCTTGGCTGAATAAGCCAAGAAGCCAGGTCCCCGAATTCTGGCTCTTTGGAGAGAAAAAA CAAACACTCGGCTGAAAAAGTCAAGCCCTCTCCTCATATCTACCCCGAGATTGAGGAGCCGCCGGCTTGGCCGGAACCCC AATCTGTTCCCCCACCCCCTTATCTGGCACAGGGTGCTGCGAGGGGACCCCTCTGCCCCTCCTGGAGCTCCCGGCGGTGGAG GGACCTGCTGCAGGGACTCGGAGCCGGAGGGGCCCCCCCGGAGCGGACAGACGAGATCGCGACATTACCGCTGCGCACGTACGGCCCTCCCACACCCGGGGGGCCAATTGCAGCCCCTCCAGTATTGGCCCTTTTCTTCTGCAGATCTCTATAATTGGA AAACTAACCATCCCCCTTTCTCGGAGGATCCCCAACGCCTCACGGGGTTGGTGGAGTCCCTTATGTTCTCTCACCAGCCT ACTTGGGATGATTGTCAACAGCTGCTGCAGACACTCTTCACAACCGAGGAGCGAGAGAATTCTGTTAGAGGCTAGAAA AAATGTTCCTGGGGCCGACGGGCGACCCACGCAGTTGCAAAATGAGATTGACATGGGATTTCCCTTGACTCGCCCCGGTTGGGACTACAACACGCTGAAAGGTAGGGAGAGCTTGAAAAATCTATCGCCAGGCTCTGGTGGCGGGTCTCCGGGGCGCCCCCA AGACGGCCCACTAATTTGGCTAAGGTAAGAGAAGTGATGCAGGGACCGAATGAACCCCCCTCTGTTTTCCTTGAGAGGCT CTTGGAAGCCTTCAGGCGGTACACCCCTTTTGATCCCACCTCAGAGGCCCAAAAAGCCTCAGTGGCTTTGGCCTTTATAG GACAGTCAGCCTTGGATATTAGAAAGAAGCTTCAGAGACTGGAAGGGTTACAGGAGGCTGAGTTACGTGATCTAGTGAAG GAGGCAGAGAAAGTATTACAAAAGGGAGACAGAAGAAGAAGAACAAAGAAGAAGAGAGAAAGGAAAGGAAAGGAAAGGAAAGGAAAGGAAAGGGAAAGGAAAGGGAAAGGAAGGACCAATGGGAAAAGGAAATTAGGTCAGGCCCTAGACAGTCAGGGAACCTGGGCAATAGGACCCCAAGAAGGACCCAATGTTAAAAGAAAAAGGACACTGGGCAAGGAACCCCAAGAAGGAACAAAGGAACAAAGGACCTTAGCCCCAAGAAGGGAAAAAAGGAACAAAGGACTGAAGGTCTTAGC TCTGAAAGAAGATAAAGACTAGGGAAGACGGGGTTCGGACCCCCTCCCCGAGCCCAGGGTAACTTTGAAGGTGGAGGGGC AACCAGTTGAGTTCCTGGTTGATACCGGAGCGAAACATTCAGTGCTACTACAGCCATTAGGAAAAACTAAAAAGATAAAAAA

TCCTGGGTGATGGGTGCCACAGGGCAACAACAGTATCCATGGACTACCCGAAGAACAGTTGACTTGGGAGTGGGACGGGT AACCCACTCGTTTCTGGTCATACCTGAGTGCCCAGCACCCCTCTTAGGTAGAGACTTATTGACCAAGATGGGAGCACAAA TTTCTTTTGAACAAGGGAAACCAGAAGTGTCTGCAAATAACAAACCTATCACTGTGTTGACCCTCCAATTAGATGACGAA TATCGACTATACTCTCCCCTAGTAAAGCCTGATCAAAATATACAATTCTGGTTGGAACAGTTTCCCCAAGCCTGGGCAGA GACAGTACCCCTTGAGTAAAGAAGCTCAAGAAGGAATTCGGCCGCATGTCCAAAGATTAATCCAACAGGGCATCCTAGTT CCTGTCCAATCTCCCTGGAATACTCCCCTGCTACCGGTTAGAAAGCCTGGGACTAATGACTATCGACCAGTACAGGACTT GAGAGAGGTCAATAAACGGGTGCAGGATATACACCCAACAGTCCCGAACCCTTATAACCTCTTGTGTGCTCTCCCACCCC AACGGAGCTGGTATACAGTATTGGACTTAAAGGATGCCTTTTTCTGCCTGAGATTACACCCCACTAGCCAACCACTTTTT GCCTTCGAATGGAGAGATCCAGGTACGGGAAGAACCGGGCAGCTCACCTGGACCCGACTGCCCCAAGGGTTCAAGAACTC CCCGACCATCTTTGACGAAGCCCTACACAGAGACCTGGCCAACTTCAGGATCCAACACCCTCAGGTGACCCTCCCAGT ACGTGGATGACCTGCTTCTGGCGGGAGCCACCAAACAGGACTGCTTAGAAGGCACGAAGGCACTACTGCTGGAATTGTCT GACCTAGGCTACAGAGCCTCTGCTAAGAAGGCCCAGATTTGCAGGAGAGAGGTAACATACTTGGGGTACAGTTTGCGGGA CGGGCAGCGATGGCTGACGGAGGCACGGAAGAAAACTGTAGTCCAGATACCGGCCCCAACCACAGCCAAACAAGTGAGAG AGTTTTTGGGGACAGCTGGATTTTGCAGACTGTGGATCCCGGGGTTTGCGACCTTAGCAGCCCCACTCTACCCACTAACC AAAGAAAAAGGGGAATTCTCCTGGGCTCCTGAGCACCAGAAGGCATTTGATGCTATCAAAAAAGGCCCTGCTGAGCGCACC TGCTCTGGCCCTCCCTGACGTAACTAAACCCTTTACCCTTTATGTGGATGAGCGTAAGGGAGTAGCCCGGGGAGTTTTAA CCCAAACTCTAGGACCATGGAGGAGACCTGTTGCCTACCTGTCAAAGAAGCTCGATCCTGTAGCCAGTGGTTGGCCCGTA TGCCTGAAGGCTATCGCAGCTGTGGCCATACTGGTCAAGGACGCTGACAAATTGACTTTGGGACAGAATATAACTGTAAT GCCTGCTTCTCACAGAGAGGGTCACGTTCGCTCCACCAGCCGCTCTCAACCCTGCCACTCTTCTGCCTGAAGAGACTGAT GAACCAGTGACTCATGATTGCCATCAACTATTGATTGAGGAGACTGGGGTCCGCAAGGACCTTACAGACATACCGCTGAC TGGAGAAGTGTTAACCTGGTTCACTGACGGAAGCAGCTATGTGGTGGAAGGTAAGAGGATGGCTGGGGCGGCGGTGGTGG AAGCCTTACATTTGCCAAAAAGGCTAGCTATTATACACTGTCCTGGACATCAGAAAGCCAAAGATCCCATATCCAGAGGG AACCAGATGGCTGACCGGGTTGCCAAGCAGGCAGCCCAGGGTGTTAACCTTCTGCCTATGATAGAAACACCCAAAGCCCC CACCTAGGAACTAAACACCTGCAGCAGTTGGTCAGAACATCTCCTTATCATGTTCTGAGGCTACCAGGAGTGGCTGATTCGGTGGTCAAACACTGTGTGCCCTGCCAGCTGGTTAATGCTAATCCTTCCAGAATACCTCCAGGAAAGAGACTAAGGGGAA GCCACCCAGGCGCTCACTGGGAAGTGGACTTCACTGAGGTAAAGCCGGCTAAATACGGAAACAAATATCTATTGGTTTTT AATAGAACCATTAAAGAGACCCTTACTAAATTGACCGCGGAGACTGGCGTTAATGATTGGATAGCTCTCCTGCCCTTTGT ACTITITAGGGTTAGGAACACCCCTGGACAGTTTGGGCTGACCCCCTATGAATTACTCTACGGGGGACCCCCCCATTGG TAGAAATTGCTTCCGTACATAGTGCTGACGTGCTGCTTTCCCAGCCTTTGTTCTCTAGGCTCAAGGCACTTGAGTGGGTG AGACAACGAGCGTGGAGGCAACTCCGGGAGGCCTACTCAGGAGGAGGAGACTTGCAGATCCCACATCGTTTCCAAGTGGG AGATTCAGTCTACGTTAGACGCCACCGTGCAGGAAACCTCGAGACTCGGTGGAAGGGCCCTTATCACGTACTTTTGACCA CACCAACGGCTGTGAAAGTCGAAGGAATCTCCACCTGGATCCATGCATCCCACGTTAAGCCGGCGCCACCTCCCGATTCG ACAGGTATTAATATTAACAGCACTCAAGGGGAGGCTCCCTTGGGGACCTGGTGGCCTGAATTATATGTCTGCCTTCGATCAGTAATCCCTGGTCTCAATGACCAGGCCACACCCCCCGATGTACTCCGTGCTTACGGGTTTTACGTTTGCCCAGGACCCC TGGAAATGGCCAGTCTCTCAGCAAGACAGAGTAAGTTACTCTTTTGTTAACAATCCTACCAGTTATAATCAATTTAATTA TGGCCATGGGAGATGGAAAGATTGGCAACAGCGGGTACAAAAAGATGTACGAAATAAGCAAATAAGCTGTCATTCGTTAG ACCTAGATTACTTAAAAATAAGTTTCACTGAAAAAGGAAAACAAGAAAATATTCAAAAGTGGGTAAATGGTATGTCTTGG GGAATAGTGTACTATGGAGGCTCTGGGAGAAAGAAAGGATCTGTTCTGACTATTCGCCTCAGAATAGAAACTCAGATGGA ACCTCCGGTTGCTATAGGACCAAATAAGGGTTTGGCCGAACAAGGACCTCCAATCCAAGAACAGAGGCCATCTCCTAACC CCTCTGATTACAATACAACCTCTGGATCAGTCCCCACTGAGCCTAACATCACTATTAAAACAGGGGCGAAACTTTTTAACCTCATCCAGGGGAGCTTTTCAAGCTCTTAACTCCACGACTCCAGAGGCTACCTCTTCTTGTTGGCTTTGCTTAGCTTCGGG CCCACCTTACTATGAGGGAATGGCTAGAGGAGGGAAATTCAATGTGACAAAGGAACATAGAGACCAATGTACATGGGGAT

LamA10

AGTCCCGAACCCTTATAACCTCTTGTGTGCTCTCCCACCCCAACGGAGCTGGTATACAGTATTGGACTTAAAGGATGCCT TITTCTGCCTGAGATTACACCCCACTAGCCAACCACTTTTTGCCTTCGAATGGAGAGATCCAGGTACGGGAAGAACCGGG CAGCTCACCTGGACCCGACTGCCCCAAGGGTTCAAGAACTCCCCGACCATCTTTGACGAAGCCCTACACAGAGACCTGGC CAACTTCAGGATCCAACACCCTCAGGTGACCCTCCCAGTACGTGGATGACCTGCTTCTGGCGGGAGCCACCAAACAGG CAACTTCAGGATCCAACACCCTCAGGTGACCCTCCTCCAGTACGTGGATGACCTGCTTCTGGCGGGAGCCACCAAACAGG ACTGCTTAGAAGGCACGAAGGCACTACTGCTGGAATTGTCTGACCTAGGCTACAGAGCCTCTGCTAAGAAGGCCCAGATT TGCAGGAGAGAGGTAACATACTTGGGGTACAGTTTGCGGGACGGGCAGCGATGGCTGACGGAGGCACGGAAGAAAACTGT AGTCCAGATACCGGCCCCAACCACAGCCAAACAAGTGAGAGAGTTTTTGGGGGACAGCTGGATTTTGCAGACTGTGGATCC CGGGGTTTGCGACCTTAGCAGCCCCACTCTACCCACTAACCAAAGAAAAAGGGGAATTCTCCTGGGCTCCTGAGCACCAG AAGGCATTTGATGCTATCAAAAAGGCCCTGCTGAGCGCACCTGCTCTGGCCCTCCTGACGTAACCCTTTACCCT TTATGTGGATGAGCGTAAGGGAGTAGCCCGGGGAGTTTTAACCCAAACTCTAGGACCATGGAGGAGACCTGTTGCCTACC TGTCAAAGAAGCTCGATCCTGTAGCCAGTGGTTGGCCCGTATGCCTGAAGGCTATCGCAGCTGTGGCCATACTGGTCAAG GACGCTGACAAATTGACTTTGGGACAGAATATAACTGTAATAGCCCCCCATGCGTTGGAGAACATCGTTCGGCAGCCCCC AGACCGATGGATGACCAACGCCCGCATGACCCACTATCAAAGCCTGCTTCTCACAGAGAGGGGTCACGTTCGCTCCACCAG AAGAAGGGTTAGAATATGTCCAACAGATACATCGTCTAACCCACCTAGGAACTAAACACCTGCAGCAGTTGGTCAGAACA
TCTCCTTATCATGTTCTGAGGCTACCAGGAGTGGCTGATTCGGTGGTCAAACACTGTGTGCCCTGCCAGCTGGTTAATGC TAATCCTTCCAGAATACCTCCAGGAAAGAGACTAAGGGGAAGCCACCCAGGCGCTCACTGGGAAGTGGACTTCACTGAGG
TAAAGCCGGCTAAATACGGAAACAATATCTATTGGTTTTTGTAGACACCTTTTCAGGATGGGTAGAGGCTTATCCTACT
AAGAAAGAGACTTCAACCGTGGTGGCTAAGAAAATACTGGAGGAAATTTTTCCAAGATTTTGCAAGATATACGTAATAGG
GTCAGACAATGGTCCAGCTTTCGTTGCCCAGGTAAGTCAGGGAATACTGCAAGATATTTGGAGGATTTGGAAACTGCATT CCCAGCCTTTGTTCTCTAGGCTCAAGGCACTTGAGTGGGTGAGACAACGAGCGTGGAGGCAACTCCGGGAGGCCTACTCA GGAGGAGGAGACTICCAGATCCACATCGTTTCCAAGTGGAGATCACGTTAGACGCAGCATCCACGTGAAGACTCCGGGAGGCCTACTCA GGAGGAGGAGGAGACTIGCAGATCCCACATCGTTTCCAAGTGGGAGATTCAGTCTACGTTAGACGCCACCGTGCAGGAAAACCT CGAGACTCGGTGGAAGGGCCCTTATCACGTACTTTTGACCACACCAACGGCTGTGAAAGTCCGAAGGAATCTCCACCTGGA TCCATGCATCCCACCGTTAAGCCGGCGCCACTCCCCGATTCGGGGTGGAAAGCCGAAAAGACTGAAAATCCCCTTAAGCTT TGTACTCCGTGCTTACGGGTTTTACGTTTGCCCAGGACCCCCAAATAATGAAGAATATTGTGGAAATCCTCAGGATTTCT TTTGCAAGCAATGGAGCTGCGTAACTTCTAATGATGGGAATTGGAAATGGCCAGTCTCTCAGCAAGACAGAGTAAGTTAC TCTTTTGTTAACAATCCTACCAGTTATAATCAATTTAATTATGGCCATGGGAGATGGAAAGATTGGCAACAGCGGGTACA AAAAGATGTACGAAATAAGCAAATAAGCTGTCATTCGTTAGACCTAGATTACTTAAAAATAAGTTTCACTGAAAAAGGAA

LamAll

GGAGTTTGAGTTTTATCGAATTTGAAACAGTGGTTTACATGGAGATTGTAGTGAAAGGATGAAAATGCAACCTGACTCTC CCAGAACCCAGGAAGTTAATAAGAAGCTCTAAATGCCCTCGAATTCCAGACCCTGTTCCCTATAGGTAAAAGATCATACT TTTTGCTGTTTTAGGGCTTGCTTTCTGCTCTGTACAAAACTTTGTGGAAGGGGAAAAACAGGCCCCTGAGTATGTGCCTC TATGCTTGAAACTTCTTGAAACTGCTCCTAACTGCTTGTTTTGGCTTCTGTAAACCTGCTTGCATAAGATAAAAAGAGGAG AAGTCAATTGCCTAACGGACCCCAGTAAGATCGGGTGTACCACAAAATGTTGAAACACATATCTTGGTGACAACATGTCT TITTAAATTGACTGGTTTGTGATATTTTGAAATGATTGGTTTGTAAAGCGCGGGCTTTGTTGTGAACCCCATAAAAGCTG TCCCGACTCCACACTCGGGGCCGCAGTCCTCTACCCCTGCGTGGTGTACGACTGTGGGCCCCCAGCGCGCTTGGAATAAAA TGGATCTGTTGGTTTCTGTGTGTCTTTGTCTTGTGCGTCCTTGTCTACAGTTTTAATATGGACAGACGGTGACGACGCCCTCTTAGTTTGACTCTCGACCATTGGACTGAAGTTAAATCCAGGGCTCATAATTTGTCAGTTCAGGTTAAGAAGAGGG ACCITGGCAGACTTTCTGTGTCTCTGAATGGCCGACATTCGATGTTGGATGGCCATCAGAGGGGACCTTTAATTCTGAGA TTATCCTGGCTGTTAAAGCAATTATTTTTCAGACTGGACCCGGCTCTCATCCCAATCAGGAGCCCTATATCCTTACGTGG CAAGATTTGGCAGAGGATCCTCCGCCATGGGTTAAACCTTGGCTGAATAAGCCAAGAAAGCCAGGTCCCCGAATTCTGGC TCTTGGAGAGAAAACAAACACTCGGCTGAAAAAGTCAAGCCCTCTCCTCATATCTACCCCGAGATTGAGGAGCCGCCGG ATCTCTATAATTGGAAAACTAACCATCCCCCTTTCTCGGAGGATCCCCAACGCCTCACGGGGTTGGTGGAGTCCCTTATG

GTTAGAGGCTAGAAAAAATGTTCCTGGGGCCGACGGGCGACCCACGCAGTTGCAAAATGAGATTGACATGGGATTTCCCT TGACTCGCCCCGGTTGGGACTACAACACGGCTGAAAGGTAGGGAGAGCTTGAAAATCTATCGCCAGGCTCTGGTGGCGGGT CGTGATCTAGTGAAGGAGGCAGAGAAAGTATATTACAAAAGGGAGACAGAAGAAAGGGGAACAAAGAAAAGAGAGAGAGA TGAAGGTGGAGGGCAACCAGTTGAGTTCCTGGTTGATACCGGAGCGAAACATTCAGTGCTACTACAGCCATTAGGAAAA CTAAAAGATAAAAAATCCTGGGTGATGGGTGCCACAGGGCAACAACAGTATCCATGGACTACCCGAAGAACAGTTGACTT GGGAGTGGGACGGGTAACCCACTGGTTTCTGGTCATACCCGAGGACTGCCCCACAAGTTATCCACGGAGAGACAGTTGACCA
AGATGGGAGCACAAATTTCTTTTGAACAAGGGAAACCAGAAGTGTCTGCAAATAACAAACCTATCACTGTGTTGACCCTC
CAATTAGATGACGAATATCGACTATACTCTCCCCTAGTAAAAGCCTGATCAAAATATACAATTCTGGTTGGAACAGTTTCC
CCAAGCCTGGCCAGAAACCGCAGGGATGGGTTTGGCAAAGCAAGTCCCCCACAAGTTATTCAACTGAAGGCCAGTGCCA CACCAGTGTCAGTCAGACAGTACCCCTTGAGTAAAGAAGCTCAAGAAGGAATTCGGCCGCATGTCCAAAGATTAATCCAA CAGGGCATCCTAGTTCCTGTCCAATCTCCCTGGAATACTCCCCTGCTACCGGTTAGAAAGCCTGGGACTAATGACTATCG ACCAGTACAGGACTTGAGAGAGGTCAATAAACGGGTGCAGGATATACACCCAACAGTCCCGAACCCTTATAACCTCTTGT CTCTACCCACTAACCAAAGAAAAAGGGGAATTCTCCTGGGCTCCTGAGCACCAGAAGGCATTTGATGCTATCAAAAAGGC AGTGGTTGGCCCGTATGCCTGAAGGCTATCGCAGCTGTGGCCATACTGGTCAAGGACGCTGACAAATTGACTTTGGGACA AACACCCAAAGCCCCAGAACCCGGACGACAGTACACCCTAGAAGACTGGCAAGAGATAAAAAAGATAGACCAGTTCTCTG ATACATCGTCTAACCCACCTAGGAACTAAACACCTGCAGCAGTTGGTCAGAACATCTCCTTATCATGTTCTGAGGCTACC AGGAGTGGCTGATTCGGTGGTCAAACACTGTGTGCCCTGCCAGCTGGTTAATGCTAATCCTTCCAGAATACCTCCAGGAA AGAGACTAAGGGGAAGCCACCCAGGCGCTCACTGGGAAGTGGACTTCACTGAGGTAAAGCCGGCTAAATACGGAAACAAA CAGGTAGAGAGGATGAATAGAACCATTAAAGAGACCCTTACTAAATTGACCGCGGAGACTGGCGTTAATGATTGGATAGC TCTCCTGCCCTTTGTGCTTTTTAGGGTTAGGAACACCCCTGGACAGTTTGGGCTGACCCCCTATGAATTACTCTACGGGG GACCCCCCCATTGGTAGAAATTGCTTCCGTACATAGTGCTGACGTGCTGCTGTTCCCAGCCTTTGTTCTCTAGGCTCAAG GCACTTGAGTGGGTGAGACAACGAGCGTGGAGGCAACTCCGGGAGGCCTACTCAGGAGGAGAGACTTGCAGATCC

LamA3A

ACCTTGGCAGACTTTCTGTGTCTCTGAATGGCCGACATTCGATGTTGGATGGCCATCAGAGGGGACCTTTAATTCTGAGA TTATCCTGGCTGTTAAAGCAATTATTTTTCAGACTGGACCCGGCTCTCATCCCAATCAGGAGCCCTATATCCTTACGTGG CAAGATTTGGCAGAGGATCCTCCGCCATGGGTTAAACCTTGGCTGAATAAGCCAAGAAAGCCAGGTCCCCGAATTCTGGC TCTTGGAGAGAAAAACAACACTCGGCTGAAAAAGTCAAGCCCTCTCCTCATATCTACCCCGAGATTGAGGAGCCGCCGGCTTGGCCCGGAAACCCCAATCTGTTCCCCCACCCCTTATCTGGCACAGGGTGCTGCGAGGGGACCCTCTGCCCCTCCTGGA TGACTCGCCCCGGTTGGGACTACAACACGGCTGAAGGTAGGGAGAGCTTGAAAATCTATCGCCAGGCTCTGGTGGCGGGT CTCCGGGGCGCCTCAAGACGGCCCACTAATTTGGCTAAGGTAAGAGAAGTGATGCAGGGACCGAATGAACCCCCCTCTGT TTTCCTTGAGAGGCTCTTGGAAGCCTTCAGGCGGTACACCCCTTTTGATCCCACCTCAGAGGCCCAAAAAAGCCTCAGTGG CGTGATCTAGTGAAGGAGGCAGAGAAAGTATATTACAAAAGGGAGACAGAAGAAAAGGGAACAAAGAAAAGAGAGAGAGA AAGAGAGGAAAGGGAAAGACGTAATAAACGGCAAGAGAAGTTTGACTAAGATCTTGGCTGCAGTGGTTGAAGGGA AAAGCAATACGGAAAGAGAGAGAGATTTTAGGAAAATTAGGTCAGGCCCTAGACAGTCAGGGAACCTGGGCAATAGGACC CCACTCGACAAGGACCAATGTGCATATTGTAAAGAAAAAGGACACTGGGCAAGGAACTGCCCCAAGAAGGGAAACAAAGG ACTGAAGGTCTTAGCTCTGGAAGAAGATAAAGACTAGGGAAGACGGGGTTCGGACCCCCTCCCCGAGCCCAGGGTAACTT TGAAGGTGGAGGGCAACCAGTTGGGTTCCTGGTTGATACCGGAGCGAAACATTCAGTGCTACTACAGCCATTAGGAAAA CTAAAAGATAAAAAATCCTGGGTGATGGGTGCCACAGGGCAACAACAGTATCCATGGACTACCCGAAGAACAGTTGACTT GGGAGTGGGACGGGTAACCCACTCGTTTCTGGTCATACCTGAGTGCCCAGCACCCCTCTTAGGTAGAGACTTATTGACCA CCAAGCCTGGGCAGAAACCGCAGGGATGGGTTTGGCAAAGCAAGTTCCCCCACAAGTTATTCAACTGAAGGCCAGTGCCA CACCAGTGTCAGTCAGACAGTACCCCTTGAGTAAAGAAGCTCAAGAAGGAATTCGGCCGCATGTCCAAAGATTAATCCAA CAGGGCATCCTAGTTCCTGTCCAATCTCCCTGGAATACTCCCCTGCTACCGGTTAGAAAGCCTGGGACTAATGACTATCG ACCAGTACAGGACTTGAGAGAGGTCAATAAACGGGTGCAGGATATACACCCAACAGTCCCGAACCCTTATAAACCTCTTGT GTGCTCTCCCACCCCAACGGAGCTGGTATACAGTATTGGACTTAAAGGATGCCTTTTTCTGCCTGAGATTACACCCCACT AGCCAACCACTTTTTGCCTTCGAATGGAGAGATCCAGGTACGGGAAGAACCGGGCAGCTCACCTGGACCCGACTGCCCCA AGGGTTCAAGAACTCCCCGACCATCTTTGACGAAGCCCTACACAGAGACCTGGCCAACTTCAGGATCCAACACCCTCAGG TGACCCTCCTCCAGTACGTGGATGACCTGCTTCTGGCGGGAGCCACCAAACAGGACTGCTTAGAAGGCACGAAGGCACTA GTACAGTTTGCGGGACGGCAGCGATGGCTGACGGAGGCACGGAAGAAAACTGTAGTCCAGATACCGGCCCCAACCACAG AGTGGTTGGCCCGTATGCCTGAAGGCTATCGCAGCTGTGGCCATACTGGTCAAGGACGCTGACAAATTGACTTTGGGACA TGACCCACTATCAAAGCCTGCTTCTCACAGAGAGGGTCACGTTCGCTCCACCAGCCGCTCTCAACCCTGCCACTCTTCTG AGACATACCGCTGACTGGAGAAGTGTTAACCTGGTTCACTGACGGAAGCAGCTATGTGGTGGAAGGTAAGAGGATGGCTG AACACCCAAAGCCCCAGAACCCGGACGACAGTACACCCTAGAAGACTGGCAAGAGATAAAAAAGATAGACCAGTTCTCTG GACCCCCCCATTGGTAGAAATTGCTTCCGTACATAGTGCTGACGTGCTGCTTTCCCAGCCTTTGTTCTCTAGGCTCAAG GCACTTGAGTGGGTGAGACAACGAGCGTGGAGGCAACTCCGGGAGGCCTACTCAGGAGGAGGAGACTTGCAGATCC

LamA

TGGTTCACTGACGGAAGCAGCTATGTGGTGGAAGGTAAGAGGATGGCTGGGGCGGCGGTGGTGGACGGGACCCGCACGAT CTGGGCCAGCAGCCTGCCGGAAGGAACTTCAGCACAAAAGGCTGAGCTCATGGCCCTCACGCAAGCTTTGCGGCTGGCCG AAGGGAAATCCATAAACATTTATACAGACAGCAGGTATGCCTTTGCGACTGCACACGTACACGGGGCCATCTATAAGCAA AAGGGGTTGCTTACCTCAGCAGGGAGGGAATAAAAGAACAAAGAACCATACACCCATATACACCCAGAGGGAACCAGATGCTAAACAATCTGCC
AAAAAGGCTAGCTATTATACACTGTCCTGGACACATCAGAAGCCAAAGATCCCATATACCAGAGGGAACCAGATGGCTGACC
GGGTTGCCAAGCAGCCAGCCAGGTGTTAACCTTCTGCCTATGATAGAAACACCCCAAAGCCCCAGAACCCGGACGACAG TACACCCTAGAAGACTGGCAAGAGATAAAAAAGATAGACCAGTTCTCTGAGACTCCGGAAGGGACCTGCTATACCTCAGA
TGGGAAGGAAATCCTGCCCCACAAAGAAGGGTTAGAATATGTCCAACAGATACATCGTCTAACCCACCTAGGAACTAAAC
ACCTGCAGCAGTTGGTCAGAACATCTCCTTATCATGTTCTGAGGCTACCAGGAGTGGCTGATTCGGTGGTCAAACACTGT
GTGCCCTGCCAGCTGGTTAATCCTAATCCTTCCAGGAATACCTCCAGGAAGAAGAGACTAAAGGCGCACCCAGGCGCTCA CTGGGAAGTGGACTTCACTGAGGTAAAGCCGGCTAAATACGGAAACAAATATCTATTGGTTTTTGTAGACACCTTTTCAG GATGGGTAGAGGCTTATCCTACTAAGAAAGAGACTTCAACCGTGGTGGCTAAGAAAATACTGGAGGAAATTTTCCAAGA ACATAGTGCTGACGTGCTTTCCCAGCCTTTGTTCTCTAGGCTCAAGGCACTTGAGTGGGTGAGACAACGAGCGTGGA GGCAACTCCGGGAGGCCTACTCAGGAGGAGGAGACTTGCAGATCCCACATCGTTTCCAAGTGGGAGATTCAGTCTACGTT AGACGCCACCGTGCAGGAAACCTCGAGACTCGGTGGAAGGGCCCTTATCACGTACTTTTGACCACACCAACGGCTGTGAA AGTCGAAGGAATCTCCACCTGGATCCATGCATCCCACGTTAAGCCGGCGCCACCTCCCGATTCGGGGTGGAAAGCCGAAA AGACTGAAAATCCCCTTAAGCTTCGCCTCCATCGCGTGGTTCCTTACTCTGTCAATAACTCCTCAAGTTAATGGTAAACG ACAGCACTCAAGGGGAGCTCCCTTGGGGACCTGGTGGCCTGAATTATATGTCTGCCTTCGATCAGTAATCCCTGGTCTC AATGACCAGGCCACACCCCCGGATGTACTCCGTGCTTACGGGTTTTACGTTTGCCCAGGACCCCCAAATAATGAAGAATA TTGTGGAAATCCTCAGGATTTCTTTTGCAAGCAATGGAGCTGCGTAACTTCTAATGATGGGAAATTGGAAATGGCCAGTCT CTCAGCAAGACAGAGTAAGTTACTCTTTTGTTAACAATCCTACCAGTTATAATCAATTTAATTATGGCCATGGGAGATGG AAAGATTGGCAACAGCGGGTACAAAAAGATGTACGAAATAAGCAAATAAGCTGTCATTCGTTAGACCTAGATTACTTAAA AATAAGTTTCACTGAAAAAGGAAAACAAGAAAATATTCAAAAGTGGGTAAATGGTATGTCTTGGGGAATAGTGTACTATG GAGGCTCTGGGAGAAAGAAAGGATCTGTTCTGACTATTCGCCTCAGAATAGAAACTCAGATGGAACCTCCGGTTGCTATA GGACCAAATAAGGGTTTGGCCGAACAAGGACCTCCAATCCAAGAACAGAGCCATCTCCTAACCCCTCTGATTACAATAC AACCTCTGGATCAGTCCCCACTGAGCCTAACATCACTATTAAAACAGGGGCGAAACTTTTTAACCTCATCCAGGGAGCTT TTCAAGCTCTTAACTCCACGACTCCAGAGGCTACCTCTTCTTGTTGGCTTTGCTTAGCTTCGGGCCCACCTTACTATGAG GGAATGCCTAGAGGAGGGAAATTCAATGTGACAAAGGAACATAGAGACCAATGTACATGGGGATCCCAAAATAAGCTTAC CCTTACTGAGGTTTCTGGAAAAGGCACCTGCATAGGGATGGTTCCCCCATCCCACCACACCCTTTGTAACCACACTGAAGCCTTTAATCCGAACCTCTGAGAGTCAGTATCTGGTACCTGGTTATGACAGGTGGTGGGCATGTAATACTGGATTAACCCCT TGTGTTTCCACCTTGGTTTTCAACCAAACTAAAGACTTTTGCGTTATGGTCCAAATTGTCCCCCGGGTGTACTACTATCCCCGAAAAAGCAGTCCTTGATGAATATGACTATAGATATAATCGGCCAAAAAGAGAGCCCATATCCCTGACACTAGCTGTAA TGAAGGAGGAATGCTGTTTTTATGTGGATCATTCAGGGGCCCATCAGAGACTCCATGAACAAGCTTAGAGAAAGGTTGGAG CGCTAGCTCTACCAGTTCTAAGATTAGAACTATTAACAAGAGAAGAAGTGGGGGAATGAAAGGATGAAAATGCAACCTGAC TCTCCCAGAACCCAGGAAGTTAATAAGAAGCTCTAAATGCCCTCGAATTCCAGACCCTGTTCCCTATAGGTAAAAGATCA GGAGAAGTCAATTGCCTAACGGACCCCAGTAAGATCGGGTGTACCACAAAATGTTGAAACACATATCTTGGTGACAACAT GAAGTTTTAAATTGACTGGTTTGTGATATTTTGAAATGATTGGTTTGTAAAGCGCGGGCTTTGTTGTGAACCCCATAAAA GCTGTCCCGACTCCACACTCGGGGCCGCAGTCCTCTACCCCTGCGTGGTGTACGACTGTGGGCCCCAGCGCGCTTGGAAT TTTTTGCTAGTCTTACAGCACCTTTATTTTTTCCATTT

LamAl

TTCCTTGAGAGGCTCTTGGAAGCCTTCAGGCGGTACACCCCTTTTGATCCCACCTCAGAGGCCCAAAAAAGCCTCAGTGGC GAAGGTGGAGGGGCAACCAGTTGAGTTCCTGGTTGATACCGGAGCGAAACATTCAGTGCTACTACAGCCATTAGGAAAAC TAAAAGATAAAAAATCCTGGGTGATGGGTGCCACAGGGCAACAACAGTATCCATGGACTACCCCGAAGAACAGTTGACTTG GGAGTGGGACGGGTAACCCACTCGTTTCTGGTCATACCTGAGTGCCCAGCACCCCTCTTAGGTAGAGACTTATTGACCAA GATGGGAGCACAAATTTCTTTTGAACAAGGGAAACCAGAAGTGTCTGCAAATAACAAACCTATCACTGTGTTGACCCTCC AATTAGATGACGAATATCGACTATACTCTCCCCTAGTAAAGCCTGATCAAAATATACAATTCTGGTTGGAACAGTTTCCC CAAGCCTGGGCAGAAACCGCAGGGATGGGTTTGGCAAAGCAAGTTCCCCCACAAGTTATTCAACTGAAGGCCAGTGCCAC ACCAGTGTCAGTCAGACAGTACCCCTTGAGTAAAGAAGCTCAAGAAGGAATTCGGCCGCATGTCCAAAGATTAATCCAAC AGGGCATCCTAGTTCCTGTCCAATCTCCCTGGAATACTCCCCTGCTACCGGTTAGAAAGCCTGGGACTAATGACTATCGACCAGTACAGGACTTGAGAGAGGTCAATAAACGGGTGCAGGATATACACCCCAACAGTCCCGAACCCTTATAACCTCTTGTG TGCTCTCCCACCCCAACGGAGCTGGTATACAGTATTGGACTTAAAGGATGCCTTTTTCTGCCTGAGATTACACCCCACTA CAAACAAGTGAGAGAGTTTTTGGGGACAGCTGGATTTTGCAGACTGTGGATCCCGGGGGTTTGCGACCTTAGCAGCCCCAC TCTACCCACTAACCAAAGAAAAAGGGGAATTCTCCTGGGCTCCTGAGCACCAGAAGGCATTTGATGCTATCAAAAAGGCC CTGCTGAGCGCACCTGCTCTGGCCCTCCCTGACGTAACTAAACCCTTTACCCTTTATGTGGATGAGCGTAAGGGAGTAGC CCGGGGAGTTTTAACCCAAACTCTAGGACCATGGAGGAGACCTGTTGCCTACCTGTCAAAGAAGCTCGATCCTGTAGCCA CUGGGAGGCCGTATGCCTGAAGGCTATCGCAGCTGTGGCCATACTGGTCAAGGACGCTGACAAATTGACTTTGGGACAG GACATACCGCTGACTGGAGAAGTGTTAACCTGGTTCACTGACGGAAGCAGCTATGTAGTGGAAGGTAAGAGGATGGCTGG TACATCGTCTAACCCACCTAGGAACTAAACACCTGCAGCAGTTGGTCAGAACATCTCCTTATCATGTTCTGAGGCTACCA GGAGTGGCTGATTCGGTGGTCAAACACTGTGTGCCCTGCCAGCTGGGTAAAGCCCGGCTAAATACGGAAACAAATATCTAT TAAGTCAGGGACTGGCCAAGATATTGGGGATTGATTGGAAACTGCATTGTGCATACAGACCCCAAAGCTCAGGACAGGTA GAGAGGATGAATAGAACCATTAAAGAGACCCTTACTAAATTGACCGCGGAGACTGGCGTTAATGATTGGATAGCTCTCCT GCCCTTTGTGCTTTTTAGGGTTAGGAACACCCCTGGACAGTTTGGGCTGACCCCCTATAAATTACTCTACGGGGGACCCCCCCATTGGTAGAAATTGCTTCCGTACATAGTGCTGACGTGCTGCTTTCCCAGCCTTTGTTCTCTAGGCTCAAGGCACTT GAGTGGGTGAGACAACGAGCGTGGAGGCAACTCCGGGAGGCCTACTCAGGAGGAGAGACTTGCAGATCCCACATCGTTT
CCAAGTGGGAGATTCAGTTAGACGCCACCGTGCAGGAAACCTCGAGACTCGGTGGAAGGCCCTTATCACGTAC
TTTTGACCACACCAACGGCTGTGAAAGTCGAAGGAATCTCCACCTGGATCCATGCATCCCACGGTTAAGCCGGCGCACCT
CCCGATTCGGGGTGGAAAGCCGAAAAGACTGAAAATCCCCTTAAGCTTCGCCTCCATCGCGTGGTTCCTTACTCTGTCAA CCTTCGATCAGTAATCCCTGGTCTCAATGACCAGGCCACACCCCCGGATGTACTCCGTGCTTACGGGTTTTACGTTTGCC GATGGGAATTGGAAATGGCCAGTCTCAGCAAGACAGAGTAAGTTACTCTTTTGTTAACAATCCTACCAGTTATAATCA TCAGATGGAACCTCCGGTTGCTATAGGACCAAATAAGGGTTTGGCCGAACAAGACCTCCAATCCAAGAACAGAGCCAT CTCCTAACCCCTCTGATTACAATACAACCTCTGGATCAGTCCCCACTGAGCCTAACATCACTATTAAAACAGGGGCGAAA AAAGAAGGAGGATTATGTGTAGCCTTGAAGGAGGAATGCTGTTTTTATGTGGATCATTCAGGGGCCATCAGAGACTCCAT GAACAAGCTTAGAGAAAGGACTGGAGAGCGCCCGCGGGTCTCGAACAACCCAGACAGGTTGCTTGTTTCAATTAAAGAAC

28/28

SEQUENCE LISTING

<110> Mayo Foundation for Medical Education and Research 5 Federspiel, Mark J. <120> Methods to inhibit infectious agent transmission associated with transplant 10<130> 150.192WO1 <150> US 60/135,631 <151> 1999-05-24 15<160> 32 <170> FastSEQ for Windows Version 4.0 <210> 1 20<211> 47 <212> DNA <213> Artificial Sequence <220> 25<223> A primer <400> 1 47 gcgcatgcag atctgatgct taaacaggta gaaattttca ccgatgg 30<210> 2 <211> 45 <212> DNA <213> Artificial Sequence 35<220> <223> A primer <400> 2 gctgctgcgt cgacttaaac ttcaacttgg tagcctgtat cttcc 45 40 <210> 3

<211> 659

<212> PRT

<213> Porcine endogenous retrovirus

5<400> 3

Met His Pro Thr Leu Ser Arg Arg His Leu Pro Ile Arg Gly Gly Lys

1 5 10 15

Pro Lys Arg Leu Lys Ile Pro Leu Ser Phe Ala Ser Ile Ala Trp Phe 20 25 30

10Leu Thr Leu Ser Ile Thr Pro Gln Val Asn Gly Lys Arg Leu Val Asp
35 40 45

Ser Pro Asn Ser His Lys Pro Leu Ser Leu Thr Trp Leu Leu Thr Asp
50 55 60

Ser Gly Thr Gly Ile Asn Ile Asn Ser Thr Gln Gly Glu Ala Pro Leu 1565 70 75 80

Gly Thr Trp Trp Pro Glu Leu Tyr Val Cys Leu Arg Ser Val Ile Pro 85 90 95

Gly Leu Asn Asp Gln Ala Thr Pro Pro Asp Val Leu Arg Ala Tyr Gly
100 105 110

20Phe Tyr Val Cys Pro Gly Pro Pro Asn Asn Glu Glu Tyr Cys Gly Asn

Pro Gln Asp Phe Phe Cys Lys Gln Trp Ser Cys Val Thr Ser Asn Asp 130 135 140

Gly Asn Trp Lys Trp Pro Val Ser Gln Gln Asp Arg Val Ser Tyr Ser

Phe Val Asn Asn Pro Thr Ser Tyr Asn Gln Phe Asn Tyr Gly His Gly
165 170 175

Arg Trp Lys Asp Trp Gln Gln Arg Val Gln Lys Asp Val Arg Asn Lys

30Gln Ile Ser Cys His Ser Leu Asp Leu Asp Tyr Leu Lys Ile Ser Phe

Thr Glu Lys Gly Lys Gln Glu Asn Ile Gln Lys Trp Val Asn Gly Met

Thr Glu Lys Gly Lys Gln Glu Ash Tie Gin Lys Tip var Ash Gly Mee 210 215 220

Ser Trp Gly Ile Val Tyr Tyr Arg Gly Ser Gly Arg Lys Lys Gly Ser

Val Leu Thr Ile Arg Leu Arg Ile Glu Thr Gln Met Glu Pro Pro Val
245 250 255

Ala Ile Gly Pro Asn Lys Gly Leu Ala Glu Gln Gly Pro Pro Ile Gln
260 265 270

40Glu Gln Arg Pro Ser Pro Asn Pro Ser Asp Tyr Asn Thr Thr Ser Gly

		275					280					285			
Ser	Val	Pro	Thr	Glu :	Pro 2	Asn	Ile	Thr	Ile	Lys	Thr	Gly	Ala	Lys	Leu
	290					295					300				
Phe	Asn	Leu	Ile	Gln	Gly .	Ala	Phe	Gln	Ala	Leu	Asn	Ser	Thr	Thr	Pro
5305					310					315					320
Glu	Ala	Thr	Ser	Ser	Cys	Trp	Leu	Cys	Leu	Ala	Ser	Gly	Pro	Pro	Tyr
				325					330					335	
Tyr	Glu	Gly	Met	Ala	Arg	Gly	Gly	Lys	Phe	Asn	Val	Thr	Lys	Glu	His
_			340					345					350		
10Arg	Asp	Gln	Cys	Thr	Trp	Gly	Ser	Gln	Asn	Lys	Leu	Thr	Leu	Thr	Glu
		355					360					365			
Val	Ser	Gly	Lys	Gly	Thr	Cys	Ile	Gly	Met	Val	Pro	Pro	Ser	His	Gln
	370					375					380				
His	Le u	Cys	Asn	His	Thr	Glu	Ala	Phe	Asn	Arg	Thr	Ser	Glu	Ser	Gln
15385					390					395					400
Tyr	Leu	Val	Pro	Gly	Tyr	Asp	Arg	Trp	Trp	Ala	Cys	Asn	Thr	Gly	Leu
				405					410					415	
Thr	Pro	Cys	Val	Ser	Thr	Leu	Val	Phe	Asn	Gln	Thr	Lys	Asp	Phe	Cys
			420					425					430		
20Val	Met	Val	Gln	Ile	Val	Pro	Arg	Val	Tyr	Tyr	Tyr	Pro	Glu	Lys	Ala
		435					440					445			
Val	Leu	Asp	Glu	Tyr	Asp	Tyr	Arg	Tyr	Asn	Arg	Pro	Lys	Arg	Glu	Pro
	450)				455					460				
Ile	Ser	Leu	Thr	Leu	Ala	Val	Met	Leu	Gly	Leu	Gly	Val	Ala	Ala	Gly
25465					470					475					480
Val	. Gly	/ Thr	Gly	Thr	Ala	Ala	Leu	Ile	Thr	Gly	Pro	Gln	Gln	Leu	Glu
				485					490					495	_
Lys	Gly	/ Lev	ser	Asn	Leu	His	Arg	Ile	Val	Thr	Glu	Asn			Ala
			500					505					510		_
30Leu	ı Glı	ı Lys	s Ser	· Val	Ser	Asn	Leu	Glu	Glu	ı Ser	Leu			Leu	Ser
		515					520					525		_	
Glu	. Va	l Val	l Leu	ı Gln	Asn	Arg	Arg	Gly	Let	ı Asp			Phe	Leu	Lys
	530					535					540			_	
Gl	ı Gl	y Gly	y Lei	ı Cys	Val	Ala	Leu	Lys	Gli			: Cys	Phe	Tyr	Val
3554					550					555			_	~ 3	560
As	p Hi	s Se	r Gly	y Ala		Arg	Asp	Ser			n Lys	s Leu	ı Arg		
				565					570 		- س	~7	- C	575	
Le	u Gl	u Ly			Arg	Glu	ı Lys			r Th:	r Gli	n GIŞ			Glu
			58					585				_	590		
40G1	y Tr	p Ph	e Ası	n Arg	, Ser	Pro	Tr	. Lei	ı Al	a Th	r Lei	л гел	ı ser	Ala	Leu

4

605 595 600 Thr Gly Pro Leu Ile Val Leu Leu Leu Leu Leu Thr Val Gly Pro Cys 620 615 610 Ile Ile Asn Lys Leu Ile Ala Phe Ile Arg Glu Arg Ile Ser Ala Val 635 630 Gln Ile Met Val Leu Arg Gln Gln Tyr Gln Ser Pro Ser Ser Arg Glu 645 650 655 Ala Gly Arg 10 <210> 4 <211> 660 <212> PRT <213> Porcine endogenous retrovirus 15 <400> 4 Met His Pro Thr Leu Ser Arg Arg His Leu Pro Ile Arg Gly Gly Lys Pro Lys Arg Leu Lys Ile Pro Leu Ser Phe Ala Ser Ile Ala Trp Phe Leu Thr Leu Ser Ile Thr Pro Gln Val Asn Gly Lys Arg Leu Val Asp 45 35 40 Ser Pro Asn Ser His Lys Pro Leu Ser Leu Thr Trp Leu Leu Thr Asp 55 25Ser Gly Thr Gly Ile Asn Ile Asn Ser Thr Gln Gly Glu Ala Pro Leu 75 70 Gly Thr Trp Trp Pro Glu Leu Tyr Val Cys Leu Arg Ser Val Ile Pro 90 Gly Leu Asn Asp Gln Ala Thr Pro Pro Asp Val Leu Arg Ala Tyr Gly 105 100 30 Phe Tyr Val Cys Pro Gly Pro Pro Asn Asn Glu Glu Tyr Cys Gly Asn 120 Pro Gln Asp Phe Phe Cys Lys Gln Trp Ser Cys Ile Thr Ser Asn Asp 130 135 35Gly Asn Trp Lys Trp Pro Val Ser Gln Gln Asp Arg Val Ser Tyr Ser 150 155 Phe Val Asn Asn Pro Thr Ser Tyr Asn Gln Phe Asn Tyr Gly His Gly 170 Arg Trp Lys Asp Trp Gln Gln Arg Val Gln Lys Asp Val Arg Asn Lys 190 180 185

Gln	Ile	Ser	Cys	His	Ser	Leu	Asp	Leu	Asp	Tyr	Leu	Lys	Ile	Ser	Phe
		195					200					205			
Thr	Glu	Lys	Gly	Lys	Gln	Glu	Asn	Ile	Gln	Lys	Trp	Val	Asn	Gly	Ile
	210					215					220				
5Ser	Trp	Gly	Ile	Val	Tyr	Tyr	Gly	Gly	Ser	Gly	Arg	Lys	Lys	Gly	Ser
225					230					235					240
Val	Leu	Thr	Ile	Arg	Leu	Arg	Ile	Glu	Thr	Gln	Met	Glu	Pro	Pro	Val
				245					250					255	
Ala	Ile	Gly	Pro	Asn	Lys	Gly	Leu	Ala	Glu	Gln	Gly	Pro	Pro	Ile	Gln
10			260					265					270		
Glu	Gln	Arg	Pro	Ser	Pro	Asn	Pro	Ser	Asp	Tyr	Asn	Thr	Thr	Ser	Gly
		275					280					285			
Ser	Val	Pro	Thr	Glu	Pro	Asn	Ile	Thr	Ile	Lys	Thr	Gly	Ala	Lys	Leu
	290					295					300				
15Phe	Ser	Leu	Ile	Gln	Gly	Ala	Phe	Gln	Ala	Leu	Asn	Ser	Thr	Thr	Pro
305					310					315					320
Glu	Ala	Thr	Ser	Ser	Cys	Trp	Leu	Cys	Leu	Ala	Ser	Gly	Pro	Pro	Tyr
				325					330					335	
Tyr	Glu	Gly	Met	Ala	Arg	Gly	Gly	Lys	Phe	Asn	Val	Thr	Lys	Glu	His
20			340					345					350		
Arg	Asp	Gln	Cys	Thr	Trp	Gly	Ser	Gln	Asn	Lys	Leu	Thr	Leu	Thr	Glu
		355					360					365			
Val	Ser	Gly	Lys	Gly	Thr	Cys	Ile	Gly	Met	Val	Pro	Pro	Ser	His	Glr
	370					375					380				
25His	Leu	Cys	Asn	His	Thr	Glu	Ala	Phe	Asn	Arg	Thr	Ser	Glu	Ser	Glr
385					390					395					400
Tyr	Leu	Val	Pro	Gly	Tyr	Asp	Arg	Trp	Trp	Ala	Cys	Asn	Thr	Gly	Let
				405					410					415	
Thr	Pro	Cys	Val	Ser	Thr	Leu	Val	Phe	Asn	Gln	Thr	Lys	Asp	Phe	Cys
30			420					425					430		
Val	Met	Val	Gln	Ile	Val	Pro	Arg	Val	Tyr	Tyr	Tyr	Pro	Glu	Lys	Ala
		435					440					445			
Val	Leu	Asp	Glu	Tyr	Asp	Tyr	Arg	Tyr	Asn	Arg	Pro	Lys	Arg	Glu	Pro
	450					455					460				
35Ile	Ser	Leu	Thr	Leu	Ala	Val	Met	Leu	Gly	Leu	Gly	Val	Ala	Ala	Gly
465					470					475					480
Val	Gly	Thr	Gly	Thr	Ala	Ala	Leu	Ile	Thr	Gly	Pro	Gln	Gln	Leu	Gli
				485					490					495	
Lys	Gly	Leu	Ser	Asn	Leu	His	Arg	Ile	Val	Thr	Glu	Asp	Leu	Gln	Ala
40			500					505					510		

Leu Glu Lys Ser Val Ser Asn Leu Glu Glu Ser Leu Thr Ser Leu Ser 515 520 Glu Val Val Leu Gln Asn Arg Arg Gly Leu Asp Leu Leu Phe Leu Lys 535 5Glu Gly Gly Leu Cys Val Ala Leu Lys Glu Glu Cys Cys Phe Tyr Val 550 555 Asp His Ser Gly Ala Ile Arg Asp Ser Met Ser Lys Leu Arg Glu Arg 570 565 Leu Glu Arg Arg Arg Glu Arg Glu Ala Asp Gln Gly Trp Phe Glu 590 580 585 10 Gly Trp Phe Asn Arg Ser Pro Trp Met Thr Thr Leu Leu Ser Ala Leu 600 Thr Gly Pro Leu Val Val Leu Leu Leu Leu Leu Thr Val Gly Pro Cys 620 615 15Leu Ile Asn Arg Phe Val Ala Phe Val Arg Glu Arg Val Ser Ala Val 630 635 Gln Ile Met Val Leu Arg Gln Gln Tyr Gln Gly Leu Leu Ser Gln Gly 650 645 Glu Thr Asp Leu 20 660 <210> 5 <211> 638 <212> PRT 25<213> Porcine endogenous retrovirus <400> 5 Met His Pro Thr Leu Asn Arg Arg His Leu Pro Ile Arg Gly Gly Lys 10 30Pro Lys Arg Leu Lys Ile Pro Leu Ser Phe Ala Ser Ile Ala Trp Phe 25 20 Leu Thr Leu Ser Ile Thr Ser Gln Thr Asn Gly Met Arg Ile Gly Asp 40 Ser Leu Asn Ser His Lys Pro Leu Ser Leu Thr Trp Leu Ile Thr Asp 55 50 Ser Gly Thr Gly Ile Asn Ile Asn Asn Thr Gln Gly Glu Ala Pro Leu 70 75 Gly Thr Trp Trp Pro Asp Leu Tyr Val Cys Leu Arg Ser Val Ile Pro 40Ser Leu Thr Ser Pro Pro Asp Ile Leu His Ala His Gly Phe Tyr Val

			100					105					110		
Cys	Pro	Gly	Pro	Pro .	Asn .	Asn	Gly	Lys	His	Cys	Gly	Asn	Pro	Arg	Asp
-		115					120					125			
Phe	Phe	Cys	Lys	Gln	Trp	Asn	Cys	Val	Thr	Ser	Asn	Asp	Gly	Tyr	Trp
5	130					135					140				
Lys	Trp	Pro	Thr	Ser	Gln	Gln	Asp	Arg	Val	Ser	Phe	Ser	Tyr	Val	Asn
145					150					155					160
Thr	Tyr	Thr	Ser	Ser	Gly	Gln	Phe	Asn	Tyr	Leu	Thr	Trp	Ile	Arg	Thr
				165					170					175	
10Gly	Ser	Pro	Lys	Cys	Ser	Pro	Ser	Asp	Leu	Asp	Tyr	Leu	Lys	Ile	Ser
			180					185					190		
Phe	Thr	Glu	Lys	Gly	Lys	Gln	Glu	Asn	Ile	Leu	Lys	Trp	Val	Asn	Gly
		195					200					205			_ •
Met	Ser	Trp	Gly	Met	Val	Tyr	Tyr	Gly	Gly	Ser	Gly	Lys	Gln	Pro	Gly
15	210					215					220		_	_	
Ser	Ile	Leu	Thr	Ile	Arg	Leu	Lys	Ile	Asn		Leu	Glu	Pro	Pro	
225					230					235			_		240
Ala	Ile	Gly	Pro	Asn	Thr	Val	Leu	Thr		Gln	Arg	Pro	Pro		GIN
				245					250			3	D	255	Cl.,
20Gly	Pro	Gly			Ser	Asn	Ile		Ser	GIY	ser	Asp		1111	GIU
			260		_		~ 1	265	T	T	Dho	602	270	Tle	Gln
Ser	Ser			Thr	Lys	Met		Ala	ьуѕ	ьeu	PHE	285		110	0111
	_	275			~	3	280		mb ~	Dro	Clu			Ser	Ser
_			Gln	Ala	Leu		ser	THE	1111	PIO	300			501	001
25	290		0	Leu	ח ד ת	295	Cly	Pro	Pro	ጥኒታ			Glv	Met	Ala
) ьеч	Cys	ьец	310	261	GIY	FIO	110	315		0	,		320
305		- 611		: Phe		Val	Thr	Lvs	Glu			Asp	Gln	Cys	Thr
Arg	ALC	у Сту	, пле	325		var	****	270			3			335	
30Trp	. Gls	, Ser	- Glr			Leu	Thr	Leu	Thr	Glu	Val	Ser	Gly	Lys	Gly
30111	, 613	, 501	340					345					350		
Thr	- Cvs	s Ile		, Lys	Val	Pro	Pro	Ser	His	Gln	His	Leu	. Cys	Asn	His
	7-	355		-2-			360					365			
Thr	: Gli			a Asn	Gln	Thr	Ser	Glu	Ser	Gln	туг	Let	ı Val	Pro	Gly
35	37					375					380				
	r Asj	p Arg	g Trp	Trp	Ala	Cys	Asn	Thr	Gly	Leu	Thr	Pro	Cys	val	Ser
385					390					395					400
Thi	r Le	u Vai	l Phe	e Asn	Gln	Thr	Lys	Asp	Phe	Cys	: Ile	Met	. Val	Gln	lle
				405					410					415	
40Va	1 Dr	o Ar	or Va	TVI	Tvr	Tvr	Pro	Glu	ı Lys	: Ala	a Ile	e Lev	ı Ası	Glu	Tyr

			420					425					430			
Agn	TUr	Ara		His	Ara	Gln	Lvs		Glu	Pro	Tle	Ser		Thr	Leu	
изъ	171	435		1120	9	0111	440	5	014			445				
Δla	Val		Leu	Glv	Leu	Glv		Ala	Ala	Glv	Val		Thr	Gly	Thr	
5	450			1		455				1	460	1				
		Leu	Val	Thr	Glv		Gln	Gln	Leu	Glu		Glv	Leu	Ser	Asn	
465					470					475					480	
	His	Ara	Ile	Val		Glu	Asp	Leu	Gln		Leu	Glu	Lys	Ser	Val	
				485			•		490				•	495		
10Ser	Asn	Leu	Glu	Glu	Ser	Leu	Thr	Ser	Leu	Ser	Glu	Val	Val	Leu	Gln	
			500					505					510			
Asn	Arg	Arg	Gly	Leu	Asp	Leu	Leu	Phe	Leu	Lys	Glu	Gly	Gly	Leu	Cys	
		515					520					525				
Val	Ala	Leu	Lys	Glu	Glu	Cys	Cys	Phe	Tyr	Val	Asp	His	Ser	Gly	Ala	
15	530					535					540					
Ile	Arg	Asp	Ser	Met	Asn	Lys	Leu	Arg	Glu	Arg	Leu	Glu	Lys	Arg	Arg	
545					550					5 5 5					560	
Arg	Glu	Lys	Glu	Thr	Thr	Gln	Gly	Trp	Phe	Glu	Gly	Trp	Phe	Asn	Arg	
				565					570					575		
20Ser	Leu	Trp	Leu	Ala	Thr	Leu	Leu	Ser	Ala	Leu	Thr	Gly	Pro	Leu	Ile	
			580					585					590			
Va1	Leu	Leu	Leu	Leu	Leu	Thr	Val	Gly	Pro	Cys	Ile	Ile	Asn	Lys	Leu	
		595					600					605				
Ile	Ala	Phe	Ile	Arg	Glu	Arg	Ile	Ser	Ala	Val	Gln	Ile	Met	Val	Leu	
25	610					615					620					
-	Gln	Gln	Tyr	Gln		Pro	Ser	Ser	Arg		Ala	Gly	Arg			
625					630					635						
	0 > 6	0.4														
30<21	1> / 2> D															
			ne e	ndog	enou	c re	trov:	irue								
421	3 > F	OICI.	ne c	naog	enou.	s re	CIOV.	1143								
<40	0> 6															
		aaa	t.gaa.	aato	ca a	ccta	actc	t cc	caga	accc	agga	aaqt	taa	taaqa	aagctc	60
															tgctgt	120
	_		_												cctga	180
			_	_		_									cttctg	240
_															agtaag	300
															ccaccc	360

```
cgaaacatgc gcaaatgtgt aactctaaaa caatttaaat taattggtcc acgaagcgcg
                                                                         420
 ggctctcgaa gttttaaatt gactggtttg tgatattttg aaatgattgg tttgtaaagc
                                                                         480
 gcgggctttg ttgtgaaccc cataaaagct gtcccgactc cacactcggg gccgcagtcc
                                                                         540
 tctacccctg cgtggtgtac gactgtgggc cccagcgcgc ttggaataaa aatcctcttg
                                                                         600
                                                                         660
Sctqtttqcat caaqaccgct tctcgtgagt gattaagggg agtcgccttt tccgagcctg
                                                                         704
 gaggttcttt ttgctagtct tacatttggg ggctcgtccg ggat
 <210> 7
 <211> 633
10<212> DNA
  <213> Porcine endogenous retrovirus
  <400> 7
  aatgaaagga tgaaaataca acctaagcta atgagaagct taaaattgtt ctgaattcca
                                                                          60
                                                                         120
15qagtttgttc cttataggta aaagattagg ttttttgctg ttttaaaaata tgcggaagta
  aaataggccc tgagtacatg tctctaggca tgaaacttct tgaaactatt tgagataaca
                                                                          180
                                                                          240
  agaaaaggga gtttctaact gcttgtttag cttctgtaaa actggttgcg ccataaagat
                                                                          300
  qttqaaatqt tqatacacat atcttggtga caacatgtct cccccacccc gaaacatgcg
                                                                          360
  caaatgtgta actctaaaac aatttaaatt aattggtcca cgaagcgcgg gctctcgaag
20ttttaaattg actggtttgt gatattttga aatgattggt ttgtaaagcg cgggctttgt
                                                                          420
  tgtgaacccc ataaaagctg tcccgactcc acactcgggg ccgcagtcct ctacccctgc
                                                                          480
  gtggtgtacg actgtgggcc ccagcgcgct tggaataaaa atcctcttgc tgtttgcatc
                                                                          540
                                                                          600
  aaqaccqctt ctcgtgagtg attaagggga gtcgcctttt ccgagcctgg aggttctttt
                                                                          633
  tgctggtctt acatttgggg gctcgtccgg gat
25
  <210> 8
  <211> 20
  <212> DNA
  <213> Porcine endogenous retrovirus
30
  <400> 8
                                                                           20
  tggaaagatt ggcaacagcg
  <210> 9
35<211> 20
  <212> DNA
  <213> Porcine endogenous retrovirus
  <400> 9
                                                                           20
 40agtgatgtta ggctcagtgg
```

```
<210> 10
 <211> 20
 <212> DNA
 <213> Porcine endogenous retrovirus
 <400> 10
                                                                            20
 ttctcctttg tcaattccgg
  <210> 11
10<211> 20
  <212> DNA
  <213> Porcine endogenous retrovirus
  <400> 11
                                                                            20
15tactttatcg ggtcccactg
  <210> 12
  <211> 20
  <212> DNA
20<213> Porcine endogenous retrovirus
  <400> 12
                                                                            20
  ctgacctgga ttagaactgg
25<210> 13
  <211> 20
  <212> DNA
  <213> Porcine endogenous retrovirus
 30<400> 13
                                                                            20
   atgttagagg atggtcctgg
   <210> 14
   <211> 22
 35<212> DNA
   <213> Porcine endogenous retrovirus
   <400> 14
                                                                             22
   acctcgagac tcggtggaag gg
 40
```

<210> 15	
<211> 24	
<212> DNA	
<213> Porcine endogenous retrovirus	
5	
<400> 15	
ctgggttctg ggagggttag gttg	24
<210> 16	
10<211> 24	
<212> DNA	
<213> Porcine endogenous retrovirus	
<400> 16	
15acgtactgga ggagggtcac ctga	24
<210> 17	
<211> 24	
<212> DNA	
20<213> Porcine endogenous retrovirus	
<400> 17	
gtcccgaacc cttataacct cttg	24
25<210> 18	
<211> 1980	
<212> DNA	
<213> Porcine endogenous retrovirus	
20.400. 10	
30<400> 18 atgcatccca cgttaagccg gcgccacctc ccgattcggg gtggaaagcc gaaaagactg	60
aaaatcccct taagcttogc ctccatcgcg tggttcctta ctctgtcaat aactcctcaa	120
gttaatggta aacgeettgt ggacageeeg aacteecata aaccettate teteacetgg	180
ttacttactg actooggtac aggtattaat attaacagca ctcaagggga ggctcccttg	240
35gggacctggt ggcctgaatt atatgtctgc cttcgatcag taatccctgg tctcaatgac	300
cagging gardigate analyticage corregations takened garding control and cagging	360
aataatgaag aatattgtgg aaatcctcag gatttctttt gcaagcaatg gagctgcgta	420
actictaatg atgggaattg gaaatggcca gtctctcagc aagacagagt aagttactct	480
tttgttaaca atcctaccag ttataatcaa tttaattatg gccatgggag atggaaagat	540
40tggcaacagc gggtacaaaa agatgtacga aataagcaaa taagctgtca ttcgttagac	600
inclinated Additionary admid and a manufactor conditions	-

ctagattact	taaaaataag	tttcactgaa	aaaggaaaac	aagaaaatat	tcaaaagtgg	660
gtaaatggta	tgtcttgggg	aatagtgtac	tatagaggct	ctgggagaaa	gaaaggatct	720
gttctgacta	ttcgcctcag	aatagaaact	cagatggaac	ctccggttgc	tataggacca	780
aataagggtt	tggccgaaca	aggacctcca	atccaagaac	agaggccatc	tcctaacccc	840
5tctgattaca	atacaacctc	tggatcagtc	cccactgagc	ctaacatcac	tattaaaaca	900
ggggcgaaac	tttttaacct	catccaggga	gcttttcaag	ctcttaactc	cacgactcca	960
gaggctacct	cttcttgttg	gctttgctta	gcttcgggcc	caccttacta	tgagggaatg	1020
gctagaggag	ggaaattcaa	tgtgacaaag	gaacatagag	accaatgtac	atggggatcc	1080
caaaataagc	ttacccttac	tgaggtttct	ggaaaaggca	cctgcatagg	gatggttccc	1140
10ccatcccacc	aacacctttg	taaccacact	gaagccttta	atcgaacctc	tgagagtcag	1200
tatctggtac	ctggttatga	caggtggtgg	gcatgtaata	ctggattaac	cccttgtgtt	1260
tccaccttgg	ttttcaacca	aactaaagac	ttttgcgtta	tggtccaaat	tgtcccccgg	1320
gtgtactact	atcccgaaaa	agcagtcctt	gatgaatatg	actatagata	taatcggcca	1380
aaaagagagc	ccatatccct	gacactagct	gtaatgctcg	gattgggagt	ggctgcaggc	1440
15gtgggaacag	gaacggctgc	cctaatcaca	ggaccgcaac	agctggagaa	aggacttagt	1500
aacctacatc	gaattgtaac	ggaaaatctc	caagccctag	aaaaatctgt	cagtaacctg	1560
gaggaatccc	taacctcctt	atctgaagtg	gttctacaga	acagaagggg	gttagatctg	1620
ttatttctaa	aagaaggagg	attatgtgta	gccttaaagg	aggaatgctg	tttttatgtg	1680
gatcattcag	gggccatcag	agactccatg	aacaagctta	gagaaaggtt	ggagaagcgt	1740
20cgaagggaaa	aggaaactac	tcaagggtgg	tttgagggat	ggttcaacag	gtctccttgg	1800
ttggctaccc	tactttctgc	tttaacagga	cccttaatag	tectectect	gttactcaca	1860
gttgggccat	gtattattaa	caagttaatt	gccttcatta	gagaacgaat	aagtgcagtc	1920
cagatcatgg	tacttagaca	acagtaccaa	agcccgtcta	gcagagaagc	tggccgctag	1980

25<210> 19

<211> 7362

<212> DNA

<213> Porcine endogenous retrovirus

30<400> 19

tacttcttgg	ggaagaccct	gggctgctaa	ctgggtcttg	gctggtccta	gtgaaaggat	60
gaaaatgcaa	cctgactctc	ccagaaccca	ggaagttaat	aagaagctct	aaataatgaa	120
aggatgaaaa	tgcaacctga	ctctcccaga	acccaggaag	ttaataagaa	gctctaaatg	180
ccctcgaatt	ccagaccctg	ttccctatag	gtaaaagatc	atactttttg	ctgttttagg	240
35gcttgctttc	tgctctgtac	aaaactttgt	ggaaggggaa	aaacaggccc	ctgagtatgt	300
gcctctatgc	ttgaaacttc	ttgaaactgc	tcctaactgc	ttgtttggct	tctgtaaacc	360
tgcttgcata	agataaaaag	aggagaagtc	aattgcctaa	cggaccccag	taagatcggg	420
tgtaccacaa	aatgttgaaa	cacatatctt	ggtgacaaca	tgtctcccc	accccgaaac	480
atgcgcaaat	gtgtaactct	aaaacaattt	aaattaattg	gtccacgaag	cgcgggctct	540
40cgaagtttta	aattgactgg	tttgtgatat	tttgaaatga	ttggtttgta	aagcgcgggc	600

tttgttgtga	accccataaa	agctgtcccg	actccacact	cggggccgca	gtcctctacc	660
cctgcgtggt	gtacgactgt	gggccccagc	gcgcttggaa	taaaaatcct	cttgctgttt	720
gcatcaagac	cgcttctcgt	gagtgattaa	ggggagtcgc	cttttccgag	cctggaggtt	780
ctttttgcta	gtcttacatt	tgggggctcg	tccgggatct	gtcgcggcca	cccctaacac	840
5ccgagaaccg	acttggaggt	aaaaaggatc	ctctttttaa	cgtgtatgca	tgtaccggcc	900
ggcgtctctg	ttctgagtgt	ctgttttcag	tggtgcgcgc	tttcggtttg	cagctgtcct	960
ctcagaccgt	aaggactggg	ggactgtgat	cagcagacgt	gctaggagga	tcacaggctg	1020
ccaccctggg	ggacgccccg	ggaggtgggg	agagccaggg	acgcctggtg	gtctccttct	1080
gtcggtcaga	ggaccgagtt	ctgttgttga	agcgaaagct	tccccctccg	cggccgtccg	1140
10actcttttgc	ctgcttgtgg	aagacgcgga	cgggtcgcgt	gtgtctggat	ctgttggttt	1200
ctgttttgtg	tgtctttgtc	ttgtgcgtcc	ttgtctacag	ttttaatatg	ggacagacgg	1260
tgacgacccc	tcttagtttg	actctcgacc	attggactga	agttaaatcc	agggctcata	1320
atttgtcagt	tcaggttaag	aagggacctt	ggcagacttt	ctgtgtctct	gaatggccga	1380
cattcgatgt	tggatggcca	tcagagggga	cctttaattc	tgagattatc	ctggctgtta	1440
15aagcaattat	ttttcagact	ggacccggct	ctcatcccaa	tcaggagccc	tatatcctta	1500
cgtggcaaga	tttggcagag	gatecteege	catgggttaa	accttggctg	aataagccaa	1560
gaaagccagg	tccccgaatt	ctggctcttg	gagagaaaaa	caaacactcg	gctgaaaaag	1620
tcaagccctc	tcctcatatc	taccccgaga	ttgaggagcc	gccggcttgg	ccggaacccc	1680
aatctgttcc	cccaccccct	tatctggcac	agggtgctgc	gaggggaccc	tctgcccctc	1740
20ctggagctcc	ggcggtggag	ggacctgctg	cagggactcg	gagccggagg	ggcgccaccc	1800
cggagcggac	agacgagatc	gcgacattac	cgctgcgcac	gtacggccct	cccacaccgg	1860
ggggccaatt	gcagcccctc	cagtattggc	ccttttcttc	tgcagatctc	tataattgga	1920
aaactaacca	tcccctttc	tcggaggatc	cccaacgcct	cacggggttg	gtggagtccc	1980
ttatgttctc	tcaccagcct	acttgggatg	attgtcaaca	gctgctgcag	acactcttca	2040
25caaccgagga	gcgagagaga	attctgttag	aggctagaaa	aaatgttcct	ggggccgacg	2100
ggcgacccac	gcagttgcaa	aatgagattg	acatgggatt	tcccttgact	cgccccggtt	2160
gggactacaa	cacggctgaa	ggtagggaga	gcttgaaaat	ctatcgccag	gctctggtgg	2220
cgggtctccg	gggcgcctca	agacggccca	ctaatttggc	taaggtaaga	gaagtgatgc	2280
agggaccgaa	tgaaccccc	: tctgttttcc	ttgagaggct	cttggaagcc	ttcaggcggt	2340
30acaccccttt	tgatcccaco	: tcagaggccc	aaaaagcctc	agtggctttg	gcctttatag	2400
gacagtcago	cttggatatt	agaaagaago	: ttcagagact	ggaagggtta	caggaggctg	2460
agttacgtga	tctagtgaag	g gaggcagaga	aagtatatta	caaaagggag	acagaagaag	2520
aaagggaaca	a aagaaaagag	g agagaaagag	g aggaaaggga	ggaaagacgt	aataaacggc	2580
aagagaagaa	a tttgactaag	g atcttggctg	g cagtggttga	agggaaaagc	aatacggaaa	2640
35gagagagag	a ttttaggaaa	a attaggtcag	g gccctagaca	gtcagggaac	ctgggcaata	2700
ggaccccact	cgacaaggad	caatgtgcat	attgtaaaga	aaaaggacac	tgggcaagga	2760
actgcccca	a gaagggaaa	c aaaggactga	a aggtettage	tctgaaagaa	gataaagact	2820
agggaagac	g gggttcgga	c cocctccccq	g agcccagggt	aactttgaag	gtggagggc	2880
aaccagttg	a gttcctggt	t gataccggag	g cgaaacatto	agtgctacta	cagccattag	2940
40gaaaactaa	a agataaaaa	a teetgggtga	a tgggtgccad	agggcaacaa	cagtatccat	3000

PCT/US00/14296

ggactacccg	aagaacagtt	gacttgggag	tgggacgggt	aacccactcg	tttctggtca	3060
tacctgagtg	cccagcaccc	ctcttaggta	gagacttatt	gaccaagatg	ggagcacaaa	3120
tttcttttga	acaagggaaa	ccagaagtgt	ctgcaaataa	caaacctatc	actgtgttga	3180
ccctccaatt	agatgacgaa	tatcgactat	actctcccct	agtaaagcct	gatcaaaata	3240
5tacaattctg	gttggaacag	tttccccaag	cctgggcaga	aaccgcaggg	at gg gtttgg	3300
caaagcaagt	tccccacaa	gttattcaac	tgaaggccag	tgccacacca	gtgtcagtca	3360
gacagtaccc	cttgagtaaa	gaagctcaag	aaggaattcg	gccgcatgtc	caaagattaa	3420
tccaacaggg	catcctagtt	cctgtccaat	ctccctggaa	tactcccctg	ctaccggtta	3480
gaaagcctgg	gactaatgac	tatcgaccag	tacaggactt	gagagaggtc	aataaacggg	3540
10tgcaggatat	acacccaaca	gtcccgaacc	cttataacct	cttgtgtgct	ctcccacccc	3600
aacggagctg	gtatacagta	ttggacttaa	aggatgcctt	tttctgcctg	agattacacc	3660
ccactagcca	accacttttt	gccttcgaat	ggagagatcc	aggtacggga	agaaccgggc	3720
agctcacctg	gacccgactg	ccccaagggt	tcaagaactc	cccgaccatc	tttgacgaag	3780
ccctacacag	agacctggcc	aacttcagga	tccaacaccc	tcaggtgacc	ctcctccagt	3840
15acgtggatga	cctgcttctg	gcgggagcca	ccaaacagga	ctgcttagaa	ggcacgaagg	3900
cactactgct	ggaattgtct	gacctaggct	acagagcctc	tgctaagaag	gcccagattt	3960
gcaggagaga	ggtaacatac	ttggggtaca	gtttgcggga	cgggcagcga	tggctgacgg	4020
aggcacggaa	gaaaactgta	gtccagatac	cggccccaac	cacagccaaa	caagtgagag	4080
agtttttggg	gacagctgga	ttttgcagac	tgtggatccc	ggggtttgcg	accttagcag	4140
20ccccactcta	cccactaacc	aaagaaaaag	gggaattctc	ctgggctcct	gagcaccaga	4200
aggcatttga	tgctatcaaa	aaggccctgc	tgagcgcacc	tgctctggcc	ctccctgacg	4260
		tatgtggatg				4320
cccaaactct	aggaccatgg	aggagacctg	ttgcctacct	gtcaaagaag	ctcgatcctg	4380
		tgcctgaagg				4440
25acgctgacaa						4500
		gaccgatgga				4560
		gtcacgttcg				4620
		. gaaccagtga				4680
		: cttacagaca				4740
30tcactgacgg						4800
		gccagcagcc				4860
					aacatttata	4920
					aagcaaaggg	4980
					agcctattag	5040
35aagccttaca						5100
					gcagcccagg	5160
					cgacagtaca	5220
					ccggaaggga	5280
					gaatatgtcc	5340
40aacagatac	a togtotaaco	c cacctaggaa	a ctaaacacct	gcagcagttg	gtcagaacat	5400

	ctccttatca	tgttctgagg	ctaccaggag	tggctgattc	ggtggtcaaa	cactgtgtgc	5460
	cctgccagct	ggttaatgct	aatccttcca	gaatacctcc	aggaaagaga	ctaaggggaa	5520
	gccacccagg	cgctcactgg	gaagtggact	tcactgaggt	aaagccggct	aaatacggaa	5580
	acaaatatct	attggttttt	gtagacacct	tttcaggatg	ggtagaggct	tatcctacta	5640
ģ	Sagaaagagac	ttcaaccgtg	gtggctaaaa	aaatactgga	ggaaattttt	ccaagatttg	5700
	gaatacctaa	ggtaataggg	tcagacaatg	gtccagcttt	cgttgcccag	gtaagtcagg	5760
	gactggccaa	gatattgggg	attgattgga	aactgcattg	tgcatacaga	ccccaaagct	5820
	caggacaggt	agagaggatg	aatagaacca	ttaaagagac	ccttactaaa	ttgaccgcgg	5880
	agactggcgt	taatgattgg	atagctctcc	tgccctttgt	actttttagg	gttaggaaca	5940
1	Occcctggaca	gtttgggctg	accccctatg	aattactcta	cgggggaccc	ccccattgg	6000
	tagaaattgc	ttccgtacat	agtgctgacg	tgctgctttc	ccagcctttg	ttctctaggc	6060
	tcaaggcact	tgagtgggtg	agacaacgag	cgtggaggca	actccgggag	gcctactcag	6120
	gaggaggaga	cttgcagatc	ccacatcgtt	tccaagtggg	agattcagtc	tacgttagac	6180
	gccaccgtgc	aggaaacctc	gagactcggt	ggaagggccc	ttatcacgta	cttttgacca	6240
1	5caccaacggc	tgtgaaagtc	gaaggaatct	ccacctggat	ccatgcatcc	cacgttaagc	6300
	cggcgccacc	tcccgattcg	gggtggaaag	ccgaaaagac	tgaaaatccc	cttaagcttc	6 36 0
	gcctccatcg	cgtggttcct	tactctgtca	ataactcctc	aagttaatgg	taaacgcctt	6420
	gtggacagcc	cgaactccca	taaaccctta	tctctcacct	ggttacttac	tgactccggt	6480
	acaggtatta	atattaacag	cactcaaggg	gaggctccct	tggggacctg	gtggcctgaa	6540
2	Ottatatgt:t	gccttcgatc	agtaatccct	ggtctcaatg	accaggccac	accccccgat	6600
	gtactccgtg	cttacgggtt	ttacgtttgc	ccaggacccc	caaataatga	agaatattgt	6660
	ggaaatcctc	aggatttctt	ttgcaagcaa	tggagctgcg	taacttctaa	tgatgggaat	6720
	tggaaatggc	cagtctctca	gcaagacaga	gtaagttact	cttttgttaa	caatcctacc	6780
	agttataato	aatttaatta	tggccatggg	agatggaaag	attggcaaca	gcgggtacaa	6840
2	25aaagatgtac	gaaataagca	aataagctgt	cattcgttag	acctagatta	cttaaaaata	6900
	agtttcactg	aaaaaggaaa	acaagaaaat	attcaaaagt	gggtaaatgg	tatgtcttgg	6960
	ggaatagtgt	actatggagg	ctctgggaga	aagaaaggat	ctgttctgac	tattcgcctc	7020
	agaatagaaa	ctcagatgga	acctccggtt	gctataggac	caaataaggg	tttggccgaa	7080
	caaggaccto	caatccaaga	acagaggcca	tctcctaacc	cctctgatta	caatacaacc	7140
:	30tctggatcag	tccccactga	gcctaacatc	actattaaaa	caggggcgaa	actttttaac	7200
	ctcatccagg	gagettttea	agctcttaac	tccacgactc	cagaggctac	ctcttcttgt	7260
	tggctttgct	tagcttcggg	g cccaccttac	tatgagggaa	tggctagagg	agggaaattc	7320
	aatgtgacaa	a aggaacatag	g agaccaatgt	acatggggat	cc		7362

<210> 20

<211> 4402

<212> DNA

40<213> Porcine endogenous retrovirus

16

<400> 20 gatettgget geagtggttg aagggaaaag caatacggaa agagagagag attttaggaa 60 aattaggtca ggccctagac agtcagggaa cctgggcaat aggaccccac tcgacaagga 120 ccaatgtgca tattgtaaag aaaaaggaca ctgggcaagg aactgcccca agaagggaaa 180 Scaaaggactg aaggtettag etetggaaga agataaagae tagggaagae ggggttegga 240 cccctcccc gagcccaggg taactttgaa ggtggagggg caaccagttg agttcctggt 300 tgataccgga gcgaaacatt cagtgctact acagccatta ggaaaactaa aagataaaaa 360 atcctgggtg atgggtgcca cagggcaaca acagtatcca tggactaccc gaagaacagt 420 tgacttggga gtgggacggg taacccactc gtttctggtc atacctgagt gcccagcacc 480 10cctcttaggt agagacttat tgaccaagat gggagcacaa atttcttttg aacaagggaa 540 accagaagtg tctgcaaata acaaacctat cactgtgttg accctccaat tagatgacga 600 atategacta tactetecce tagtaaagee tgatcaaaat atacaattet ggttggaaca 660 gtttccccaa gcctgggcag aaaccgcagg gatgggtttg gcaaagcaag ttcccccaca 720 agttattcaa ctgaaggcca gtgccacacc agtgtcagtc agacagtacc ccttgagtaa 780 15agaageteaa gaaggaatte ggeegeatgt ecaaagatta atecaacagg geateetagt 840 teetgtecaa teteeetgga atacteeeet getaceggtt agaaageetg ggaetaatga 900 ctatcgacca gtacaggact tgagagaggt caataaacgg gtgcaggata tacacccaac 960 agtecegaac cettataace tettgtgtge teteceacee caacggaget ggtatacagt 1020 attggactta aaggatgeet ttttetgeet gagattacae eecactagee aaccaetttt 1080 20tgccttcgaa tggagagatc caggtacggg aagaaccggg cagctcacct ggacccgact 1140 geeccaaggg tteaagaact eeccgaceat etttgaegaa geectacaea gagaeetgge 1200 caacttcagg atccaacace ctcaggtgac cetectccag tacgtggatg acctgettet 1260 ggcgggagcc accaaacagg actgcttaga aggcacgaag gcactactgc tggaattgtc 1320 1380 tgacctaggc tacagagcct ctgctaagaa ggcccagatt tgcaggagag aggtaacata 25cttggggtac agtttgcggg acgggcagcg atggctgacg gaggcacgga agaaaactgt 1440 agtccagata ccggccccaa ccacagccaa acaagtgaga gagtttttgg ggacagctgg 1500 attttgcaga ctgtggatcc cggggtttgc gaccttagca gccccactct acccactaac 1560 caaagaaaaa ggggaattct cctgggctcc tgagcaccag aaggcatttg atgctatcaa 1620 1680 aaaggccctg ctgagcgcac ctgctctggc cctccctgac gtaactaaac cctttaccct 30ttatgtggat gagcgtaagg gagtagcccg gggagtttta acccaaactc taggaccatg 1740 gaggagacct gttgcctacc tgtcaaagaa gctcgatcct gtagccagtg gttggcccgt 1800 atgectgaag getategeag etgtggeeat aetggteaag gaegetgaea aattgaettt 1860 gggacagaat ataactgtaa tagcccccca tgcgttggag aacatcgttc ggcagccccc 1920 agaccgatgg atgaccaacg cccgcatgac ccactatcaa agcctgcttc tcacagagag 1980 35ggtcacgttc gctccaccag ccgctctcaa ccctgccact cttctgcctg aagagactga 2040 tgaaccagtg actcatgatt gccatcaact attgattgag gagactgggg tccgcaagga 2100 ccttacagac ataccgctga ctggagaagt gttaacctgg ttcactgacg gaagcagcta 2160 tgtggtggaa ggtaagagga tggctggggc ggcggtggtg gacgggaccc gcacgatctg 2220 ggccagcagc ctgccggaag gaacttcagc acaaaaggct gagctcatgg ccctcacgca 2280 40agctttgcgg ctggccgaag ggaaatccat aaacatttat acagacagca ggtatgcctt 2340

tgcgactgca	cacgtacacg	gggccatcta	taagcaaagg	gggttgctta	cctcagcagg	2400
gagggaaata	aagaacaaag	aggaaattct	aagcctatta	gaagccttac	atttgccaaa	2460
aaggctagct	attatacact	gtcctggaca	tcagaaagcc	aaagatccca	tatccagagg	2520
		ttgccaagca				2580
		cagaacccgg				2640
		tctctgagac				2700
		aagaagggtt				2760
		tgcagcagtt				2820
		cggtggtcaa				2880
10taatccttcc						2940
		taaagccggc				3000
		gggtagaggc				3060
		aggaaatttt				3120
		tcgttgccca				3180
15gattgattgg						3240
		cccttactaa				3300
		tgctttttag				3360
		acgggggacc				3420
		cccagccttt				3480
20gagacaacga						3540
		gagattcagt				3600
		cttatcacgt				3660
		a tccatgcatc				3720
		a ctgaaaatcc				3780
					ccgaactccc	3840
					aatattaaca	3900
					tgccttcgat	3960
					gcttacgggt	4020
					caggatttct	4080
30tttgcaage	a atggagetg	c gtaacttcta	atgatgggaa	a ttggaaatg	g ccagtctctc	4140
					t caatttaatt	4200
					a cgaaataagc	4260
					t gaaaaaggaa	4320
аасаадааа	a tattcaaaa	g tgggtaaatg	gtatgtctt	g gggaatagt	g tactatggag	4380
	g aaagaaagg		_			4402
2220003334	, <u>-</u>					

<210> 21

<211> 6076

<212> DNA

^{40&}lt;213> Porcine endogenous retrovirus

<400> 21						
	ggtgagaggc	tgtggcatag	ggaggcctcc	ctggaggagg	ttgaaaggat	60
		ccagaaccca				120
		tataggtaaa				180
5ctttctgctc						240
		actgctccta				300
		aagtcaattg				360
		atcttggtga				420
		aatttaaatt				480
10ttttaaattg	actggtttgt	gatattttga	aatgattggt	ttgtaaagcg	cgggctttgt	540
		tcccgactcc				600
		ccagcgcgct				660
		attaagggga				720
tgctagtctt	acatttgggg	gctcgtccgg	gatctgtcgc	ggccacccct	aacacccgag	780
15aaccgacttg						840
		ttcagtggtg				900
accgtaagga	ctgggggact	gtgatcagca	gacgtgctag	gaggatcaca	ggctgccacc	960
		tggggagagc				1020
		gttgaagcga				1080
20tttgcctgct	tgtggaagac	gcggacgggt	cgcgtgtgtc	tggatctgtt	ggtttctgtt	1140
ttgtgtgtct	ttgtcttgtg	cgtccttgtc	tacagtttta	atatgggaca	gacggtgacg	1200
		cgaccattgg				1260
tcagttcagg	ttaagaaggg	accttggcag	actttctgtg	tctctgaatg	gccgacattc	1320
gatgttggat	ggccatcaga	ggggaccttt	aattctgaga	ttatcctggc	tgttaaagca	1380
25attatttttc	agactggacc	cggctctcat	cccaatcagg	agccctatat	ccttacgtgg	1440
caagatttgg	cagaggatco	tccgccatgg	gttaaacctt	ggctgaataa	gccaagaaag	1500
ccaggtcccc	gaattetgge	tcttggagag	aaaaacaaac	actcggctga	aaaagtcaag	1560
ccctctcctc	atatctacco	: cgagattgag	gagccgccgg	cttggccgga	accccaatct	1620
gttcccccac	ccccttatct	ggcacagggt	gctgcgaggg	gaccctctgc	ccctcctgga	1680
30gctccggcgg	tggagggaco	tgctgcaggg	actcggagcc	ggaggggcgc	cacccggag	1740
cggacagacg	agatcgcgad	attaccgctg	cgcacgtacg	gccctcccac	accggggggc	1800
					ttggaaaact	1860
aaccatccc	ctttctcgg	a ggatccccaa	cgcctcacgg	ggttggtgga	gtcccttatg	1920
					cttcacaacc	1980
					cgacgggcga	2040
					: cggttgggac	2100
					ggtggcgggt	2160
					gatgcaggga	2220
					g gcggtacacc	2280
40ccttttgat	c ccacctcag	a ggcccaaaaa	a gcctcagtgg	ctttggcctt	tataggacag	2340

tcagccttgg	atattagaaa	gaagcttcag	agactggaag	ggttacagga	ggctgagtta	2400
cgtgatctag	tgaaggaggc	agagaaagta	tattacaaaa	gggagacaga	agaagaaagg	2460
gaacaaagaa	aagagagaga	aagagaggaa	agggaggaaa	gacgtaataa	acggcaagag	2520
aagaatttga	ctaagatctt	ggctgcagtg	gttgaaggga	aaagcaatac	ggaaagagag	2580
5agagatttta	ggaaaattag	gtcaggccct	agacagtcag	ggaacctggg	caataggacc	2640
ccactcgaca	aggaccaatg	tgcatattgt	aaagaaaaag	gacactgggc	aaggaactgc	2700
cccaagaagg	gaaacaaagg	actgaaggtc	ttagctctgg	aagaagataa	agactaggga	2760
agacggggtt	cggaccccct	ccccgagccc	agggtaactt	tgaaggtgga	ggggcaacca	2820
gttgggttcc	tggttgatac	cggagcgaaa	cattcagtgc	tactacagcc	attaggaaaa	2880
10ctaaaagata	aaaaatcctg	ggtgatgggt	gccacagggc	aacaacagta	tccatggact	2940
acccgaagaa	cagttgactt	gggagtggga	cgggtaaccc	actcgtttct	ggtcatacct	3000
gagtgcccag	cacccctctt	aggtagagac	ttattgacca	agatgggagc	acaaatttct	3060
tttgaacaag	ggaaaccaga	agtgtctgca	aataacaaac	ctatcactgt	gttgaccctc	3120
caattagatg	acgaatatcg	actatactct	cccctagtaa	agcctgatca	aaatatacaa	3180
15ttctggttgg	aacagtttcc	ccaagcctgg	gcagaaaccg	cagggatggg	tttggcaaag	3240
caagttcccc	cacaagttat	tcaactgaag	gccagtgcca	caccagtgtc	agtcagacag	3300
taccccttga	gtaaagaagc	tcaagaagga	attcggccgc	atgtccaaag	attaatccaa	3360
cagggcatcc	tagttcctgt	ccaatctccc	tggaatactc	ccctgctacc	ggttagaaag	3420
cctgggacta	atgactatcg	accagtacag	gacttgagag	aggtcaataa	acgggtgcag	3480
20gatatacacc	caacagtccc	gaacccttat	aacctcttgt	gtgctctccc	accccaacgg	3540
agctggtata	cagtattgga	cttaaaggat	gcctttttct	gcctgagatt	acaccccact	3600
agccaaccac	tttttgcctt	cgaatggaga	gatccaggta	cgggaagaac	cgggcagctc	3660
acctggaccc	gactgcccca	agggttcaag	aactccccga	ccatctttga	cgaagcccta	3720
cacagagacc	tggccaactt	caggatccaa	caccctcagg	tgaccctcct	ccagtacgtg	3780
25gatgacctgc	ttctggcggg	agccaccaaa	caggactgct	tagaaggcac	gaaggcacta	3840
ctgctggaat	tgtctgacct	aggctacaga	gcctctgcta	agaaggccca	gatttgcagg	3900
agagaggtaa	catacttggg	gtacagtttg	cgggacgggc	agcgatggct	gacggaggca	3960
cggaagaaaa	ctgtagtcca	gataccggcc	ccaaccacag	ccaaacaagt	gagagagttt	4020
ttggggacag	ctggattttg	cagactgtgg	atcccggggt	ttgcgacctt	agcagcccca	4080
30ctctacccac	taaccaaaga	aaaaggggaa	ttctcctggg	ctcctgagca	ccagaaggca	4140
tttgatgcta	tcaaaaaggc	cctgctgagc	gcacctgctc	tggccctccc	tgacgtaact	4200
aaacccttta	ccctttatgt	ggatgagcgt	aagggagtag	cccggggagt	tttaacccaa	4260
actctaggad	catggaggag	acctgttgcc	tacctgtcaa	agaagctcga	tcctgtagcc	4320
agtggttggd	ccgtatgcct	gaaggctato	gcagctgtgg	ccatactggt	caaggacgct	4380
35gacaaattga	ctttgggaca	gaatataact	gtaatagccc	accatgcgtt	ggagaacatc	4440
gttcggcag	ccccagaccg	atggatgaco	aacgcccgca	tgacccacta	tcaaagcctg	4500
cttctcacag	g agagggtcac	gttcgctcca	ccagccgctc	tcaaccctgo	cactcttctg	4560
cctgaagaga	a ctgatgaaco	agtgactcat	gattgccatc	aactattgat	tgaggagact	4620
ggggtccgca	a aggaccttac	agacatacco	ctgactggag	aagtgttaac	ctggttcact	4680
40gacggaagc	a gctatgtggt	ggaaggtaag	g aggatggctg	gggcggcggt	ggtggacggg	4740

PCT/US00/14296

	acccgcacga	tctgggccag	cagcctgccg	gaaggaactt	cagcacaaaa	ggctgagctc	4800
	atggccctca	cgcaagcttt	gcggctggcc	gaagggaaat	ccataaacat	ttatacagac	4860
	agcaggtatg	cctttgcgac	tgcacacgta	cacggggcca	tctataagca	aagggggttg	4920
	cttacctcag	cagggaggga	aataaagaac	aaagaggaaa	ttctaagcct	attagaagcc	4980
!	Sttacatttgc	caaaaaggct	agctattata	cactgtcctg	gacatcagaa	agccaaagat	5040
	cccatatcca	gagggaacca	gatggctgac	cgggttgcca	agcaggcagc	ccagggtgtt	5100
	aaccttctgc	ctatgataga	aacacccaaa	gccccagaac	ccggacgaca	gtacacccta	5160
	gaagactggc	aagagataaa	aaagatagac	cagttctctg	agactccgga	agggacctgc	5220
	tatacctcag	atgggaagga	aatcctgccc	cacaaagaag	ggttagaata	tgtccaacag	5280
1	Oatacatcgtc	taacccacct	aggaactaaa	cacctgcagc	agttggtcag	aacatctcct	5340
	tatcatgttc	tgaggctacc	aggagtggct	gattcggtgg	tcaaacactg	tgtgccctgc	5400
	cagctggtta	atgctaatcc	ttccagaata	cctccaggaa	agagactaag	gggaagccac	5460
	ccaggcgctc	actgggaagt	ggacttcact	gaggtaaagc	cggctaaata	cggaaacaaa	5520
	tatctattgg	tttttgtaga	caccttttca	ggatgggtag	aggcttatcc	tactaagaaa	5580
1	5gagacttcaa	ccgtggtggc	taagaaaata	ctggaggaaa	tttttccaag	atttggaata	5640
	cctaaggtaa	tagggtcaga	caatggtcca	gctttcgttg	cccaggtaag	tcagggactg	5700
	gccaagatat	tggggattga	ttggaaactg	cattgtgcat	acagacccca	aagctcagga	5760
	caggtagaga	ggatgaatag	aaccattaaa	gagaccctta	ctaaattgac	cgcggagact	5820
	ggcgttaatg	attggatagc	tctcctgccc	tttgtgcttt	ttagggttag	gaacacccct	5880
2	0ggacagtttg	ggctgacccc	ctatgaatta	ctctacgggg	gaccccccc	attggtagaa	5940
	attgcttccg	tacatagtgc	tgacgtgctg	ctttcccagc	ctttgttctc	taggctcaag	6000
	gcacttgagt	gggtgagaca	acgagcgtgg	aggcaactcc	gggaggccta	ctcaggagga	6060
	ggagacttgc	agatcc					6076

25

<210> 22

<211> 4918

<212> DNA

30<213> Porcine endogenous retrovirus

<400> 22

```
tggattttgc agactgtgga tcccggggtt tgcgacctta gcagccccac tctacccact
                                                                         60
 aaccaaagaa aaaggggaat tctcctgggc tcctgagcac cagaaggcat ttgatgctat
                                                                        120
35caaaaaggcc ctgctgagcg cacctgctct ggccctccct gacgtaacta aaccctttac
                                                                        180
 cctttatgtg gatgagcgta agggagtagc ccggggagtt ttaacccaaa ctctaggacc
                                                                        240
 atggaggaga cctgttgcct acctgtcaaa gaagctcgat cctgtagcca gtggttggcc
                                                                        300
 cgtatgcctg aaggctatcg cagctgtggc catactggtc aaggacgctg acaaattgac
                                                                        360
  tttgggacag aatataactg taatagcccc ccatgcgttg gagaacatcg ttcggcagcc
                                                                        420
                                                                         480
40cccagaccga tggatgacca acgcccgcat gacccactat caaagcctgc ttctcacaga
```

540	ctgaagagac	actcttctgc	caaccctgcc	cagccgctct	ttcgctccac	gagggtcacg
600	gggtccgcaa	gaggagactg	actattgatt	attgccatca	gtgactcatg	tgatgaacca
660	acggaagcag	tggttcactg	agtgttaacc	tgactggaga	gacataccgc	ggaccttaca
720	cccgcacgat	gtggacggga	ggcggcggtg	ggatggctgg	gaaggtaaga	ctatgtggtg
780	tggccctcac	gctgagctca	agcacaaaag	aaggaacttc	agcctgccgg	5ctgggccagc
840	gcaggtatgc	tatacagaca	cataaacatt	aagggaaatc	cggctggccg	gcaagctttg
900	ttacctcagc	agggggttgc	ctataagcaa	acggggccat	gcacacgtac	ctttgcgact
960	tacatttgcc	ttagaagcct	tctaagccta	aagaggaaat	ataaagaaca	agggagggaa
1020	ccatatccag	gccaaagatc	acatcagaaa	actgtcctgg	gctattatac	aaaaaggcta
1080	accttctgcc	cagggtgtta	gcaggcagcc	gggttgccaa	atggctgacc	10agggaaccag
1140	aagactggca	tacaccctag	cggacgacag	ccccagaacc	acacccaaag	tatgatagaa
1200	atacctcaga	gggacctgct	gactccggaa	agttctctga	aagatagacc	agagataaaa
1260	tacatcgtct	gtccaacaga	gttagaatat	acaaagaagg	atcctgcccc	tgggaaggaa
1320	atcatgttct	acatctcctt	gttggtcaga	acctgcagca	ggaactaaac	aacccaccta
1380	agctggttaa	gtgccctgcc	caaacactgt	attcggtggt	ggagtggctg	15gaggctacca
1440	caggcgctca	ggaagccacc	gagactaagg	ctccaggaaa	tccagaatac	tgctaatcct
1500	atctattggt	ggaaacaaat	ggctaaatac	aggtaaagcc	gacttcactg	ctgggaagtg
1560	agacttcaac	actaagaaag	ggcttatcct	gatgggtaga	accttttcag	ttttgtagac
1620	ctaaggtaat	tttggaatac	ttttccaaga	tggaggaaat	aagaaaatac	cgtggtggct
1680	ccaagatatt	cagggactgg	ccaggtaagt	ctttcgttgc	aatggtccag	20agggtcagac
1740	aggtagagag	agctcaggac	cagaccccaa	attgtgcata	tggaaactgc	ggggattgat
1800	gcgttaatga	gcggagactg	taaattgacc	agacccttac	accattaaag	gatgaataga
1860	gacagtttgg	aacacccctg	tagggttagg	ttgtgctttt	ctcctgccct	ttggatagct
1920	ttgcttccgt	ttggtagaaa	accccccca	tctacggggg	tatgaattac	gctgaccccc
1980	cacttgagtg	aggctcaagg	tttgttctct	tttcccagcc	gacgtgctgc	25acatagtgct
2040	gagacttgca	tcaggaggag	ggaggcctac	ggcaactccg	cgagcgtgga	ggtgagacaa
2100	gtg caggaa a	agacgccacc	agtctacgtt	tgggagattc	cgtttccaag	gatcccacat
2160	cggctgtgaa	accacaccaa	cgtacttttg	gcccttatca	cggtggaagg	cctcgagact
2220	cacctcccga	aagccggcgc	atcccacgtt	ggatccatgc	atctccacct	agtcgaagga
2280	atcgcgtggt	cttcgcctcc	tccccttaag	agactgaaaa	aaagccgaaa	30ttcggggtgg
2340	agcccgaact	ccttgtggac	atggtaaacg	cctcaagtta	gtcaataact	tccttactct
2400	attaatatta	cggtacaggt	ttactgactc	acctggttac	cttatctctc	cccataaacc
2460	gtctgccttc	tgaattatat	cctggtggcc	cccttgggga	aggggaggct	acagcactca
2520	cgtgcttacg	cgatgtactc	ccacaccccc	aatgaccagg	ccctggtctc	gatcagtaat
2580	cctcaggatt	ttgtggaaat	atgaagaata	ccccaaata	ttgcccagga	35ggttttacgt
2640	tggccagtct	gaattggaaa	ctaatgatgg	tgcgtaactt	gcaatggagc	tcttttgcaa
2700	aatcaattta	taccagttat	ttaacaatcc	tactcttttg	cagagtaagt	ctcagcaaga
2760	gtacgaaata	acaaaaagat	aacagcgggt	aaagattggc	tgggagatgg	attatggcca
2820	actgaaaaag	aataagtttc	attacttaaa	ttagacctag	ctgtcattcg	agcaaataag
2880	gtgtactatg	ttggggaata	atggtatgtc	aagtgggtaa	aaatattcaa	40gaaaacaaga

```
gaggetetgg gagaaagaaa ggatetgtte tgaetatteg eeteagaata gaaacteaga
                                                                        2940
 tggaacctcc ggttgctata ggaccaaata agggtttggc cgaacaagga cctccaatcc
                                                                        3000
 aagaacagag gccatctcct aacccctctg attacaatac aacctctgga tcagtcccca
                                                                        3060
                                                                        3120
 ctgagcctaa catcactatt aaaacagggg cgaaactttt taacctcatc cagggagctt
Sttcaagctct taactccacg actccagagg ctacctcttc ttgttggctt tgcttagctt
                                                                        3180
 cgggcccacc ttactatgag ggaatggcta gaggagggaa attcaatgtg acaaaggaac
                                                                        3240
 atagagacca atgtacatgg ggatcccaaa ataagcttac ccttactgag gtttctggaa
                                                                        3300
                                                                        3360
 aaggcacctg catagggatg gttcccccat cccaccaaca cctttgtaac cacactgaag
 cctttaatcg aacctctgag agtcagtatc tggtacctgg ttatgacagg tggtgggcat
                                                                        3420
10gtaatactgg attaacccct tgtgtttcca ccttggtttt caaccaaact aaagactttt
                                                                        3480
  gcgttatggt ccaaattgtc ccccgggtgt actactatcc cgaaaaagca gtccttgatg
                                                                        3540
 aatatgacta tagatataat cggccaaaaa gagagcccat atccctgaca ctagctgtaa
                                                                        3600
  tgctcggatt gggagtggct gcaggcgtgg gaacaggaac ggctgcccta atcacaggac
                                                                        3660
  cgcaacagct ggagaaagga cttagtaacc tacatcgaat tgtaacggaa gatctccaag
                                                                        3720
15ccctagaaaa atctgtcagt aacctggagg aatccctaac ctccttatct gaagtggttc
                                                                        3780
  tacagaacag aagggggtta gatctgttat ttctaaaaga aggaggatta tgtgtagcct
                                                                        3840
                                                                        3900
  tgaaggagga atgctgtttt tatgtggatc attcaggggc catcagagac tccatgaaca
  agcttagaga aaggttggag aagcgtcgaa gggaaaagga aactactcaa gggtggtttg
                                                                        3960
                                                                        4020
  agggatggtt caacaggtct ccttggttgg ctaccctact ttctgcttta acaggaccct
                                                                        4080
20taatagtcct cctcctgtta ctcacagttg ggccatgtat tattaacaag ttaattgcct
                                                                        4140
  tcattagaga acgaataagt gcagtccaga tcatggtact tagacaacag taccaaagcc
  cgtctagcag agaagctggc cgctagctct accagttcta agattagaac tattaacaag
                                                                        4200
  agaagaagtg gggaatgaaa ggatgaaaat gcaacctgac tctcccagaa cccaggaagt
                                                                        4260
  taataagaag ctctaaatgc cctcgaattc cagaccctgt tccctatagg taaaagatca
                                                                        4320
25tactttttgc tgttttaggg cttgctttct gctctgtaca aaactttgtg gaaggggaaa
                                                                        4380
  aacaggcccc tgagtatgtg cctctatgct tgaaacttct tgaaactgct cctaactgct
                                                                        4440
  tgtttggctt ctgtaaacct gcttgcataa gataaaaaga ggagaagtca attgcctaac
                                                                        4500
  ggaccccagt aagatcgggt gtaccacaaa atgttgaaac acatatcttg gtgacaacat
                                                                        4560
  gtctccccca ccccgaaaca tgcgcaaatg tgtaactcta aaacaattta aattaattgg
                                                                        4620
30tccacgaagc gcgggctctc gaagttttaa attgactggt ttgtgatatt ttgaaatgat
                                                                        4680
  tggtttgtaa agcgcgggct ttgttgtgaa ccccataaaa gctgtcccga ctccacactc
                                                                        4740
  ggggccgcag tcctctaccc ctgcgtggtg tacgactgtg ggccccagcg cgcttggaat
                                                                        4800
  aaaaatcctc ttgctgtttg catcaagacc gcttctcgtg agtgattaag gggagtcgcc
                                                                        4860
  ttttccgagc ctggaggttc tttttgctag tcttacagca cctttatttt ttccattt
                                                                        4918
```

35

<210> 23

<211> 7873

<212> DNA

<213> Porcine endogenous retrovirus

<400> 23 ggatctgttg gtttctgttt tgtgtgtctt tgtcttgtgc gtccttgtct acagttttaa 60 tatgggacag acggtgacga cccctcttag tttgactctc gaccattgga ctgaagttaa 120 180 atccagggct cataatttgt cagttcaggt taagaaggga ccttggcaga ctttctgtgt 240 Sctctgaatgg ccgacattcg atgttggatg gccatcagag gggaccttta attctgagat tatectgget gttaaageaa ttatttttea gaetggaeee ggeteteate eeaateagga 300 geoctatate ettaegtgge aagatttgge agaggateet eegecatggg ttaaaeettg 360 420 gctgaataag ccaagaaagc caggtccccg aattctggct cttggagaga aaaacaaaca ctcggctgaa aaagtcaagc cctctcctca tatctacccc gagattgagg agccgccggc 480 10ttggccggaa ccccaatctg ttcccccacc cccttatctg gcacagggtg ctgcgagggg 540 accetetgee cetectggag etceggeggt ggagggaeet getgeaggga eteggageeg 600 gaggggcgcc accccggagc ggacagacga gatcgcgaca ttaccgctgc gcacgtacgg 660 720 ccctcccaca ccggggggcc aattgcagcc cctccagtat tggccctttt cttctgcaga 780 tetetataat tggaaaacta accateceee ttteteggag gateeecaae geeteaeggg 840 15gttggtggag tcccttatgt tctctcacca gcctacttgg gatgattgtc aacagctgct 900 gcagacactc ttcacaaccg aggagcgaga gagaattctg ttagaggcta gaaaaaatgt tectggggee gaegggegae ceaegeagtt geaaaatgag attgaeatgg gattteeett 960 1020 gactegeece ggttgggaet acaacaegge tgaaggtagg gagagettga aaatetateg 1080 ccaggetetg gtggegggte teeggggege etcaagaegg eecaetaatt tggetaaggt 1140 20aagagaagtg atgcagggac cgaatgaacc cccctctgtt ttccttgaga ggctcttgga agcettcagg cggtacaccc cttttgatcc cacctcagag gcccaaaaaag cctcagtggc 1200 tttggccttt ataggacagt cagccttgga tattagaaag aagcttcaga gactggaagg 1260 gttacaggag gctgagttac gtgatctagt aaaggaggca gagaaagtat attacaaaag 1320 ggagacagaa gaagaaaggg aacaaagaaa agagagaaa agagaggaaa gggaggaaag 1380 25acgtaataaa cggcaagaga agaatttgac taagatcttg gctgcagtgg ttgaagggaa 1440 1500 aagcaatacg gaaagagaga gagattttag gaaaattagg tcaggcccta gacagtcagg 1560 gaacctgggc aataggaccc cactcgacaa ggaccaatgt gcatattgta aagaaaaagg acactgggca aggaactgcc ccaagaaggg aaacaaagga ctgaaggtct tagctctgga 1620 agaagataaa gactagggaa gacggggttc ggaccccctc cccgagccca gggtaacttt 1680 1740 30gaaggtggag gggcaaccag ttgagtteet ggttgatace ggagegaaac atteagtget 1800 actacagcca ttaggaaaac taaaagataa aaaatcctgg gtgatgggtg ccacagggca 1860 acaacagtat ccatggacta cccgaagaac agttgacttg ggagtgggac gggtaaccca 1920 ctcgtttctg gtcatacctg agtgcccagc acccctctta ggtagagact tattgaccaa gatgggagca caaatttctt ttgaacaagg gaaaccagaa gtgtctgcaa ataacaaacc 1980 35tatcactgtg ttgaccctcc aattagatga cgaatatcga ctatactctc ccctagtaaa 2040 gcctgatcaa aatatacaat tctggttgga acagtttccc caagcctggg cagaaaccgc 2100 agggatgggt ttggcaaagc aagttccccc acaagttatt caactgaagg ccagtgccac 2160 2220 accagtgtca gtcagacagt acccettgag taaagaaget caagaaggaa tteggeegca 2280 tgtccaaaga ttaatccaac agggcatcct agttcctgtc caatctccct ggaatactcc 40cctgctaccg gttagaaagc ctgggactaa tgactatcga ccagtacagg acttgagaga 2340

ggtgaataaa	caaatacaaa	atatacaccc	aacagtcccg	aacccttata	acctcttgtg	2400
		gctggtatac				2460
		gccaaccact				2520
		cctggacccg				2580
5catctttgac						2640
		atgacctgct				2700
		tgctggaatt				2760
		gagaggtaac				2820
		ggaagaaaac				2880
10caaacaagtg						2940
		tctacccact				3000
		ttgatgctat				3060
		aaccctttac				3120
		ctctaggacc				3180
15gaagctcgat						3240
		acaaattgac				3300
		ttcggcagcc				3360
		ttctcacaga				3420
		ctgaagagac				3480
20actattgatt						3540
		acggaagcag				3600
		cccgcacgat				3660
		tggccctcac				3720
		gcaggtatgc				3780
25ctataagcaa						3840
		tacatttgcc				3900
-		ccatatccag				3960
		accttctgcc				4020
=		aagactggca				4080
30gactccggaa						4140
		tacatcgtct				4200
		atcatgttct				4260
		agctgggtaa				4320
		tcaggatggg				4380
35caaccgtggt						4440
		tccagctttc				4500
		actgcattgt				4560
		taaagagacc				4620
					ccctggacag	4680
40tttgggctga						4740

tccgtacata g	tgctgacgt	gctgctttcc (cagcctttgt	tctctaggct	caaggcactt	4800
gagtgggtga g						4860
ttgcagatcc C						4920
ggaaacctcg a						4980
5gtgaaagtcg a						5040
cccgattcgg g	atagaaagc	cgaaaagact	gaaaatcccc	ttaagcttcg	cctccatcgc	5100
gtggttcctt a						5160
gaactcccat a						5220
tattaacagc a						5280
10ccttcgatca g						5340
ttacgggttt t						5400
ggatttcttt t						5460
agtctctcag C						5520
atttaattat g						5580
15aaataagcaa a						5640
aaaaggaaaa						5700
ctatggaggc t						5760
tcagatggaa						5820
aatccaagaa						5880
20ccccactgag						5940
		ccacgactcc				6000
		atgagggaat				6060
ggaacataga	gaccaatgta	catggggatc	ccaaaataag	cttaccctta	ctgaggtttc	6120
tagaaaaaaa	acctocatao	ggatggttcc	cccatcccac	caacaccttt	gtaaccacac	6180
25tgaagccttt						6240
					aaactaaaga	6300
					aagcagtcct	6360
					tgacactagc	6420
tgtaatgctc	ggattgggag	tggctgcagg	cgtgggaaca	ggaacggctg	ccctaatcac	6480
30aggaccgcaa	cagctggaga	aaggacttag	taacctacat	cgaattgtaa	cggaagatct	6540
ccaacccta	gaaaaatcto	tcagtaacct	ggaggaatco	ctaacctcct	: tatctgaagt	6600
ggttgtagag	aacagaagg	ggttagatct	gttatttcta	aaagaaggag	g gattatgtgt	6660
ageettgaag	gaggaatgct	gtttttatgt	ggatcattca	ggggccatca	a gagactccat	6720
gaacaagett	agagaaagga	a ctggagagcg	cccgcgggt	tcgaacaac	cagacaggtt	6780
35gcttgtttca	attaaagaag	totcoaagta	accgctgago	taaagccag	ttagagaaag	6840
gttggagaag	catcaaaaa	aaaaggaaa	tactcaagag	tggtttgag	g gatggttcaa	6900
					a tagtcctcct	6960
					a ttagagaacg	7020
					t ctagcagaga	7080
					a agaagtgggg	7140
404901990090						

PCT/US00/14296

26

26	
aatgaaagga tgaaaatgca acctgactct cccagaaccc aggaagttaa taagaagctc	7200
taaatgccct cgaattccag accetgttcc ctataggtaa aagatcatac tttttgctgt	7260
tttagggett getttetget etgtacaaaa etttgtggaa ggggaaaaac aggeeeetga	7320
gtatgtgcct ctatgcttga aacttcttga aactgctcct aactgcttgt ttggcttctg	7380
Staaacctgct tgcataagat aaaaagagga gaagtcaatt gcctaacgga ccccagtaag	7440
atogggtgta coacaaaatg ttgaaacaca tatottggtg acaacatgto toccocacco	7500
cgaaacatgc gcaaatgtgt aactctaaaa caatttaaat taattggtcc acgaagcgcg	7560
ggctctcgaa gttttaaatt gactggtttg tgatattttg aaatgattgg tttgtaaagc	7620
gcgggctttg ttgtaaaccc cataaaagct gtcccgactc cacactcggg gccgcagtcc	7680
10tctacccctg cgtggtgtac gactgtgggc cccagcgcgc ttggaataaa aatcctcttg	7740
ctgtttgcat caagaccgct tctcgtgagt gattaagggg agtcgccttt tccgagcctg	7800
gaggttettt ttgetagtet tacaacagea eeteagtttt gtteetaaga agtetgegge	7860
cctcacccag tca	7873
15	
<210> 24	
<211> 20	
<212> DNA	
20<213> Rous sarcoma virus	
<400> 24	
gggacgaggt tatgccgctg	20
25<210> 25	
<211> 20	
<212> DNA	
<213> Rous sarcoma virus	
30<400> 25	20
gggcgtgcgc gcattaccac	20
<210> 26	
<211> 20	
35<212> DNA	
<213> Rous sarcoma virus	
<400> 26	20
gaccgaccca gggaacaatc	20

```
<210> 27
 <211> 20
 <212> DNA
 <213> Rous sarcoma virus
 <400> 27
                                                                            20
 atgaggaaaa ttgcgggtgg
 <210> 28
10<211> 20
  <212> DNA
  <213> Artificial Sequence
  <220>
15<223> A primer
  <400> 28
                                                                            20
  ggaatgtgac tggtaatgga
20<210> 29
  <211> 20
  <212> DNA
  <213> Artificial Sequence
25<220>
  <223> A primer
   <400> 29
                                                                             20
  gccttagtga tggtgatggt
 30
   <210> 30
   <211> 21
   <212> DNA
   <213> Mus musculus
 35
   <400> 30
                                                                             21
   ccatccgtct tcatcttccc t
   <210> 31
 40<211> 21
```

```
<212> DNA
 <213> Mus musculus
 <400> 31
                                                                          21
Stggtgcggtg tccttgtagt t
 <210> 32
 <211> 6076
 <212> DNA
10<213> Porcine endogenous retrovirus
  <400> 32
                                                                          60
 ggagtttgag ttttatcgaa tttgaaacag tggtttacat ggagattgta gtgaaaggat
  gaaaatgcaa cctgactctc ccagaaccca ggaagttaat aagaagctct aaatgccctc
                                                                         120
15gaattccaga ccctgttccc tataggtaaa agatcatact ttttgctgtt ttagggcttg
                                                                         180
                                                                         240
  ctttctgctc tgtacaaaac tttgtggaag gggaaaaaca ggcccctgag tatgtgcctc
                                                                         300
  tatgcttgaa acttcttgaa actgctccta actgcttgtt tggcttctgt aaacctgctt
  gcataagata aaaagaggag aagtcaattg cctaacggac cccagtaaga tcgggtgtac
                                                                         360
                                                                         420
  cacaaaatgt tgaaacacat atcttggtga caacatgtct cccccacccc gaaacatgcg
                                                                         480
20caaatgtgta actctaaaac aatttaaatt aattggtcca cgaagcgcgg gctctcgaag
  ttttaaattg actggtttgt gatattttga aatgattggt ttgtaaagcg cgggctttgt
                                                                         540
                                                                         600
  tgtgaacccc ataaaagctg tcccgactcc acactcgggg ccgcagtcct ctacccctgc
  gtggtgtacg actgtgggcc ccagcgcgct tggaataaaa atcctcttgc tgtttgcatc
                                                                         660
  aagaccgctt ctcgtgagtg attaagggga gtcgcctttt ccgagcctgg aggttctttt
                                                                         720
25tgctagtctt acatttgggg gctcgtccgg gatctgtcgc ggccacccct aacacccgag
                                                                         780
  aaccgacttg gaggtaaaaa ggatcctctt tttaacgtgt atgcatgtac cggccggcgt
                                                                         840
                                                                         900
  ctctgttctg agtgtctgtt ttcagtggtg cgcgctttcg gtttgcagct gtcctctcag
                                                                         960
  accgtaagga ctgggggact gtgatcagca gacgtgctag gaggatcaca ggctgccacc
  ctgggggacg ccccgggagg tggggagagc cagggacgcc tggtggtctc cttctgtcgg
                                                                         1020
                                                                         1080
30tcagaggacc gagttctgtt gttgaagcga aagcttcccc ctccgcggcc gtccgactct
                                                                         1140
  tttgcctgct tgtggaagac gcggacgggt cgcgtgtgtc tggatctgtt ggtttctgtt
                                                                         1200
  ttgtgtgtct ttgtcttgtg cgtccttgtc tacagtttta atatgggaca gacggtgacg
                                                                         1260
  acceptetta gittgactet egaceatigg actgaagita aatecaggge teataatitg
  tragttragg traagaaggg accttggrag actttrtgtg trtctgaatg grogarattr
                                                                         1320
                                                                         1380
 35gatgttggat ggccatcaga ggggaccttt aattctgaga ttatcctggc tgttaaagca
                                                                         1440
   attatttttc agactggacc cggctctcat cccaatcagg agccctatat ccttacgtgg
                                                                         1500
   caagatttgg cagaggatcc tccgccatgg gttaaacctt ggctgaataa gccaagaaag
                                                                         1560
   ccaggtcccc gaattctggc tcttggagag aaaaacaaac actcggctga aaaagtcaag
   ccctctcctc atatctaccc cgagattgag gagccgccgg cttggccgga accccaatct
                                                                         1620
 40gttcccccac ccccttatct ggcacagggt gctgcgaggg gaccctctgc ccctcctgga
                                                                         1680
```

gctccggcgg	tggagggacc	tgctgcaggg	actcggagcc	ggaggggcgc	caccccggag	1740
cggacagacg	agatcgcgac	attaccgctg	cgcacgtacg	gccctcccac	accggggggc	1800
caattgcagc	ccctccagta	ttggcccttt	tcttctgcag	atctctataa	ttggaaaact	1860
aaccatcccc	ctttctcgga	ggatccccaa	cgcctcacgg	ggttggtgga	gtcccttatg	1920
5ttctctcacc	agcctacttg	ggatgattgt	caacagctgc	tgcagacact	cttcacaacc	1980
gaggagcgag	agagaattct	gttagaggct	agaaaaaatg	ttcctggggc	cgacgggcga	2040
cccacgcagt	tgcaaaatga	gattgacatg	ggatttccct	tgactcgccc	cggttgggac	2100
tacaacacgg	ctgaaggtag	ggagagcttg	aaaatctatc	gccaggctct	ggtggcgggt	2160
ctccggggcg	cctcaagacg	gcccactaat	ttggctaagg	taagagaagt	gatgcaggga	2220
10ccgaatgaac	cccctctgt	tttccttgag	aggctcttgg	aagccttcag	gcggtacacc	2280
ccttttgatc	ccacctcaga	ggcccaaaaa	gcctcagtgg	ctttggcctt	tataggacag	2340
tcagccttgg	atattagaaa	gaagcttcag	agactggaag	ggttacagga	ggctgagtta	2400
cgtgatctag	tgaaggaggc	agagaaagta	tattacaaaa	gggagacaga	agaagaaagg	2460
gaacaaagaa	aagagagaga	aagagaggaa	agggaggaaa	gacgtaataa	acggcaagag	2520
15aagaatttga	ctaagatctt	ggctgcagtg	gttgaaggga	aaagcaatac	ggaaagagag	2580
a ga gatttta	ggaaaattag	gtcaggccct	agacagtcag	ggaacctggg	caataggacc	2640
ccactcgaca	aggaccaatg	tgcatattgt	aaagaaaaag	gacactgggc	aaggaactgc	2700
cccaagaagg	gaaacaaagg	actgaaggtc	ttagctctgg	aagaagataa	agactaggga	2760
agacggggtt	cggaccccct	ccccgagccc	agggtaactt	tgaaggtgga	ggggcaacca	2820
20gttgagttcc	tggttgatac	cggagcgaaa	cattcagtgc	tactacagcc	attaggaaaa	2880
ctaaaagata	aaaaatcctg	ggtgatgggt	gccacagggc	aacaacagta	tccatggact	2940
acccgaagaa	cagttgactt	gggagtggga	cgggtaaccc	actcgtttct	ggtcatacct	3000
gagtgcccag	cacccctctt	aggtagagac	ttattgacca	agatgggagc	acaaatttct	3060
tttgaacaag	ggaaaccaga	agtgtctgca	aataacaaac	ctatcactgt	gttgaccctc	3120
25caattagatg	acgaatatcg	actatactct	cccctagtaa	agcctgatca	aaatatacaa	3180
ttctggttgg	aacagtttcc	ccaagcctgg	gcagaaaccg	cagggatggg	tttggcaaag	3240
caagttcccc	cacaagttat	tcaactgaag	gccagtgcca	caccagtgtc	agtcagacag	3300
taccccttga	gtaaagaagc	tcaagaagga	attcggccgc	atgtccaaag	attaatccaa	3360
cagggcatcc	tagttcctgt	ccaatctccc	tggaatactc	ccctgctacc	ggttagaaag	3420
30cctgggacta	atgactatcg	accagtacag	gacttgagag	aggtcaataa	acgggtgcag	3480
gatatacacc	caacagtccc	gaacccttat	aacctcttgt	gtgctctccc	accccaacgg	3540
agctggtata	cagtattgga	cttaaaggat	gcctttttct	gcctgagatt	acaccccact	3600
agccaaccac	tttttgcctt	cgaatggaga	gatccaggta	cgggaagaac	cgggcagctc	3660
acctggaccc	gactgcccca	agggttcaag	aactccccga	ccatctttga	cgaagcccta	3720
35cacagagacc	tggccaactt	caggatccaa	caccctcagg	tgaccctcct	ccagtacgtg	3780
gatgacctgc	ttctggcggg	agccaccaaa	caggactgct	tagaaggcac	gaaggcacta	3840
ctgctggaat	tgtctgacct	aggctacaga	gcctctgcta	agaaggccca	gatttgcagg	3900
agagaggtaa	catacttggg	gtacagtttg	cgggacgggc	agcgatggct	gacggaggca	3960
cggaagaaaa	ctgtagtcca	gataccggcc	ccaaccacag	ccaaacaagt	gagagaattt	4020
40ttggggacag	ctggattttg	cagactgtgg	atcccggggt	ttgcgacctt	agcagcccca	4080
	-					

PCT/US00/14296

ctctaccca	taaccaaaga	aaaaggggaa	ttctcctggg	ctcctgagca	ccagaaggca	4140
tttgatgcta	tcaaaaaggc	cctgctgagc	gcacctgctc	tggccctccc	tgacgtaact	4200
aaacccttta	ccctttatgt	ggatgagcgt	aagggagtag	cccggggagt	tttaacccaa	4260
actctagga	catggaggag	acctgttgcc	tacctgtcaa	agaagctcga	tcctgtagcc	4320
5agtggttgg	ccgtatgcct	gaaggctatc	gcagctgtgg	ccatactggt	caaggacgct	4380
gacaaattg	a ctttgggaca	gaatataact	gtaatagccc	cccatgcgtt	ggagaacatc	4440
gttcggcag	ccccagaccg	atggatgacc	aacgcccgca	tgacccacta	tcaaagcctg	4500
cttctcaca	g agagggtcac	gttcgctcca	ccagccgctc	tcaaccctgc	cactcttctg	4560
cctgaagag	a ctgatgaacc	agtgactcat	gattgccatc	aactattgat	tgaggagact	4620
10ggggtccgc	a aggaccttac	agacataccg	ctgactggag	aagtgttaac	ctggttcact	4680
gacggaagc	a gctatgtggt	ggaaggtaag	aggatggctg	gggcggcggt	ggtggacggg	4740
acccgcacg	a tctgggccag	cagcctgccg	gaaggaactt	cagcacaaaa	ggctgagctc	4800
atggccctc	a cgcaagcttt	gcggctggcc	gaagggaaat	ccataaacat	ttatacagac	4860
agcaggtat	g cctttgcgac	tgcacacgta	cacggggcca	tctataagca	aagggggttg	4920
15cttacctca	g cagggaggga	aataaagaac	aaagaggaaa	ttctaagcct	attagaagcc	4980
ttacatttg	c caaaaaggct	agctattata	cactgtcctg	gacatcagaa	agccaaagat	5040
cccatatcc	a gagggaacca	gatggctgac	cgggttgcca	agcaggcagc	ccagggtgtt	5100
aaccttctg	c ctatgataga	aacacccaaa	gccccagaac	ccggacgaca	gtacacccta	5160
gaagactgg	c aagagataaa	aaagatagac	cagttctctg	agactccgga	agggacctgc	5220
20tatacctca	g atgggaagga	aatcctgccc	cacaaagaag	ggttagaata	tgtccaacag	5280
atacatcgt	c taacccacct	aggaactaaa	cacctgcagc	agttggtcag	aacatctcct	5340
tatcatgtt	c tgaggctacc	aggagtggct	gattcggtgg	tcaaacactg	tgtgccctgc	5400
cagctggtt	a atgctaatco	ttccagaata	cctccaggaa	agagactaag	gggaagccac	5460
ccaggcgct	c actgggaagt	ggacttcact	gaggtaaagc	cggctaaata	cggaaacaaa	5520
25tatctattg	g tttttgtaga	caccttttca	ggatgggtag	aggcttatcc	tactaagaaa	5580
gagacttca	a ccgtggtggc	taagaaaata	ctggaggaaa	tttttccaag	attt ggaa ta	5640
cctaaggta	a tagggtcaga	caatggtcca	gctttcgttg	cccaggtaag	tcagggactg	5700
	t tggggattaa					5760
caggtagag	a ggatgaatag	, aaccattaaa	gagaccctta	ctaaattgac	cgcggagact	5820
	g attggatago					5880
_	g ggctgacccc					5940
	g tacatagtgo					6000
gcacttgag	t gggtgagaca	acgagcgtgg	aggcaactcc	gggaggccta	ctcaggagga	6060
ggagacttg	c agatcc					6076

INTERNATIONAL SEARCH REPORT

Inte .ional Application No
PCT/US 00/14296

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N15/48 C12N15/62 C07K19/00 A61K48/00 C07K14/15 C12N7/01 C1201/68C07K16/10 According to International Patent Classification (IPC) or to both national classification and IPC B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) C12N C07K A61K C12Q IPC 7 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practical, search terms used) BIOSIS, CHEM ABS Data C. DOCUMENTS CONSIDERED TO BE RELEVANT Citation of document, with indication, where appropriate, of the relevant passages Relevant to claim No. 15-20 AKIYOSHI D. E. ET AL.: "Identification of χ a full-length cDNA for an endogenous retrovirus of miniature swine" JOURNAL OF VIROLOGY, vol. 72, no. 5, May 1998 (1998-05), pages 4503-4507, XP002126425 ISSN: 0022-538X 1-14, the whole document Υ 21-28, 30,32-39 -/--Further documents are listed in the continuation of box C. Patent family members are listed in annex. Special categories of cited documents: "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the *A* document defining the general state of the art which is not considered to be of particular relevance invention "E" earlier document but published on or after the international "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled citation or other special reason (as specified) document referring to an oral disclosure, use, exhibition or other means in the art. document published prior to the international filing date but later than the priority date claimed *&* document member of the same patent family Date of mailing of the international search report Date of the actual completion of the international search 04/10/2000 21 September 2000

Fax: (+31-70) 340-3016

5

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentiaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Authorized officer

Mand1, B

INTERNATIONAL SEARCH REPORT

in\ _ational Application No PCT/US 00/14296

Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
VANBROCKLIN M. ET AL.: "Expression of a murine leukemia virus Gag-Escherichia coli RNase HI fusion polyprotein significantly inhibits virus spread." JOURNAL OF VIROLOGY, vol. 71, no. 4, 1997, pages 3312-3318, XP002148090 ISSN: 0022-538X the whole document	1-14, 21-28, 32-39
WO 98 53104 A (MEDICAL RES COUNCIL ;STOYE JONATHAN PAUL (GB); WEISS ROBIN ANTHONY) 26 November 1998 (1998-11-26)	29,31
the whole document	30
NATSOULIS G. ET AL.: "NEW ANTIVIRAL STRATEGY USING CAPSID-NUCLEASE FUSION PROTEINS" NATURE, vol. 352, no. 6336, 15 August 1991 (1991-08-15), pages 632-635, XP000608400 ISSN: 0028-0836 the whole document	15-28
SUN LQ. ET AL.: "Ribozyme-mediated suppression of Moloney murine leukemia virus and human immunodeficiency virus type I replication in permissive cells." PROC. NATL. ACAD. SCI. USA, vol. 91, October 1994 (1994-10), pages 9715-9719, XP002148091 the whole document	13
WO 96 32494 A (UNIV ALABAMA RES FOUND) 17 October 1996 (1996-10-17) the whole document	15-28
PATIENCE C. ET AL.: "INFECTION OF HUMAN CELLS BY AN ENDOGENOUS RETROVIRUS OF PIGS" NATURE MEDICINE, vol. 3, no. 3, 1 March 1997 (1997-03-01), pages 282-286, XP002037074 ISSN: 1078-8956 the whole document	1-39
	RNase HI fusion polyprotein significantly inhibits virus spread." JOURNAL OF VIROLOGY, vol. 71, no. 4, 1997, pages 3312-3318, XP002148090 ISSN: 0022-538X the whole document WO 98 53104 A (MEDICAL RES COUNCIL;STOYE JONATHAN PAUL (GB); WEISS ROBIN ANTHONY) 26 November 1998 (1998-11-26) the whole document NATSOULIS G. ET AL.: "NEW ANTIVIRAL STRATEGY USING CAPSID-NUCLEASE FUSION PROTEINS" NATURE, vol. 352, no. 6336, 15 August 1991 (1991-08-15), pages 632-635, XP000608400 ISSN: 0028-0836 the whole document SUN LQ. ET AL.: "Ribozyme-mediated suppression of Moloney murine leukemia virus and human immunodeficiency virus type I replication in permissive cells." PROC. NATL. ACAD. SCI. USA, vol. 91, October 1994 (1994-10), pages 9715-9719, XP002148091 the whole document WO 96 32494 A (UNIV ALABAMA RES FOUND) 17 October 1996 (1996-10-17) the whole document PATIENCE C. ET AL.: "INFECTION OF HUMAN CELLS BY AN ENDOGENOUS RETROVIRUS OF PIGS" NATURE MEDICINE, vol. 3, no. 3, 1 March 1997 (1997-03-01), pages 282-286, XP002037074 ISSN: 1078-8956

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-28,32-39

A method to inhibit or prevent infectious agent transmission in a mammalian transplant recipient; an isolated and purified nucleic acid molecule comprising at least a portion of a pig endogenous retrovirus, the corresponding polypeptides and methods of using them; a method to detect human tropic pig endogenous retroviruses; a fusion protein comprising a capsid or envelope protein of PERV and a degradative enzyme, the corresponding nucleic acid and a host cell comprising it.

2. Claims: 29-31

An antibody that specifically bings pig endogenous retrovirus.

INTERNATIONAL SEARCH REPORT

	Infort	PCT/US 00/14296			
Patent document cited in search report		Publication date		atent family nember(s)	Publication date
WO 9853104	A	26-11-1998	AU EP GB US	7442698 A 0981651 A 2333102 A 6100034 A	11-12-1998 01-03-2000 14-07-1999 08-08-2000
W0 9632494	A	17-10-1996	AU CA EP US	5482896 A 2230925 A 0873415 A 6001985 A	30-10-1996 17-10-1996 28-10-1998 14-12-1999

Jonal Application No.

Into